

# Homework 3

Luke Simon

2026-05-25

## Table of contents

|                                           |           |
|-------------------------------------------|-----------|
| <b>Problem 1. Giant kelp fronds</b>       | <b>2</b>  |
| a. . . . .                                | 2         |
| b. . . . .                                | 3         |
| c. . . . .                                | 4         |
| Assumption check . . . . .                | 4         |
| Running the test . . . . .                | 5         |
| d. . . . .                                | 6         |
| e. . . . .                                | 6         |
| f. . . . .                                | 6         |
| <b>Problem 2. Personal data</b>           | <b>7</b>  |
| a. . . . .                                | 7         |
| Categorical vs Response . . . . .         | 7         |
| Predictor vs Reponse . . . . .            | 8         |
| <b>Problem 3. Affective visualization</b> | <b>12</b> |
| a. . . . .                                | 12        |
| b. . . . .                                | 13        |
| c. . . . .                                | 13        |
| d. . . . .                                | 14        |
| e. . . . .                                | 14        |
| <b>Problem 4. Statistical critique</b>    | <b>14</b> |
| a. . . . .                                | 14        |
| b. . . . .                                | 15        |
| c. . . . .                                | 15        |
| d. . . . .                                | 15        |

```

# read in packages
library(tidyverse)
library(here)
library(janitor)
library(readxl)
library(patchwork)
library(ggeffects)

# read in data and create new object
kelp <- read_csv(
  here("data", "temp-kelp.csv"))

# creating new data frame with personal data
personal_data <- read_csv(here("data", "personal-datasheet.csv")) |>
  # standardizing column names
  clean_names() |>
  # dropping rows with na observations
  drop_na() |>
  # changing the stop column to display TRUE/FALSE instead of 1/0
  mutate(stop = as.logical(stop))

# creating new data frame for model predictions
personal_num <- personal_data |>
  # changing departure time column to numeric value
  mutate(departure_time = as.numeric(departure_time)) |>
  # selecting needed columns for model
  select(departure_time, walk_minutes)

```


## Problem 1. Giant kelp fronds

a.

There are two appropriate tests for testing the strength of the relationship between temperature and giant kelp frond elongation rate (correlation). These two are the Pearson's correlation, which assumes normally distributed continuous data, and Spearman's rank correlation, which does not require normality but does require continuous data. Pearson's correlation is the correct test to choose when there are no outliers in the data and the distribution is clearly normal, in all other cases Spearman's is more appropriate.

b.

```
# base layer: ggplot, data from kelp data frame
ggplot(data = kelp,
       mapping = aes(
         x = temp_c, # x-axis: temperature
         y = kelp_elong, # y-axis: elongation rate
         color = temp_c)) + # color gradient by temperature
# first layer: scatter plot
geom_point() +
# labeling axes and adding title
labs(x = expression(Temperature~(degree*C)),
     y = expression("Giant kelp elongation rate (cm day"^-1~")"),
     title = "As temperature increases, elongation rate decreases") +
# changing theme
theme_bw() +
# removing legend
theme(legend.position = "none")
```



C.


### Assumption check

Relationship appears linear, likely meaning Pearson's is the accurate test to run. Going to check normality of each variable, as each is continuous, to confirm that Pearson's is the correct test.

```
# creating object for temperature QQ plot
tempQQ <-
# baselayer: ggplot
ggplot(data = kelp, # using kelp data frame
       aes(sample = temp_c)) + # y-axis for QQ plot
# first layer: QQ reference line
geom_qq_line(color = "cornflowerblue") + # displaying line in blue
# second layer: QQ plot
geom_qq() +
# labeling title
labs(title = "Temperature QQ")+
# changing theme
theme_bw()

# creating object for elongation QQ plot
elongQQ <-
# baselayer: ggplot
ggplot(data = kelp, # using kelp data frame
       aes(sample = kelp_elong)) + # y-axis for QQ plot
# first layer: QQ reference line
geom_qq_line(color = "red") + # displaying line in red
# second layer: QQ plot
geom_qq() +
# labeling title
labs(title = "Kelp elongation QQ") +
# changing theme
theme_bw()

# using patchwork package to display plots side-by-side
tempQQ | elongQQ
```



To figure out which test was accurate to run in this scenario I first checked the linearity of the initial plot, there appears to be a negative linear relationship between temperature and giant kelp elongation rate. Then, to check the normality (I am assuming this data is independent here and we have established that they both are continuous) I created QQ plots of each variable. The QQ plots suggest both variables (and therefore all the data) are normally distributed (centering around the reference line with only slight deviations at the tails) and therefore suggests that a Pearson's correlation test is the appropriate test to perform.

### Running the test

```
# running correlation test
cor.test(kelp$temp_c, kelp$kelp_elong,
         method = "pearson") # pearson's test
```

Pearson's product-moment correlation

```
data: kelp$temp_c and kelp$kelp_elong
t = -5.189, df = 30, p-value = 1.366e-05
alternative hypothesis: true correlation is not equal to 0
95 percent confidence interval:
```

```
-0.8359665 -0.4460157
sample estimates:
      cor
-0.6877489
```

**d.**

To evaluate the strength of the relationship between temperature and giant kelp frond elongation rate, I used a Pearson's correlation test. I found a strong negative relationship between temperature and elongation rate (Pearson's  $r = -0.69$ ,  $t(30) = -5.2$ ,  $p < 0.001$ ,  $\alpha = 0.05$ )

**e.**

Through a Pearson's correlation test I was able to determine that there is a clear negative relationship between ocean temperature and giant kelp elongation rate in the sample area. Essentially, this means that within the sample zone in areas with a higher sea temperature we see a distinctly lower elongation rate for giant kelp fronds. This is also true in the reverse, meaning that in the areas sampled with a lower temperature the elongation rates of the kelp was seen to be higher.

**f.**

```
# running correlation test
cor.test(kelp$temp_c, kelp$kelp_elong,
         method = "spearman", exact = FALSE) # spearman's rank test
```

Spearman's rank correlation rho

```
data: kelp$temp_c and kelp$kelp_elong
S = 9216.1, p-value = 1.29e-05
alternative hypothesis: true rho is not equal to 0
sample estimates:
      rho
-0.6891684
```

If I had used a Spearman's rank correlation test initially, as opposed to the Pearson's correlation I had chosen, I would have been led to the same decision in regards to the relationship of the two variables and I would have interpreted the results the same. This makes sense as a Spearman

test has the same requirements for use as the Pearson but the Pearson is a parametric test, essentially, because we can use a Pearson's test we can also use a Spearman but the Pearson's will provide more information in the test that can be useful.

## Problem 2. Personal data


a.

### Categorical vs Response

```
# base layer: gg plot
ggplot(data = personal_data, # pulling from personal data frame
       aes(x = location, # x-axis
           y = walk_minutes, # y-axis
           color = location)) + # coloring by location
# first layer: box plot
geom_boxplot(color = "grey30", # changing box plot outline color
             linewidth = 0.5,) + # making border lines thicker
# second layer: jitter plot
geom_jitter(height = 0, # removing vertical jitter
            width = 0.1, # slight horizontal jitter
            alpha = 0.9, # making points slightly opaque
            size = 2) + # making point size larger
# changing theme
theme_bw() +
# removing legend
theme(legend.position = "none",
      panel.grid = element_blank()) +
# labeling x- & y-axes and adding title
labs(x = "Location",
     y = "Walk time (minutes)",
     title = "Box plots of total walk time separated by location",
     subtitle = "Most recent observation 2026-05-26") +
# changing colors based on location
scale_color_manual(values = c("Campbell" = "aquamarine3",
                              "387" = "orange"))
```

## Box plots of total walk time separated by location

Most recent observation 2026-05-26







## Predictor vs Reponse

### Model Fitting

```
personalmodel <- lm(  
  walk_minutes ~ departure_time,  
  data = personal_num  
)  
par(mfrow = c(2,2))  
plot(personalmodel)
```





- residuals normally distributed (QQ)
- residuals are homoscedastic (Residuals vs fitted/scale-location)
- no outliers influencing model coeff estimates (leverage)

```
summary(personalmodel)
```

Call:

```
lm(formula = walk_minutes ~ departure_time, data = personal_num)
```

Residuals:

| Min     | 1Q      | Median  | 3Q     | Max    |
|---------|---------|---------|--------|--------|
| -3.6270 | -0.4453 | -0.0661 | 0.6986 | 3.9486 |

Coefficients:

|                | Estimate   | Std. Error | t value | Pr(> t )     |
|----------------|------------|------------|---------|--------------|
| (Intercept)    | 279.271170 | 40.049689  | 6.973   | 3.28e-07 *** |
| departure_time | -0.009594  | 0.001457   | -6.585  | 8.23e-07 *** |

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.572 on 24 degrees of freedom

Multiple R-squared: 0.6437, Adjusted R-squared: 0.6289

F-statistic: 43.37 on 1 and 24 DF, p-value: 8.229e-07

```
# creating new object for predictions
preds <- ggpredict(
  personalmodel, # model object
  terms = "departure_time" # predictor
)

# displaying prediction
preds
```

# Predicted values of walk\_minutes


| departure_time | Predicted | 95% CI       |
|----------------|-----------|--------------|
| 26940          | 20.81     | 19.04, 22.58 |
| 27120          | 19.08     | 17.80, 20.36 |
| 27300          | 17.35     | 16.50, 18.21 |
| 27360          | 16.78     | 16.03, 17.52 |
| 27420          | 16.20     | 15.53, 16.87 |
| 27540          | 15.05     | 14.40, 15.71 |
| 27600          | 14.48     | 13.76, 15.19 |
| 27900          | 11.60     | 10.21, 12.99 |

**Figure**

```

# base layer: ggplot
ggplot(data = personal_data,
       aes(x = departure_time, # x-axis
           y = walk_minutes, # y-axis
           color = location, # coloring by location
           shape = stop)) + # changing shape based on stops
# first layer: points
geom_point(size = 2) + # making points larger
# second layer: ribbon representing confidence interval
geom_ribbon(data = preds,
           aes(x = x,
               y = predicted,
               ymin = conf.low,
               ymax = conf.high),
           alpha = 0.1,
           inherit.aes = FALSE) +
# third layer: line representing model predictions
geom_line(data = preds,
          aes(x = x,
              y = predicted),
          inherit.aes = FALSE) +
# changing theme
theme_bw() +
# removing legend
theme(legend.position = "none",
      panel.grid = element_blank()) +
# labeling x- & y-axes and adding title
labs(x = "Departure time",
     y = "Walk time (minutes)",
     title = "Negative relationship between walk time and departure time",
     subtitle = "Most recent observation 2026-05-26") +
# changing colors based on location
scale_color_manual(values = c("Campbell" = "aquamarine3",
                              "387" = "orange"))

```



b.

**Figure 1. Median walk time is identical across locations.** Points represent individual daily walk times to two different locations, Building 387 ( $n = 11$ ) and Campbell Hall ( $n = 15$ ), on UCSB's campus from Isla Vista. Boxplots display median walk time value for each location along with interquartile range and total spread. Location groupings are separated by color (387: orange, Campbell: aquamarine). Data collected by Luke Simon between the dates of 2026-04-20 and 2026-05-26.

**Figure 2. Negative relationship between departure time and walk time.** Points represent individual daily walk times to two different locations, Building 387 ( $n = 11$ ) and Campbell Hall ( $n = 15$ ), on UCSB's campus from Isla Vista. Point shape represents if a stop was taken on the walk (triangle = TRUE, circle = FALSE), while point color represents location (387: orange, Campbell: aquamarine). Line represents linear model of departure time vs walk time. Ribbon represents 95% confidence interval from predictions. Data collected by Luke Simon between the dates of 2026-04-20 and 2026-05-26.

### Problem 3. Affective visualization

a.


For my affective data visualization I want to create a network map of all of my data as nodes. These nodes would vary in shape and color which would each correspond to some detail from

the data set. My current thought process, though this will likely change, is that color would operate on a gradient for walk time and node shape would correspond to arrival time (number of sides = minutes after 7:50). I would also include some signifier for if I took a stop and connect each node by location. Node location relative to each other is based on temperature (x-axis) and wake up time (y-axis) and node border color signifies location.

**b.**

**c.**

- This draft only includes the first 12 observations



d.

This piece is a network display of my daily walk time to my 8 am class, heavily influenced by the creation of real network maps I have made for my job and shape and color work of Posavec and Lupi. This was created on the free and open source network mapping platform draw.io. Paths are created to represent which of the two buildings I was going to (in chronological order), colors follow rainbow order (red-purple) for walk time (red = 12, purple = 21) and the number of sides of each shape represents how many minutes after 7:50 I arrived, horizontal location is by daily temperature, while vertical is by wake-up time.


e.

Slideshow link

## Problem 4. Statistical critique

a.

The author is trying to answer the question of whether or not competition between herbaceous and woody plants depends on the availability of anabolic nutrients relative to catabolic nutrients. To do so, the selected test was the Kruskal-Wallis test where the response variable is either tree height or tree abundance and the predictor is treatment type. They are doing this so that they can analyze if there is a difference in tree height/abundance across treatment type.



**b.**

This plot is fairly visually clear and concise. The axes both make sense, though perhaps under the x-axis they could label with “Treatment” but this is fairly clear in context of the paper. Median values for each treatment type are clearly visible and easy to relate to the y-axis.

**c.**

I think that the authors handle visual clutter very well. This is going to be a cramped figure no matter what due to the sheer number of treatment types. I think the line widths for each element are thoughtfully chosen to make them distinct yet not too close together. The figure can appear to be a lot at first but it is quite clear once some time is spent with it. The data:ink ratio isn't great, but there is not much that can be done to fix this due to the large spread of the treatments.

**d.**

The change I would make would be to replace the SP and AS with their full name SP -> superphosphate and AS -> ammonium sulphate. I also think including treatment on the x-axis would be very helpful for making the figure stand alone. To help with clutter caused by expanding the names I think you could do away with the repeat of the treatment types for the second plot and utilize one concise x-axis. If the authors wanted to split this into two figures they could also include the underlying data, but I think this would be too crowded in its current side-by-side configuration.