

# Modelling in public health

SE Scientific Communication, 2019

Lukas Richter

28. June 2019

1 Background

2 Statistical models

3 Dynamic models

4 Conclusion

# Background

Infectious diseases are a threat to human health  
cost lives and also money

Outbreaks

“sporadic” cases

different transmission chains



# Outbreaks



## Influenza pandemic “swine flu”

Global, 2009-2010, 100.000–400.000 deaths

## Ebola

West Africa, 2014-2016, 11.000 deaths

DRC, since 2018, 1.900 deaths

## Zika Virus

Brazil, 2015-2016, ca. 215.000 cases (CHECK!) worldwide

## Measles

Europe, 2019, ca. 6.300 cases

Austria, 2019, more than 120 cases

## Objective/Goal

during outbreak:

- exploit all data

- inform response team in real time

non outbreak situations:

- health programmes are usually implemented over a long period of time

- with broad benefits to many in the community

- find high impact and cost-effective interventions

- allow evidence based decisions

- policy evaluation (vaccine programmes, WHO disease elimination targets)

benefits of model:

- cheap: Clinical trials are seldom large enough to capture these effects

- often little or no data to analyse (new emerging diseases)

- mathematical models can help to assess potential threats and impacts

- early in the process, and later aid in interpreting data from complex and multifactorial systems

# Types of models

dynamic, mathematical (SIR)


statistical (e.g. Poisson regression)

Bayesian statistics

spacial stats/models


-> visualise outcome / communication

# Example of outbreak analytics workflow.



# Intervention effect - Invasive Pneumococcal Disease (IPD)

**Figure:** Monthly incidence of (A) PCV10 ST-IPD and (B) non-PCV10 ex ST 6A-/19A-IPD, among the  $\geq 50$  years old, observed and modelled by a segmented negative binominal regression, Austria, January 2009-February 2017, shown are overall and seasonal trends.





## Serfling-like Model


$$\begin{aligned}\log(Y_t) = & \log(pop_t) + \beta_0 + \beta_1 t + \beta_2 \sin\left(\frac{2\pi t}{12}\right) \\ & + \beta_3 \cos\left(\frac{2\pi t}{12}\right) + \beta_5 (t - t_0)^+ \\ & + \mathbb{1}_{t-t_0>0} \left[ \beta_4 + \beta_6 \sin\left(\frac{2\pi t}{12}\right) + \beta_7 \cos\left(\frac{2\pi t}{12}\right) \right]\end{aligned}$$

with

$$(x)^+ = \begin{cases} x, & \text{if } x > 0, \\ 0, & \text{otherwise.} \end{cases}$$

# Transmission model of Zika Virus

| Vars  | Description  |
|-------|--|
| $S_h$ | Susceptible Humans                                       |
| $I_h$ | Infected/Infectious humans                               |
| $R_h$ | Humans recovered from infection (with lifelong immunity) |
| $S_v$ | Susceptible vectors                                      |
| $E_v$ | Exposed vectors  |



adapted from  
<https://www.reconlearn.org/> and Ferguson et al., 2016

# Transmission model of Zika Virus

## Humans/Host

$$\frac{dS_h}{dt} = \mu_h N_h - \frac{\beta_h b}{N_h} S_h I_v - \mu_h S_h$$


$$\frac{dI_h}{dt} = \frac{\beta_h b}{N_h} S_h I_v - (\gamma_h + \mu_h) I_h$$

$$\frac{dR_h}{dt} = \gamma_h I_h - \mu_h I_h$$

## Vectors

$$\frac{dS_v}{dt} = \mu_v N_v - \frac{\beta_v b}{N_h} I_h S_v - \mu_v S_v$$

$$\frac{dI_v}{dt} = \frac{\beta_v b}{N_h} I_h S_v - \mu_v I_v$$



adapted from  
<https://www.reconlearn.org/> and Ferguson et al., 2016

## other applications

GO

Ebola

Influenza (Nielsen) Nielsen et al., 2019

Foodborne outbreaks to identify the source of infection (poisson model)

## Conclusion/Wrap up

We saw some examples of applied modelling  
plays an increasingly important role in helping to guide the most high  
impact and cost-effective prevent disease  
models can be critical tools in guiding public health action.  
always comes with limitations (as other studies)  
decision makers benefit, so do the affected people still a lot to do -  
However, there are a number of challenges in achieving a successful  
interface between modelling and public health.



# Any questions?

# References

- [1] Lukas Richter et al. "Invasive Pneumococcal Diseases in Children and Adults before and after Introduction of the 10-Valent Pneumococcal Conjugate Vaccine into the Austrian National Immunization Program". In: *PloS One* 14.1 (2019), e0210081. ISSN: 1932-6203. DOI: 10.1371/journal.pone.0210081. pmid: 30629620.
- [2] Neil M. Ferguson et al. "Countering the Zika Epidemic in Latin America". In: *Science* 353.6297 (July 22, 2016), pp. 353–354. ISSN: 0036-8075, 1095-9203. DOI: 10.1126/science.aag0219. pmid: 27417493. URL: <https://science.sciencemag.org/content/353/6297/353> (visited on 06/19/2019).
- [3] J. Nielsen et al. "European All-Cause Excess and Influenza-Attributable Mortality in the 2017/18 Season: Should the Burden of Influenza B Be Reconsidered?". In: *Clinical Microbiology and Infection* (Feb. 18, 2019). ISSN: 1198-743X. DOI: 10.1016/j.cmi.2019.02.011. URL: <http://www.sciencedirect.com/science/article/pii/S1198743X19300588> (visited on 04/08/2019).
- [4] Polonsky Jonathan A. et al. "Outbreak Analytics: A Developing Data Science for Informing the Response to Emerging Pathogens". In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 374.1776 (July 8, 2019), p. 20180276. DOI: 10.1098/rstb.2018.0276. URL: <https://royalsocietypublishing.org/doi/10.1098/rstb.2018.0276> (visited on 06/18/2019).
- [5] R. E. Serfling, I. L. Sherman, and W. J. Houseworth. "Excess Pneumonia-Influenza Mortality by Age and Sex in Three Major Influenza A2 Epidemics, United States, 1957-58, 1960 and 1963". In: *American Journal of Epidemiology* 86.2 (Sept. 1967), pp. 433–441. ISSN: 0002-9262. DOI: 10.1093/oxfordjournals.aje.a120753. pmid: 6058395.