Analysis 1: Mid-semester Examination (corrected)

Date: September 23, 2026 **Time:** 3 hours **Marks:** 100

This is a corrected version of the paper; changes are marked *Correction*. Part labels in Q. V are relabelled (a)–(c); the original printed [a], [a], [b].

Answer *any four* questions. Each question carries 25 marks.

Write the answer to each question on a separate set of sheets. Write your name and roll number on each set.

You may use definitions of limit of a sequence, limsup and liminf of a sequence; that there exists a subsequence converging to the limsup; ratio and root tests for convergence of a series; definitions of continuity and uniform continuity; results related to sequences; continuity of a function. In problems related to continuity/uniform continuity you can use results related to sequences after stating them precisely, not by their name. You can also use any part of any question asked in the mid-sem while answering any other part, even if you have not answered the part you are quoting. Anything else, you have to write a proof.

I. Let $(a_n : n \geq 1) \subseteq \mathbb{R}$ be a sequence and let $s_n = \sum_{k=1}^{n} a_k$.

(a) If $(s_n : n \geq 1)$ converges to $s \in \mathbb{R}$, show that $(a_n : n \geq 1)$ converges to zero. [5]

(b) If $a_n = (-1)^n b_n$, where b_n satisfies $0 \leq b_{n+1} \leq b_n$ for all $n \geq 1$ and $b_n \to 0$, show that s_n converges. [8]

Correction: the paper does not require $b_n \to 0$, and without it the statement is false: $b_n = 1$ for all n gives partial sums alternating between -1 and 0, which do not converge.

(c) Let $0 \leq a_{n+1} \leq a_n$ for all n. For $m \geq 1$ let $b_m = 2^m a_{2^m}$ (thus $b_3 = 8a_8$). Show that the series $\sum_m b_m$ is convergent if and only if the series $\sum_n a_n$ is convergent. [12]

II. Let $(a_n : n \geq 1) \subseteq \mathbb{R}$ be a sequence such that [25]

(a) for any subsequence $(a_{n_k} : k \geq 1)$ of $(a_n : n \geq 1)$, writing $b_k = a_{n_k}$, there is a subsequence $(b_{k_j} : j \geq 1)$ of $(b_k : k \geq 1)$ such that $c_j = b_{k_j}$ converges;

(b) there is $\alpha \in \mathbb{R}$ such that for all subsequences $(a_{n_k} : k \geq 1)$ of $(a_n : n \geq 1)$ that converge, $(a_{n_k} : k \geq 1)$ converges to α.

Show that $(a_n : n \geq 1)$ converges to α.

III. Let $f : (0, 1) \to \mathbb{R}$ be a function.

(a) If f is continuous, show that for any sequence $(a_n : n \geq 1) \subseteq (0, 1)$ converging to $a \in (0, 1)$, the sequence $(f(a_n) : n \geq 1)$ converges to $f(a)$. [10]

(b) Suppose that for any sequence $(a_n : n \geq 1) \subseteq (0, 1)$ converging to $a \in (0, 1)$, the sequence $(f(a_n) : n \geq 1)$ converges to $f(a)$. Show that f is continuous. [15]

IV. Let $\{a_n : n \geq 0\}$, $\{b_n : n \geq 0\}$ be sequences such that $f(x) = \sum_{n=0}^{\infty} a_n x^n$ and $g(x) = \sum_{n=0}^{\infty} b_n x^n$ are convergent for all $x \in \mathbb{R}$.

(a) Show that f, g are continuous functions on $\mathbb{R}$. [6]

(b) Let $h(x) = f(x)g(x)$ for $x \in \mathbb{R}$. Show that h also admits a power series expansion $h(x) = \sum_{n=0}^{\infty} c_n x^n$ for $x \in \mathbb{R}$. Find an expression for c_n, $n \geq 0$, in terms of $\{a_n\}$ and $\{b_n\}$. [7]

Correction: the paper defines h only on $(-1, 1)$ but asks for an expansion valid on all of $\mathbb{R}$. Since f, g converge everywhere, h should be defined for $x \in \mathbb{R}$.

(c) Let $e(x) = 1 + \sum_{n=1}^{\infty} \frac{x^n}{n!}$ for $x \in \mathbb{R}$. Show that e is a continuous function and that $e(x + y) = e(x)e(y)$ for all $x, y \in \mathbb{R}$. [8]

(d) Show that $e(x) > 0$ for all $x \in \mathbb{R}$. [4]

V. Let $f, g: \mathbb{R} \to \mathbb{R}$ be continuous functions.

(a) Let $h(x) = f(x)g(x)$ for $x \in \mathbb{R}$. Show that h is a continuous function. [7]

(b) Show that for all $a, b \in \mathbb{R}$ with $a < b$, and $0 < \alpha < 1$, there is $c \in [a, b]$ such that $f(c) = \alpha f(a) + (1 - \alpha) f(b)$. [10]

Correction: the paper asks for $c \in (a, b)$, which is false when $f(a) = f(b)$: take $f(x) = (x - a)(x - b)$, so the target value is 0 but $f < 0$ on (a, b). Either allow $c \in [a, b]$ (as here) or assume $f(a) \neq f(b)$, in which case an open-interval c does exist.

(c) Suppose $\lim_{x \to \infty} f(x) = u \in \mathbb{R}$ and $\lim_{x \to -\infty} f(x) = v \in \mathbb{R}$. Show that f is bounded and also uniformly continuous on $\mathbb{R}$. [8]