
Sensitivity families at J = 56 (t(46)); bars = 95% MC intervals; size from 2,000 and power from 1,000 replications
a Size of true-null contrasts
b Power, J = 56, d = 0.3 unless stated
S1 between-R2 0 (within 0.3)
S1 between-R2 0.6 (within 0.3)
S2 total R2 0.1
S2 total R2 0.2
S3 bounded 0-100, ceiling, 5-pt steps
S4 MAR on pre-test
S4 MNAR on outcome
S4 attrition A/B/C 10/15/20%
S5 consent varies by cluster
S5 consent by arm and pre-test
S5 two clusters lost
S6 arm C residual SD x1.2
S8 offering shift 0.3
S9 crossed tutors (share 0.03)
S7 partial null (0.3, 0): C vs A
S8 offering effects +0.1/-0.1
S10 B leakage, (0.3, 0): C vs A
S7 (dB, dC) = (0.3, 0.15)
S7 (dB, dC) = (0.3, -0.3)
S7 (dB, dC) = (0.3, 0)
S8 offering effects 0.4/0.2
S10 covert AI in A: c 0.2, delta -0.3
S10 covert AI in A: c 0.2, delta 0.0
S10 covert AI in A: c 0.2, delta 0.3
S10 covert AI in A: c 0.4, delta -0.3
S10 covert AI in A: c 0.4, delta 0.0
S10 covert AI in A: c 0.4, delta 0.3
S10 B leakage 0.2, dC 0.3
S10 B leakage 0.2, dC 0.15
S10 B leakage 0.2, dC 0
0.02
0.04
0.06
0.08
0.2
0.4
0.6
0.8
1.0
Size of a true-null contrast (alpha 0.05)
Holm-adjusted power, B vs A
LMM t(J-c) (planned)
CR2 with Bell-McCaffrey df
ICC 0.05 (filled)
ICC 0.10 (open)