Title

R. Seto

Wed 12th Dec, 2012 12:17 Noon

Hydrodynamic force

One-body limit:

$$\mathbf{F}_i^{\text{self}} = -6\pi\mu a(\mathbf{v}_i - \mathbf{U}^{\infty}(\mathbf{r}_i)) \tag{1}$$

$$T_i^{\text{self}} = -8\pi\mu a^3(\boldsymbol{\omega}_i - \boldsymbol{\Omega}^{\infty}) \tag{2}$$

Melrose and Ball [2004] uses the leading squeeze mode of lubrication force:

$$\boldsymbol{F}_{ij}^{\text{pair}} = -\alpha_{\text{n}}(h_{ij}) \{ (\boldsymbol{v}_i - \boldsymbol{v}_j) \cdot \boldsymbol{n}_{ij} \} \boldsymbol{n}_{ij}$$
(3)

$$T_{ij}^{\text{pair}} = \mathbf{0},$$
 (4)

where

$$\alpha_{\rm n}(h_{ij}) = \frac{3\pi\mu a^2}{2h_{ij}} \tag{5}$$

Hydrodynamic interaction acting on particle i is given by

$$\boldsymbol{F}_{i}^{\mathrm{H}} = \boldsymbol{F}_{i}^{\mathrm{self}} + \sum_{j} \boldsymbol{F}_{ij}^{\mathrm{pair}}$$
 (6)

$$T_i^{\mathrm{H}} = T_i^{\mathrm{self}}$$
 (7)

Dimensionless equations

The unit of length, velocity and force are given: $L_0 = a$, $v_0 = a\dot{\gamma}$, and $F_0 \equiv 6\pi\mu av_0$. The dimensionless variables are introduced: $\tilde{h}_{ij} = h_{ij}/L_0$, $\tilde{\boldsymbol{v}}_i = \boldsymbol{v}_i/U_0$, and $\tilde{\boldsymbol{F}}_{ij}^{\text{pair}} = \boldsymbol{F}_{ij}^{\text{pair}}/F_0$. The relations are written as follows:

$$\tilde{\mathbf{F}}_i^{\text{self}} = -(\tilde{\mathbf{v}}_i - \tilde{\mathbf{U}}^{\infty}(\tilde{\mathbf{r}}_i)) \tag{8}$$

$$\tilde{\boldsymbol{F}}_{ij}^{\text{pair}} = -\frac{1}{4\tilde{h}_{ij}} \left\{ (\tilde{\boldsymbol{v}}_i - \tilde{\boldsymbol{v}}_i) \cdot \boldsymbol{n}_{ij} \right\} \boldsymbol{n}_{ij}$$
(9)

Matrix form

$$\boldsymbol{F}_{ij}^{\text{pair}} = -\frac{1}{4h_{ij}} \{ (\boldsymbol{v}_i - \boldsymbol{v}_j) \cdot \boldsymbol{n}_{ij} \} \boldsymbol{n}_{ij}$$
(10)

$$= -\frac{1}{4h_{ij}} \left[\left\{ (\boldsymbol{v}_i - \boldsymbol{U}_i^{\infty}) - (\boldsymbol{v}_j - \boldsymbol{U}_j^{\infty}) + \boldsymbol{U}_i^{\infty} - \boldsymbol{U}_j^{\infty} \right\} \cdot \boldsymbol{n}_{ij} \right] \boldsymbol{n}_{ij}$$
(11)

 $\Delta \boldsymbol{v} \equiv (\boldsymbol{v} - \boldsymbol{U}^{\infty})$

$$\Delta \boldsymbol{v} \cdot \boldsymbol{n} \boldsymbol{n} = (\Delta v_x n_x + \Delta v_y n_y + \Delta v_z n_z) n_x \boldsymbol{i}$$
(12)

$$+ (\Delta v_x n_x + \Delta v_u n_u + \Delta v_z n_z) n_u \mathbf{j} \tag{13}$$

$$+ (\Delta v_x n_x + \Delta v_y n_y + \Delta v_z n_z) n_z \mathbf{k} \tag{14}$$

$$= \begin{pmatrix} n_x n_x & n_y n_x & n_z n_x \\ n_x n_y & n_y n_y & n_z n_y \\ n_x n_z & n_y n_z & n_z n_z \end{pmatrix} \begin{pmatrix} \Delta v_x \\ \Delta v_y \\ \Delta v_z \end{pmatrix}$$
(15)

$$= \mathbf{n} \otimes \mathbf{n} \Delta \mathbf{v} \tag{16}$$

$$\boldsymbol{F}_{i} = -\Delta \boldsymbol{v}_{i} - \sum_{j} \left(\frac{1}{4h_{ij}} \boldsymbol{n}_{ij} \otimes \boldsymbol{n}_{ij} \Delta \boldsymbol{v}_{i} - \frac{1}{4h_{ij}} \boldsymbol{n}_{ij} \otimes \boldsymbol{n}_{ij} \Delta \boldsymbol{v}_{j} + \frac{1}{4h_{ij}} (\boldsymbol{U}_{i}^{\infty} - \boldsymbol{U}_{j}^{\infty}) \cdot \boldsymbol{n}_{ij} \boldsymbol{n}_{ij} \right)$$
(17)

$$\begin{pmatrix}
\vdots \\
\mathbf{F}_{i} \\
\vdots \\
\mathbf{F}_{j} \\
\vdots
\end{pmatrix} = \begin{pmatrix}
\vdots \\
\mathbf{F}_{i}^{\text{self}} + \mathbf{F}_{ij}^{\text{pair}} \\
\vdots \\
\mathbf{F}_{j}^{\text{self}} + \mathbf{F}_{ji}^{\text{pair}} \\
\vdots
\end{pmatrix}$$
(18)

$$= -\begin{pmatrix} \cdots & \cdots & \cdots & \cdots & \cdots \\ \cdots & 1 + \frac{1}{4h_{ij}} \boldsymbol{n}_{ij} \otimes \boldsymbol{n}_{ij} & \cdots & -\frac{1}{4h_{ij}} \boldsymbol{n}_{ij} \otimes \boldsymbol{n}_{ij} & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ \cdots & -\frac{1}{4h_{ji}} \boldsymbol{n}_{ji} \otimes \boldsymbol{n}_{ji} & \cdots & 1 + \frac{1}{4h_{ji}} \boldsymbol{n}_{ji} \otimes \boldsymbol{n}_{ji} & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots \end{pmatrix} \begin{pmatrix} \vdots \\ \Delta \boldsymbol{v}_i \\ \vdots \\ \Delta \boldsymbol{v}_j \\ \vdots \end{pmatrix}$$
(19)

$$+ \begin{pmatrix} \vdots \\ -\frac{1}{4h_{ij}} (\boldsymbol{U}_{i}^{\infty} - \boldsymbol{U}_{j}^{\infty}) \cdot \boldsymbol{n}_{ij} \boldsymbol{n}_{ij} \\ \vdots \\ -\frac{1}{4h_{ji}} (\boldsymbol{U}_{j}^{\infty} - \boldsymbol{U}_{i}^{\infty}) \cdot \boldsymbol{n}_{ji} \boldsymbol{n}_{ji} \\ \vdots \end{pmatrix}$$

$$(20)$$

References

J. R. Melrose and R. C. Ball. Continuous shear thickening transitions in model concentrated colloids—the role of interparticle forces. *J. Rheol.*, 48:937–960, 2004.