



PARTICIPANT NAME: LINDSEY STICKLER
DATE OF BIRTH: 04/22/1983
PHONE: +1 615 440 8522
EMAIL: lindsey@alyndata.com

FORM SUBMISSION DATE: 05/19/2025
EMERGENCY CONTACT: LINDSEY STICKLER
EMERGENCY CONTACT PHONE: +1 615 440 8522

This report is based on the information that you have provided. PLEASE notify your facilitator of any changes.

MEDICAL HISTORY CONSIDERATIONS

HIGH BLOOD PRESSURE

Ayahuasca consumption can cause a temporary increase in your blood pressure and heart rate. While this effect is typically modest and short lived, it's important for individuals who have hypertension to make sure their blood pressure is well controlled before attending a retreat. If you have a home blood pressure cuff that you trust, check your blood pressure several times before attending a retreat. If you are concerned about your blood pressure, have it checked by your health care provider. Review your blood pressure results with your medical care provider prior to attending a retreat.

DIABETICS WHO REQUIRE INSULIN

Ayahuasca consumption can lead to changes in your blood sugar levels. That, combined with eating less on the day of the ceremony, can add to the challenges of regulating your blood sugar. Before attending a retreat, it is very important that you have an excellent understanding of how to measure your blood sugar and how to adjust your insulin dosing prior to and after ceremony. If you have any questions about this topic, PLEASE discuss this with your health care provider.

If you check your blood sugar levels at home, bring your glucometer with you to the retreat.

Finally, it is important that you discuss this issue with your facilitator to ensure that everyone involved in your experience is comfortable with the situation.

SEIZURE

Ayahuasca can make it more likely for an individual who has a history of seizures to have a seizure. This is because Ayahuasca can lower the seizure threshold. If you have ever had a seizure, this

MUST be discussed this with your facilitator before attending a retreat. Your facilitator may ask that you speak to your health care provider and get his/her approval.

MENTAL HEALTH HISTORY

MAJOR BIPOLAR DISORDER / MANIC DEPRESSION

For individuals with a history of bipolar disorder (manic depression), ayahuasca poses significant risks due to its intense psychoactive effects. The profound nature of these experiences can trigger or worsen manic episodes, depressive symptoms, or psychotic states, potentially resulting in prolonged psychological instability or distress. Even when symptoms are currently managed, ayahuasca's powerful emotional and cognitive impact may disrupt your treatment and undermine long-term mental health stability.

Given these substantial risks, you **must discuss this thoroughly with both your facilitator and your mental health professional** before participating in any plant medicine experience.

MEDICATIONS

Current Medications:

1. Metformin
2. Atenolol
3. Sertraline [Zoloft]
4. Albuterol

DRUG INTERACTION FEEDBACK & RECOMMENDATIONS:

MAJOR The risk or severity of serotonin syndrome can be increased when Sertraline is combined with Ayahuasca.

- **Recommendation:** This use of a monoamine oxidase inhibitor with sertraline, or within 14 days of discontinuing sertraline, is contraindicated. PLEASE do not stop this, or any medication, without discussing it with your healthcare provider. Stopping any medication suddenly can cause uncomfortable and potentially dangerous side effects.
 - Guest should be off medication for minimum **8 days** prior to event and remain off medication for **2 days** after the event.

MODERATE Ayahuasca may increase the hypotensive activities of Atenolol.

- **Recommendation:** Monoamine oxidase inhibitors (MAOIs) should be used with caution in combination with antihypertensive drugs, including thiazide diuretics and β -blockers, since exaggerated hypotensive effects may result. The blood pressure of patients taking

such a combination of medications should be monitored regularly to ensure that levels never become too low or perhaps to modify dosages of either the MAOIs or antihypertensives. Moreover, the occurrence of orthostatic hypotension correlates with peak plasma MAOI levels. Indeed, minimizing peak levels by more frequent administration of smaller doses may be effective in attenuating the orthostatic response. Hypotension related to the hydrazine MAOIs (e.g., phenelzine) is less amenable to changes in the dosing interval. Interventions targeted at maintaining intravascular volume (e.g., encouraging fluid intake, increasing dietary salt, administration of salt-retaining corticosteroids such as fludrocortisone, or the use of support stockings) may be helpful. The occurrence of hypotension with the MAOIs that are not irreversible in their action is significantly less than with the irreversible MAOIs. PLEASE do not stop this, or any medication, without discussing it with your healthcare provider. Stopping any medication suddenly can cause uncomfortable and potentially dangerous side effects.

DRUG AND ALCOHOL USE:

LINDSEY STICKLER has agreed to the following standards regarding the use of the following leading up to and after the event:

- **Alcohol:** Avoid for 48 hours prior to event and for 48 hours after the event
- **Amphetamines:** Avoid for 7 days prior to event and for 48 hours after the event
- **Cocaine** (Occasional Use): Avoid for 72 hours prior to event and for 36 hours after the event
- **Cocaine** (Regular Use): Avoid for 10 days prior to event and for 48 hours after the event
- **MDMA:** Avoid for 72 hours prior to event and for 48 hours after the event
- **Narcotics:** Avoid for 72 hours prior to event and for 72 hours after the event.
- **Cannabis:** LINDSEY STICKLER has agreed to discuss cannabis use if applicable with the Facilitator.

SUPPLEMENTS:

For better or worse, neither supplement manufacturers nor their products are regulated by government agencies to the degree that prescription pharmaceuticals are regulated. As a result, it is incredibly difficult as a consumer to have the same degree of confidence about what goes into supplements. In addition, the regulations around accurate labeling is far less strenuous for the supplement manufacturers. This makes it very difficult to have confidence about predicting potential interactions between supplements and ayahuasca. There are several common supplements that should ABSOLUTELY be avoided, as they are known to be problematic when mixed with ayahuasca.

These include:

Boswellia	Kava
Rhodiola Rosea	Yohimbe
Ephedra	Kratom
Scotch Broom	Ginseng
Licorice Root	Sinicuichi
Kanna	Nutmeg
St. Johns Wort	Methylene blue

As such, it is generally recommended that all supplements should be stopped at least one week prior to participating in a ceremony.

OVER THE COUNTER (OTC) MEDICATIONS:

There are several common OTC medications that should be avoided for **at least one week** prior to ceremony. These medications contain ingredients that, if combined with ayahuasca, can lead to unpleasant and potentially harmful side effects.

1. Cough and Cold Medications containing dextromethorphan

- Dextromethorphan (found in many cough suppressants, such as Robitussin DM, Delsym, and NyQuil)
- Often (but not always) indicated by DM in the name of the drug.

2. Decongestants

- Pseudoephedrine (Sudafed, found in many cold and allergy medications)
- Phenylephrine (found in many nasal sprays and oral decongestants, such as Sudafed PE, Theraflu, or Tylenol Cold)
- Risk: Can cause a hypertensive crisis when combined with MAOIs.

3. Risk: Can cause a hypertensive crisis when combined with MAOIs.

- Chlorpheniramine (found in many cold medicines like Coricidin HBP)

b. brompheniramine (Bromfed, Dimetapp, Bromfenex, Dimetane, BPN, Lodrane)

4. **Weight Loss Supplements**

a. Many OTC weight loss supplements contain stimulants, such as ephedrine, synephrine, or caffeine.

5. **Stimulants (Caffeine-containing products)**

a. High doses of caffeine (energy drinks, weight loss supplements, and some pain relievers like Excedrin)

6. **Sleep aids that contain L-Tryptophan**

QUESTIONS/CONCERNS FROM GUEST:

This is my first event and I'm nervous.

;