

## GEE for Mean and Association (Continued)

Model:

$$\begin{aligned}g(\mu_{ij}) &= \mathbf{X}_{ij}^T \boldsymbol{\beta} \\ \ln(\nu_{ijk}) &= \mathbf{Z}_{ijk}^T \boldsymbol{\alpha}\end{aligned}$$

where

$$\begin{aligned}\mu_{ij} &= E(Y_{ij}) = \Pr[Y_{ij} = 1] \\ \nu_{ijk} &= OR(Y_{ij}, Y_{ik})\end{aligned}$$

e.g.  $\ln(\nu_{ijk}) = \alpha_0 + \alpha_1 |t_{ij} - t_{ik}|^{-1}$ , no constraint on  $\alpha$ 's.

## GEE2

$$\mathbf{U}_{\beta}(\beta, \alpha) = \sum_{i=1}^m \frac{\partial \boldsymbol{\mu}_i^T}{\partial \beta} \mathbf{V}_i^{-1} (\mathbf{Y}_i - \boldsymbol{\mu}_i) = 0$$

$$\mathbf{U}_{\alpha}(\beta, \alpha) = \sum_{i=1}^m \frac{\partial \boldsymbol{\eta}_i^T}{\partial \alpha} \mathbf{H}_i^{-1} (\mathbf{W}_i - \boldsymbol{\eta}_i) = 0$$

where

$\mathbf{V}_i$  = working covariance of  $\mathbf{Y}_i$

$\mathbf{W}_i = (R_{i1}R_{i2}, R_{i2}R_{i3}, \dots, R_{i,n_i-1}R_{i,n_i})^T$

$\boldsymbol{\eta}_i = E(\mathbf{W}_i)$

$\mathbf{H}_i$  = working covariance of  $\mathbf{W}_i$

## GEE2: Remarks

- One can set  $\mathbf{H}_i = \text{diag}\{\text{var}(R_{i1}R_{i2}), \dots, \text{var}(R_{i,n_i-1}R_{in_i})\}$  then  $\boldsymbol{\eta}_i$  and  $\mathbf{H}_i$  are fully determined by the marginal mean  $\boldsymbol{\mu}_i$  and the marginal odds ratios  $\boldsymbol{\nu}_i$ .
- The exact covariance of  $\mathbf{W}_i$  depends on higher order moments of  $\mathbf{Y}_i$ .
- In principle, we could continue modeling the higher order moments... until we modeled them all...

## GEE2: $\eta_i$ and $\mu_i, \nu_i$

Question: How is  $\eta_i$  related to  $\mu_i, \nu_i$ ?

$$\begin{aligned}\eta_{ijk} &= E(W_{ijk}) = \frac{E(Y_{ij}Y_{ik}) - \mu_{ij}\mu_{ik}}{\{\mu_{ij}(1 - \mu_{ij})\mu_{ik}(1 - \mu_{ik})\}^{1/2}} \\ &= \frac{\mu_{ijk} - \mu_{ij}\mu_{ik}}{\{\mu_{ij}(1 - \mu_{ij})\mu_{ik}(1 - \mu_{ik})\}^{1/2}} \\ \mu_{ijk} &= Pr[Y_{ij} = Y_{ik} = 1] \\ &= \begin{cases} \frac{1 - (\mu_{ij} + \mu_{ik})(1 - \nu_{ijk}) - \{[1 - (\mu_{ij} + \mu_{ik})(1 - \nu_{ijk})]^2 - 4(\nu_{ijk} - 1)\nu_{ijk}\mu_{ij}\mu_{ik}\}^{1/2}}{2(\nu_{ijk} - 1)}, & \text{if } \nu_{ijk} \neq 1 \\ \mu_{ij}\mu_{ik}, & \text{if } \nu_{ijk} = 1 \end{cases}\end{aligned}$$

## GEE2: Distribution of $\hat{\beta}$ and $\hat{\alpha}$

$$\begin{aligned}\Rightarrow \mathbf{U}(\delta) &= \sum_i \frac{\partial(\mu_i, \eta_i)^T}{\partial \delta} \begin{pmatrix} \mathbf{V}_i^{-1} & 0 \\ 0 & \mathbf{H}_i^{-1} \end{pmatrix} \begin{pmatrix} \mathbf{Y}_i - \mu_i \\ \mathbf{W}_i - \eta_i \end{pmatrix} \\ &= \sum_i \mathbf{C}_i^T \mathbf{B}_i \begin{pmatrix} \mathbf{Y}_i - \mu_i \\ \mathbf{W}_i - \eta_i \end{pmatrix}\end{aligned}$$

where  $\delta = (\beta^T, \alpha^T)^T$ . Then:

$$\hat{\delta} \sim N(\delta, \mathbf{V}_\delta) \text{ asymptotically}$$

where

$$\begin{aligned}\mathbf{V}_\delta &= \frac{1}{m} \left( \sum_{i=1}^m \mathbf{C}_i^T \mathbf{B}_i \mathbf{C}_i \right)^{-1} \left\{ \sum_{i=1}^m \mathbf{C}_i^T \mathbf{B}_i \begin{pmatrix} \mathbf{Y}_i - \mu_i \\ \mathbf{W}_i - \eta_i \end{pmatrix} \right. \\ &\quad \left. \begin{pmatrix} \mathbf{Y}_i - \mu_i \\ \mathbf{W}_i - \eta_i \end{pmatrix}^T \mathbf{B}_i^T \mathbf{C}_i \right\} \left( \sum_{i=1}^m \mathbf{C}_i^T \mathbf{B}_i \mathbf{C}_i \right)^{-1}\end{aligned}$$

## Alternative Estimating Equation

An alternative EE:

$$\mathbf{U}(\boldsymbol{\delta}) = \sum_i \frac{\partial(\mu_i, \eta_i)^T}{\partial \boldsymbol{\delta}} \text{cov}^{-1} \begin{pmatrix} \mathbf{Y}_i \\ \mathbf{W}_i \end{pmatrix} \begin{pmatrix} \mathbf{Y}_i - \mu_i \\ \mathbf{W}_i - \eta_i \end{pmatrix}$$

Now  $\text{cov}^{-1} \begin{pmatrix} \mathbf{Y}_i \\ \mathbf{W}_i \end{pmatrix}$  may not be a block diagonal matrix.

## Dis/Advantages

**Advantage:** If  $\text{cov}^{-1} \begin{pmatrix} \mathbf{Y}_i \\ \mathbf{W}_i \end{pmatrix}$  is correctly specified, then  $\hat{\delta}$  would be more efficient and  $\mathbf{U}(\delta)$  is optimal.

**Disadvantage:** If  $\text{cov}^{-1} \begin{pmatrix} \mathbf{Y}_i \\ \mathbf{W}_i \end{pmatrix}$  is misspecified and contains off-block-diagonal elements, then misspecification of the  $\nu$  (OR) model would result in a biased estimator of the mean parameter vector  $\beta$ .

# Alternating Logistic Regression (ALR)

Reference: Carey, Zeger, Diggle (1993)

Model:

$$\begin{aligned}g(\mu_{ij}) &= \mathbf{X}_{ij}^T \boldsymbol{\beta} \\ \ln(\nu_{ijk}) &= \mathbf{Z}_{ijk}^T \boldsymbol{\alpha}\end{aligned}$$

GEE2 can be fitted as follows:

1. Assign an initial value  $\boldsymbol{\alpha}_0$ .
2. Use GEE1 to estimate  $\boldsymbol{\beta}$  and  $\mu_{ij}$ .
3. Fit the following offset logistic model:

$$\text{logit Pr}[Y_{ij} = 1 | Y_{ik} = 1] = Y_{ik} \mathbf{Z}_{ijk}^T \boldsymbol{\alpha} + \ln \left( \frac{\mu_{ij} - \mu_{ijk}}{1 - \mu_{ij} - \mu_{ik} + \mu_{ijk}} \right)$$

4. Iterate between 2 and 3.



# Indonesian Infectious Disease Data (Again)

- Data:
  - $m = 275$  children examined every 3 months up to 6 consecutive quarters
  - Outcome=respiratory infection (Y/N)
  - Covariates=XERO, Gender, Age, Height, Season
- Models considered:
  - **Model 1:** GEE1 (cross-sectional age effect:  $\beta_1 age_{ij} + \beta_2 age_{ij}^2$ )
  - **Model 2:** GEE2 (cross-sectional age effect)

$$\ln \nu_{ijk} = \alpha$$

- **Model 3:** GEE2 (distinguish cross-sectional and longitudinal effects, no seasonal effect)

$$\beta_1 age_{i1} + \beta_2 age_{i1}^2 + \beta_3 (age_{ij} - age_{i1}) + \beta_4 (age_{ij} - age_{i1})^2$$

# Indonesian Infectious Disease Data: Results

Variable	Model			
	1	2	3	4
Intercept	-1.47 (0.36)	-2.05 (0.21)	-1.76 (0.25)	-2.21 (0.32)
Gender	-0.66 (0.44)	-0.49 (0.24)	-0.53 (0.24)	-0.53 (0.24)
Height for age	-0.11 (0.041)	-0.042 (0.023)	-0.051 (0.025)	-0.048 (0.024)
Seasonal cosine	-	-0.59 (0.17)	-	-0.54 (0.21)
Seasonal sine	- -	-0.16 (0.14)	- -	-0.016 (0.18)
Xerophthalmia	0.44 (1.15)	0.50 (0.44)	0.53 (0.45)	0.64 (0.44)
Age	-0.089 (0.027)	-0.030 (0.008)	- -	- -
Age <sup>2</sup>	-0.0026 (0.0011)	-0.0010 (0.0004)	- -	- -
Age at entry	- -	- -	-0.053 (0.013)	-0.053 (0.013)
(Age at entry) <sup>2</sup>	- -	- -	-0.0013 (0.0005)	-0.0013 (0.0005)
Follow-up time	- -	- -	-0.19 (0.071)	-0.082 (0.099)
(Follow-up) <sup>2</sup>	- -	- -	0.013 (0.004)	0.007 (0.007)

# Indonesian Infectious Disease Data: Cross Sectional

EXAMPLES

161




Fig. 8.1. Prevalence of respiratory infection as a function of age for three different models. — : Model 1; ..... : Model 2; --- : Model 3.

# Indonesian Infectious Disease Data: Longitudinal




Fig. 8.3. The logarithm of the risk of respiratory infection as a function of follow-up time relative to the risk at an individual's first visit.