

Optimization Theory

Lecture 03

Fudan University

luoluo@fudan.edu.cn

1 Subdifferential Calculus

2 Regularity Conditions

1 Subdifferential Calculus

2 Regularity Conditions

Subdifferential Calculus

Let f_1 and f_2 be proper convex functions on \mathbb{R}^d , then

$$\partial(f_1 + f_2)(\mathbf{x}) \supseteq \partial f_1(\mathbf{x}) + \partial f_2(\mathbf{x}).$$

If the sets $\text{ri}(\text{dom } f_1)$ and $\text{ri}(\text{dom } f_2)$ have a point in common (overlap sufficiently), we have

$$\partial(f_1 + f_2)(\mathbf{x}) = \partial f_1(\mathbf{x}) + \partial f_2(\mathbf{x}).$$

We define the relative interior $\text{ri}(\mathcal{C})$ for convex $\mathcal{C} \subseteq \mathbb{R}^d$ as

$$\text{ri}(\mathcal{C}) = \{\mathbf{z} \in \mathcal{C} : \text{for every } \mathbf{x} \in \mathcal{C} \text{ such that} \\ \text{there exist a } \mu > 1 \text{ such that } (1 - \mu)\mathbf{x} + \mu\mathbf{z} \in \mathcal{C}\}.$$

It means every line segment in \mathcal{C} having \mathbf{z} as one endpoint can be prolonged beyond \mathbf{z} without leaving \mathcal{C} .

Subdifferential Calculus

Nonempty subdifferential and convexity:

- ① If any $\mathbf{x} \in \text{dom } f$ satisfies $\partial f(\mathbf{x}) \neq \emptyset$, then f is convex.
- ② If $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex and \mathbf{x} belongs to the interior of $\text{dom } f$, then $\partial f(\mathbf{x}) \neq \emptyset$.

Theorem (Hyperplane Separation Theorem)

Let $\mathcal{X} \subseteq \mathbb{R}^d$ is a convex set and \mathbf{x}_0 belongs to its boundary. Then, there exists a nonzero vector $\mathbf{w} \in \mathbb{R}^d$ such that

$$\langle \mathbf{w}, \mathbf{x} \rangle \leq \langle \mathbf{w}, \mathbf{x}_0 \rangle.$$

The subgradient of a convex function may not exist at a boundary point of the domain.

As an example, consider the function

$$f(x) = -\sqrt{x}$$

defined on $[0, +\infty)$, where we have $\partial f(0) = \emptyset$.

Subdifferential Calculus

Given matrix $\mathbf{A} \in \mathbb{R}^{d \times m}$ and vector $\mathbf{b} \in \mathbb{R}^d$, define

$$h(\mathbf{x}) = f(\mathbf{Ax} + \mathbf{b}),$$

where f is a proper convex on \mathbb{R}^d . Then $h(\mathbf{x})$ is convex and

$$\partial h(\mathbf{x}) \supseteq \mathbf{A}^\top \partial f(\mathbf{Ax} + \mathbf{b}).$$

If the range of \mathbf{A} contains a point of $\text{ri}(\text{dom } h)$, then

$$\partial h(\mathbf{x}) = \mathbf{A}^\top \partial f(\mathbf{Ax} + \mathbf{b}).$$

Optimal Condition

Theorem

Consider proper closed convex function f and closed convex set $\mathcal{C} \subseteq (\text{dom } f)^\circ$. A point $\mathbf{x}^* \in \mathcal{C}$ is a solution of convex optimization problem

$$\min_{\mathbf{x} \in \mathcal{C}} f(\mathbf{x})$$

if and only if

$$\mathbf{0} \in \partial(f(\mathbf{x}^*) + \mathbb{1}_{\mathcal{C}}(\mathbf{x}^*)).$$

Equivalently, there exists a subgradient $\mathbf{g}^* \in \partial f(\mathbf{x}^*)$, such that any $\mathbf{y} \in \mathcal{C}$ satisfies

$$\langle \mathbf{g}^*, \mathbf{y} - \mathbf{x}^* \rangle \geq 0.$$

In particular, the point \mathbf{x}^* is the solution of the problem in unconstrained case if

$$\mathbf{0} \in \partial f(\mathbf{x}^*).$$

Outline

1 Subdifferential Calculus

2 Regularity Conditions

Regularity Conditions

The following regularity conditions are useful in the convergence analysis of convex optimization problems.

- 1 We say that a function $f : \mathcal{C} \rightarrow \mathbb{R}$ is G -Lipschitz continuous if for all $\mathbf{x}, \mathbf{y} \in \mathcal{C}$, we have

$$|f(\mathbf{x}) - f(\mathbf{y})| \leq G \|\mathbf{x} - \mathbf{y}\|_2.$$

- 2 We say a differentiable function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is L -smooth if it has L -Lipschitz continuous gradient. That is, for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$, we have

$$\|\nabla f(\mathbf{x}) - \nabla f(\mathbf{y})\|_2 \leq L \|\mathbf{x} - \mathbf{y}\|_2.$$

- 3 If the function

$$g(\mathbf{x}) = f(\mathbf{x}) - \frac{\mu}{2} \|\mathbf{x}\|_2^2$$

is convex for some $\mu > 0$, we say f is μ -strongly convex.

Strong Convexity

Theorem

The function $f : \mathcal{C} \rightarrow \mathbb{R}$ defined on convex set \mathcal{C} is μ -strongly-convex if and only if

$$f(\alpha \mathbf{x} + (1 - \alpha) \mathbf{y}) \leq \alpha f(\mathbf{x}) + (1 - \alpha) f(\mathbf{y}) - \frac{\mu \alpha (1 - \alpha)}{2} \|\mathbf{x} - \mathbf{y}\|_2^2$$

for all $\mathbf{x}, \mathbf{y} \in \mathcal{C}$ and $\alpha \in [0, 1]$.

Theorem

If a function f is differentiable on open set \mathcal{C} , then it is μ -strongly convex on \mathcal{C} if and only if

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{\mu}{2} \|\mathbf{y} - \mathbf{x}\|_2^2$$

holds for any $\mathbf{x}, \mathbf{y} \in \mathcal{C}$.

Strong Convexity

If there exists some

$$\mathbf{x}^* = \arg \min_{\mathbf{x} \in \mathcal{C}} f(\mathbf{x}),$$

then it is the unique minimizer.

Moreover, the solution is stable such that any approximate solution $\hat{\mathbf{x}}$ satisfying

$$f(\mathbf{x}) \leq f(\mathbf{x}^*) + \epsilon$$

leads to

$$\|\mathbf{x}^* - \hat{\mathbf{x}}\|_2^2 \leq \frac{2\epsilon}{\mu}.$$

Lipschitz Continuity and Smoothness

Theorem

A convex function f is G -Lipschitz continuous on $\text{dom } f$ if

$$\max_{\mathbf{g} \in \partial f(\mathbf{x})} \{\|\mathbf{g}\|_2\} \leq G$$

for all $\mathbf{x} \in \text{dom } f$.

Theorem

A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is L -smooth (possibly nonconvex), then it holds

$$|f(\mathbf{y}) - f(\mathbf{x}) - \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle| \leq \frac{L}{2} \|\mathbf{y} - \mathbf{x}\|_2^2$$

holds for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$.

Theorem

A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex and L -smooth, then we have

- ① $0 \leq f(\mathbf{y}) - f(\mathbf{x}) - \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle \leq \frac{L}{2} \|\mathbf{y} - \mathbf{x}\|_2^2$
- ② $f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2L} \|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2^2 \leq f(\mathbf{y})$
- ③ $\frac{1}{L} \|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2^2 \leq \langle \nabla f(\mathbf{x}) - \nabla f(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle$

for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$.