

# Multivariate Statistical Analysis

## Lecture 14

Fudan University

luoluo@fudan.edu.cn

- 1 Multivariate Analysis of Variance
- 2 Multivariate Linear Regression
- 3 Bayesian Multivariate Linear Regression

- 1 Multivariate Analysis of Variance
- 2 Multivariate Linear Regression
- 3 Bayesian Multivariate Linear Regression

# Multivariate Analysis of Variance

We consider testing the equality of means with common covariance.

Let  $\mathbf{x}_\alpha^{(g)}$  be an observation from the  $g$ -th population  $\mathcal{N}_p(\boldsymbol{\mu}^{(g)}, \boldsymbol{\Sigma})$  for  $\alpha = 1, \dots, N_g$  and  $g = 1, \dots, q$ . We wish to test the hypothesis

$$H_0 : \boldsymbol{\mu}_1 = \cdots = \boldsymbol{\mu}_g.$$

# Multivariate Analysis of Variance

The likelihood function is

$$L(\boldsymbol{\mu}^{(1)}, \dots, \boldsymbol{\mu}^{(g)}, \boldsymbol{\Sigma}) \\ = \prod_{g=1}^q \frac{1}{(2\pi)^{\frac{pN_g}{2}} (\det(\boldsymbol{\Sigma}))^{\frac{N_g}{2}}} \exp \left( -\frac{1}{2} \sum_{\alpha=1}^{N_g} (\mathbf{x}_{\alpha}^{(g)} - \boldsymbol{\mu}^{(g)})^{\top} \boldsymbol{\Sigma}^{-1} (\mathbf{x}_{\alpha}^{(g)} - \boldsymbol{\mu}^{(g)}) \right).$$

- ① We let  $\boldsymbol{\theta} = \{\boldsymbol{\mu}^{(1)}, \dots, \boldsymbol{\mu}^{(q)}, \boldsymbol{\Sigma}\}$  be the parameters.
- ② The set  $\Omega$  is the space in which  $\boldsymbol{\Sigma}$  is positive definite and each  $\boldsymbol{\mu}^{(g)}$  is any  $p$ -dimensional vector.
- ③ The set  $\omega$  is the space in which  $\boldsymbol{\mu}^{(1)} = \dots = \boldsymbol{\mu}^{(g)}$  ( $p$ -dimensional vectors) and  $\boldsymbol{\Sigma}$  is positive definite matrix.

# Multivariate Analysis of Variance

The likelihood ratio criterion is

$$\lambda = \frac{\sup_{\theta \in \omega} L(\theta)}{\sup_{\theta \in \Omega} L(\theta)} = \frac{(\det(\hat{\Sigma}_{\Omega}))^{\frac{N}{2}}}{(\det(\hat{\Sigma}_{\omega}))^{\frac{N}{2}}},$$

where

$$\hat{\Sigma}_{\Omega} = \frac{1}{N} \sum_{g=1}^q \sum_{\alpha=1}^{N_g} (\mathbf{x}_{\alpha}^{(g)} - \bar{\mathbf{x}}^{(g)}) (\mathbf{x}_{\alpha}^{(g)} - \bar{\mathbf{x}}^{(g)})^{\top}$$

and

$$\hat{\Sigma}_{\omega} = \frac{1}{N} \sum_{g=1}^q \sum_{\alpha=1}^{N_g} (\mathbf{x}_{\alpha}^{(g)} - \bar{\mathbf{x}}) (\mathbf{x}_{\alpha}^{(g)} - \bar{\mathbf{x}})^{\top}.$$

# Multivariate Analysis of Variance

We can write

$$N\hat{\Sigma}_{\omega} = \mathbf{A} + \mathbf{B},$$

where

$$\mathbf{A} = N\hat{\Sigma}_{\Omega} = \sum_{g=1}^q \sum_{\alpha=1}^{N_g} (\mathbf{x}_{\alpha}^{(g)} - \bar{\mathbf{x}}^{(g)}) (\mathbf{x}_{\alpha}^{(g)} - \bar{\mathbf{x}}^{(g)})^{\top} \sim \mathcal{W}_p(\mathbf{\Sigma}, N - q)$$

and

$$\mathbf{B} = \sum_{g=1}^q N_g (\bar{\mathbf{x}}^{(g)} - \bar{\mathbf{x}}) (\bar{\mathbf{x}}^{(g)} - \bar{\mathbf{x}})^{\top} \sim \mathcal{W}_p(\mathbf{\Sigma}, q - 1)$$

are independent.

# Wilks' Lambda distribution

For two independent random matrices  $\mathbf{A} \sim \mathcal{W}_p(\mathbf{\Sigma}, n)$  and  $\mathbf{B} \sim \mathcal{W}_p(\mathbf{\Sigma}, m)$  with  $n \geq p$ , the ratio

$$\frac{\det(\mathbf{A})}{\det(\mathbf{A} + \mathbf{B})}$$

has Wilks' Lambda distribution with degrees of freedom  $n$  and  $m$ , which is typically written as

$$\frac{\det(\mathbf{A})}{\det(\mathbf{A} + \mathbf{B})} \sim \Lambda_{p,n,m}.$$



## Theorem

Let  $\mathbf{A} \sim \mathcal{W}_p(\mathbf{\Sigma}, n)$  and  $\mathbf{B} \sim \mathcal{W}_p(\mathbf{\Sigma}, m)$  be two independent Wishart distributed variables, then we can write

$$\frac{\det(\mathbf{A})}{\det(\mathbf{A} + \mathbf{B})} = \prod_{i=1}^p u_i \sim \Lambda_{p,n,m},$$

where  $u_1, \dots, u_p$  are independent distributed as

$$u_i \sim \text{Beta} \left( \frac{n+1-i}{2}, \frac{m}{2} \right).$$

# Generalized Variance

Let  $\mathbf{A} \sim \mathcal{W}_p(\mathbf{\Sigma}, n)$ , we can follow above theorem to show

$$\det(\mathbf{A}) = \det(\mathbf{\Sigma}) \prod_{i=1}^p v_i$$

with some independent random variables  $v_1, \dots, v_p$ ?

# Properties of Wishart Distribution

Let  $\mathbf{A} \sim \mathcal{W}_p(\boldsymbol{\Sigma}, n)$  and partition  $\mathbf{A}$  and  $\boldsymbol{\Sigma}$  into  $q$  and  $p - q$  rows and columns as

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{bmatrix} \quad \text{and} \quad \boldsymbol{\Sigma} = \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{bmatrix},$$

then we have

(a)  $\mathbf{A}_{11} \sim \mathcal{W}_q(\boldsymbol{\Sigma}_{11}, n)$  and  $\mathbf{A}_{22} \sim \mathcal{W}_{p-q}(\boldsymbol{\Sigma}_{22}, n)$ ;

(b) if  $q = 1$ , then

$$\mathbf{a}_{21} \mid \mathbf{A}_{22} \sim \mathcal{N}_{p-q}(\mathbf{A}_{22} \boldsymbol{\Sigma}_{22}^{-1} \boldsymbol{\sigma}_{21}, \sigma_{11.2}^2 \mathbf{A}_{22})$$

where  $\sigma_{11.2}^2 = \sigma_{11} - \boldsymbol{\sigma}_{21}^\top \boldsymbol{\Sigma}_{22}^{-1} \boldsymbol{\sigma}_{21}$ ;

(c) if  $n > p - q$ , then

$$\mathbf{A}_{11.2} = \mathbf{A}_{11} - \mathbf{A}_{12} \mathbf{A}_{22}^{-1} \mathbf{A}_{21} \sim \mathcal{W}_q(\boldsymbol{\Sigma}_{11.2}, n - p + q)$$

is independent on  $\mathbf{A}_{22}$  and  $\mathbf{A}_{12}$ , where  $\boldsymbol{\Sigma}_{11.2} = \boldsymbol{\Sigma}_{11} - \boldsymbol{\Sigma}_{12} \boldsymbol{\Sigma}_{22}^{-1} \boldsymbol{\Sigma}_{21}$ .

# Outline

- 1 Multivariate Analysis of Variance
- 2 Multivariate Linear Regression
- 3 Bayesian Multivariate Linear Regression

# Multivariate Linear Regression

Given dataset  $\{(\mathbf{x}_i, \mathbf{y}_i)\}_{i=1}^N$ , where  $\mathbf{x}_i \in \mathbb{R}^p$  and  $\mathbf{y}_i \in \mathbb{R}^q$  are the feature and the corresponding output of the  $i$ -th data.

We suppose

$$\mathbf{y}_i = \mathbf{B}^\top \mathbf{x}_i + \epsilon_i \quad \text{with} \quad \mathbf{B} \in \mathbb{R}^{p \times q} \quad \text{and} \quad \epsilon_i \stackrel{\text{i.i.d}}{\sim} \mathcal{N}_q(\mathbf{0}, \mathbf{\Sigma})$$

for  $i = 1, \dots, N$ ,  $\mathbf{\Sigma} \succ 0$  and  $N > p$ .

We regard  $\mathbf{B} \in \mathbb{R}^{p \times q}$  and  $\mathbf{\Sigma} \in \mathbb{R}^{q \times q}$  as parameters, then

$$\epsilon_i = \mathbf{y}_i - \mathbf{B}^\top \mathbf{x}_i \sim \mathcal{N}_q(\mathbf{0}, \mathbf{\Sigma}).$$

# Multivariate Linear Regression

We denote

$$\mathbf{X} = \begin{bmatrix} \mathbf{x}_1^\top \\ \vdots \\ \mathbf{x}_N^\top \end{bmatrix} \in \mathbb{R}^{N \times p}, \quad \mathbf{Y} = \begin{bmatrix} \mathbf{y}_1^\top \\ \vdots \\ \mathbf{y}_N^\top \end{bmatrix} \in \mathbb{R}^{N \times q} \quad \text{and} \quad \mathbf{E} = \begin{bmatrix} \boldsymbol{\epsilon}_1^\top \\ \vdots \\ \boldsymbol{\epsilon}_N^\top \end{bmatrix} \in \mathbb{R}^{N \times q},$$

and suppose  $\mathbf{X}$  is full rank.

# MLE for Multivariate Linear Regression

We construct the likelihood function for  $\epsilon_1, \dots, \epsilon_N$  as follows

$$\begin{aligned} L(\mathbf{B}, \mathbf{\Sigma}) &= \prod_{\alpha=1}^N \frac{1}{\sqrt{(2\pi)^p \det(\mathbf{\Sigma})}} \exp \left( -\frac{1}{2} (\mathbf{B}^\top \mathbf{x}_\alpha - \mathbf{y}_\alpha)^\top \mathbf{\Sigma}^{-1} (\mathbf{B}^\top \mathbf{x}_\alpha - \mathbf{y}_\alpha) \right) \\ &= \frac{1}{(2\pi)^{Np/2} (\det(\mathbf{\Sigma}))^{N/2}} \exp \left( -\frac{1}{2} \text{tr} \left( (\mathbf{XB} - \mathbf{Y}) \mathbf{\Sigma}^{-1} (\mathbf{XB} - \mathbf{Y})^\top \right) \right). \end{aligned}$$

The maximum likelihood estimators are

$$\hat{\mathbf{B}} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{Y} \quad \text{and} \quad \hat{\mathbf{\Sigma}} = \frac{1}{N} \mathbf{Y}^\top (\mathbf{I} - \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top) \mathbf{Y}.$$

# MLE for Multivariate Linear Regression

We write

$$\mathbf{B} = [\beta_1 \quad \cdots \quad \beta_q] \in \mathbb{R}^{q \times p} \quad \text{and} \quad \hat{\mathbf{B}} = [\hat{\beta}_1 \quad \cdots \quad \hat{\beta}_q] \in \mathbb{R}^{q \times p}.$$

Then the joint distribution of  $\hat{\beta}_1, \dots, \hat{\beta}_N$  is normal and we have

- ①  $\mathbb{E}[\hat{\beta}_i] = \beta_i$ ;
- ②  $\text{Cov}[\hat{\beta}_i, \hat{\beta}_j] = \sigma_{ij}(\mathbf{X}^\top \mathbf{X})^{-1}$ ;
- ③  $\hat{\Sigma} \sim \mathcal{W}_q\left(\frac{1}{N}\Sigma, N - p\right)$ .



- 1 Multivariate Analysis of Variance
- 2 Multivariate Linear Regression
- 3 Bayesian Multivariate Linear Regression

# Bayesian Multivariate Linear Regression

We can additionally suppose each  $b_{ij}$  independently follows

$$b_{ij} \sim \mathcal{N}(0, \tau^2),$$

then the posterior likelihood function is

$$\begin{aligned} L(\mathbf{B}, \mathbf{\Sigma}) &= \prod_{i=1}^N \frac{1}{\sqrt{(2\pi)^p \det(\mathbf{\Sigma})}} \exp \left( -\frac{1}{2} (\mathbf{B}^\top \mathbf{x}_i - \mathbf{y}_i)^\top \mathbf{\Sigma}^{-1} (\mathbf{B}^\top \mathbf{x}_i - \mathbf{y}_i) \right) \\ &\quad \cdot \prod_{i=1}^p \prod_{j=1}^q \frac{1}{\sqrt{2\pi\tau^2}} \exp \left( -\frac{b_{ij}^2}{2\tau^2} \right) \\ &\propto \frac{1}{(\det(\mathbf{\Sigma}))^{N/2}} \exp \left( -\frac{1}{2} \text{tr}((\mathbf{XB} - \mathbf{Y})\mathbf{\Sigma}^{-1}(\mathbf{XB} - \mathbf{Y})^\top) - \frac{1}{2\tau^2} \|\mathbf{B}\|_F^2 \right), \end{aligned}$$

which leads to

$$\text{vec}(\hat{\mathbf{B}}) = (\mathbf{I}_q \otimes \tau^2 \mathbf{X}^\top \mathbf{X} + \mathbf{\Sigma} \otimes \mathbf{I}_p)^{-1} \text{vec}(\tau^2 \mathbf{X}^\top \mathbf{Y}).$$

# Bayesian Multivariate Linear Regression

We typically suppose

$$\beta_{(i)} \stackrel{\text{i.i.d}}{\sim} \mathcal{N}_q(\mathbf{0}, \tau^2 \mathbf{\Sigma}), \quad \text{where} \quad \mathbf{B} = \begin{bmatrix} \beta_{(1)}^\top \\ \vdots \\ \beta_{(p)}^\top \end{bmatrix} \in \mathbb{R}^{p \times q},$$

then the posterior likelihood function is

$$\begin{aligned} & L(\mathbf{B}, \mathbf{\Sigma}) \\ &= \prod_{i=1}^N \frac{1}{\sqrt{(2\pi)^p \det(\mathbf{\Sigma})}} \exp \left( -\frac{1}{2} (\mathbf{B}^\top \mathbf{x}_i - \mathbf{y}_i)^\top \mathbf{\Sigma}^{-1} (\mathbf{B}^\top \mathbf{x}_i - \mathbf{y}_i) \right) \\ & \quad \cdot \prod_{j=1}^p \frac{1}{\sqrt{(2\pi)^q \det(\mathbf{\Sigma})}} \exp \left( -\frac{1}{2\tau^2} \beta_{(j)}^\top \mathbf{\Sigma}^{-1} \beta_{(j)} \right) \\ & \propto \frac{1}{(\det(\mathbf{\Sigma}))^{N/2}} \exp \left( -\frac{1}{2} \text{tr} \left( (\mathbf{X}\mathbf{B} - \mathbf{Y}) \mathbf{\Sigma}^{-1} (\mathbf{X}\mathbf{B} - \mathbf{Y})^\top \right) - \frac{1}{2\tau^2} \mathbf{B} \mathbf{\Sigma}^{-1} \mathbf{B}^\top \right). \end{aligned}$$

# Bayesian Multivariate Linear Regression

We have

$$\hat{\mathbf{B}}_{\lambda} = (\mathbf{X}^{\top} \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^{\top} \mathbf{Y},$$

and

$$\hat{\boldsymbol{\Sigma}}_{\lambda} = \frac{1}{N} \mathbf{Y}^{\top} (\mathbf{I} - \mathbf{X}(\mathbf{X}^{\top} \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^{\top}) \mathbf{Y},$$

where  $\lambda = 1/\tau^2$ .