

# Multivariate Statistical Analysis

## Lecture 05

Fudan University

luoluo@fudan.edu.cn

- 1 Characteristic Function
- 2 Maximum Likelihood Estimation

1 Characteristic Function

2 Maximum Likelihood Estimation

# Characteristic Function

The characteristic function of a  $p$ -dimensional random vector  $\mathbf{x}$  is

$$\phi(\mathbf{t}) = \mathbb{E} \left[ \exp(\mathbf{i} \mathbf{t}^\top \mathbf{x}) \right]$$

defined for every real vector  $\mathbf{t} \in \mathbb{R}^p$ .

For the complex-valued function  $g(z)$  be written as

$$g(z) = g_1(z) + \mathbf{i} g_2(z),$$

where  $g_1(z)$  and  $g_2(z)$  are real-valued, the expected value of  $g(z)$  is

$$\mathbb{E}[g(z)] = \mathbb{E}[g_1(z)] + \mathbf{i} \mathbb{E}[g_2(z)].$$

## Theorem

*If the  $p$ -dimensional random vector  $\mathbf{x}$  has the density  $f(\mathbf{x})$  and the characteristic function  $\phi(\mathbf{t})$ , then*

$$f(\mathbf{x}) = \frac{1}{(2\pi)^p} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp(-i \mathbf{t}^\top \mathbf{x}) \phi(\mathbf{t}) dt_1 \dots dt_p.$$

- If the random variable have a density, the characteristic function determines the density function uniquely.
- If the random variable does not have a density, the characteristic function uniquely defines the probability of any continuity interval.

# Characteristic Function

## Theorem

The characteristic function of  $\mathbf{x}$  distributed according to  $\mathcal{N}_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  is

$$\phi(\mathbf{t}) = \exp \left( i \mathbf{t}^\top \boldsymbol{\mu} - \frac{1}{2} \mathbf{t}^\top \boldsymbol{\Sigma} \mathbf{t} \right).$$

for every  $\mathbf{t} \in \mathbb{R}^p$ .

Sketch of the proof:

- 1 The characteristic function of  $\mathbf{y} \sim \mathcal{N}_p(\mathbf{0}, \mathbf{I})$  is  $\phi_0(\mathbf{t}) = \exp(-\mathbf{t}^\top \mathbf{t}/2)$ .
- 2 For  $\mathbf{x} \sim \mathcal{N}_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ , we have  $\mathbf{x} = \mathbf{A}\mathbf{y} + \boldsymbol{\mu}$  such that  $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}^\top$ .
- 3 Using  $\phi_0(\mathbf{t})$  to present the characteristic function of  $\mathbf{x} \sim \mathcal{N}_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ .

# Characteristic Function

## Theorem

The characteristic function of  $\mathbf{x}$  distributed according to  $\mathcal{N}_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  is

$$\phi(\mathbf{t}) = \exp \left( i \mathbf{t}^\top \boldsymbol{\mu} - \frac{1}{2} \mathbf{t}^\top \boldsymbol{\Sigma} \mathbf{t} \right).$$

for every  $\mathbf{t} \in \mathbb{R}^p$ .

This theorem directly implies  $\mathbf{x} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  leads to  $\mathbf{C}\mathbf{x} \sim \mathcal{N}(\mathbf{C}\boldsymbol{\mu}, \mathbf{C}\boldsymbol{\Sigma}\mathbf{C}^\top)$ .



characteristic function



trick of matrix

## Theorem

*If every linear combination of the components of a random vector  $\mathbf{y}$  is normally distributed, then  $\mathbf{y}$  is normally distributed.*

In other words, if the  $p$ -dimensional random vector  $\mathbf{y}$  leads to the univariate random variable

$$\mathbf{u}^\top \mathbf{y}$$

is normally distributed for any fixed  $\mathbf{u} \in \mathbb{R}^p$ , then  $\mathbf{y}$  is normally distributed.

This is another definition of multivariate normal distribution.



# Example

## Theorem

We let

$$\mathbf{x} \sim \mathcal{N}_p(\boldsymbol{\mu}_1, \boldsymbol{\Sigma}_1), \quad \mathbf{y} \sim \mathcal{N}_p(\boldsymbol{\mu}_2, \boldsymbol{\Sigma}_2) \quad \text{and} \quad \mathbf{z} = \mathbf{x} + \mathbf{y}.$$

Suppose that  $\mathbf{x}$  and  $\mathbf{y}$  are independent, then we have

$$\mathbf{z} \sim \mathcal{N}_p(\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2, \boldsymbol{\Sigma}_1 + \boldsymbol{\Sigma}_2).$$



characteristic function



this result

# Outline

1 Characteristic Function

2 Maximum Likelihood Estimation

# The Maximum Likelihood Estimators

## Theorem

If  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N$  constitute a sample from  $\mathcal{N}_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  with  $N > p$ , the maximum likelihood estimators of  $\boldsymbol{\mu}$  and  $\boldsymbol{\Sigma}$  are

$$\hat{\boldsymbol{\mu}} = \bar{\mathbf{x}} = \frac{1}{N} \sum_{\alpha=1}^N \mathbf{x}_{\alpha} \quad \text{and} \quad \hat{\boldsymbol{\Sigma}} = \frac{1}{N} \sum_{\alpha=1}^N (\mathbf{x}_{\alpha} - \bar{\mathbf{x}})(\mathbf{x}_{\alpha} - \bar{\mathbf{x}})^{\top}$$

respectively.

# The Maximum Likelihood Estimators

The likelihood function is

$$L = \frac{1}{(2\pi)^{\frac{pN}{2}} (\det(\mathbf{\Sigma}))^{\frac{N}{2}}} \exp \left[ -\frac{1}{2} \sum_{\alpha=1}^N (\mathbf{x}_{\alpha} - \boldsymbol{\mu})^{\top} \mathbf{\Sigma}^{-1} (\mathbf{x}_{\alpha} - \boldsymbol{\mu}) \right].$$

The vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N$  are fixed at the sample values and  $L$  is a function of  $\boldsymbol{\mu}$  and  $\mathbf{\Sigma}$ .

The logarithm of the likelihood function is

$$\ln L = -\frac{pN}{2} \ln 2\pi - \frac{N}{2} \ln (\det(\mathbf{\Sigma})) - \frac{1}{2} \sum_{\alpha=1}^N (\mathbf{x}_{\alpha} - \boldsymbol{\mu})^{\top} \mathbf{\Sigma}^{-1} (\mathbf{x}_{\alpha} - \boldsymbol{\mu}).$$

# The Maximum Likelihood Estimators

There are some results for estimating the covariance.

## Theorem

*The function  $h : \mathbb{S}_{++}^p \rightarrow \mathbb{R}$  such that*

$$h(\mathbf{X}) = -\log \det(\mathbf{X})$$

*is convex, where  $\mathbb{S}_{++}^p = \{\mathbf{X} \in \mathbb{R}^{p \times p} : \mathbf{X} \succ \mathbf{0}\}$ .*

## Theorem

*If  $\mathbf{D} \in \mathbb{R}^{p \times p}$  is positive definite, the maximum of*

$$f(\mathbf{G}) = -N \ln \det(\mathbf{G}) - \text{tr}(\mathbf{G}^{-1} \mathbf{D})$$

*with respect to positive definite matrices  $\mathbf{G}$  exists, occurs at  $\mathbf{G} = \frac{1}{N} \mathbf{D}$ .*