



Check for updates

OPINION ARTICLE

Integrative Healthcare for Optimal Health

[version 1; peer review: awaiting peer review]

Ravinder Mamtani¹, Amit Abraham ¹, Sohaila Cheema¹, Karima Chaabna¹,
Darshan H Mehta ²⁻⁴, Javaid Sheikh⁵

¹Institute for Population Health, Weill Cornell Medicine - Qatar, Doha, Qatar²Benson-Henry Institute for Mind Body Medicine, Massachusetts General Hospital, Boston, Massachusetts, USA³Osher Center for Integrative Health, Brigham and Women's Hospital, Boston, Massachusetts, USA⁴Harvard Medical School, Boston, Massachusetts, USA⁵Office of the Dean, Weill Cornell Medicine - Qatar, Doha, Qatar

V1 First published: 14 Feb 2026, 15:257
<https://doi.org/10.12688/f1000research.177886.1>

Latest published: 14 Feb 2026, 15:257
<https://doi.org/10.12688/f1000research.177886.1>

Open Peer Review

Approval Status Awaiting Peer Review

Any reports and responses or comments on the article can be found at the end of the article.

Abstract

Conventional medicine and public health have collectively improved health and saved the lives of millions. Yet, as we confront the escalating crisis of non-communicable diseases (NCDs), it becomes increasingly clear that conventional approaches alone are no longer sufficient. A new, evidence-based paradigm shift is necessary. Integrative health, which combines conventional medicine with complementary and lifestyle medicine, offers that shift. To assess its effectiveness, we identified and reviewed 14 clinical trials published within the past 5 years in widely read, high-impact factor journals (and their sub-specialty counterparts) – including *The New England Journal of Medicine*, the *Lancet*, the *BMJ* and *JAMA*. These trials directly compared integrative health interventions with conventional medical treatments across conditions such as insomnia, obesity, age-related disorders, Parkinsonism, and motor neuron disease. The included trials assessed outcomes related to mental health, pain management, and quality of life. Across these studies, we found that integrative health interventions were consistently associated with improved health outcomes. While we acknowledge that stronger evidence is needed, the research findings represent an important step toward exploring a well-balanced, evidence-based approach to integrative health.

Keywords

Integrative Health, Integrative Medicine, Complementary medicine, Alternative medicine, Non-communicable disease

Corresponding author: Amit Abraham (ama2006@qatar-med.cornell.edu)

Author roles: **Mamtani R:** Conceptualization, Data Curation, Formal Analysis, Writing – Original Draft Preparation, Writing – Review & Editing; **Abraham A:** Conceptualization, Data Curation, Formal Analysis, Writing – Original Draft Preparation, Writing – Review & Editing; **Cheema S:** Conceptualization, Data Curation, Formal Analysis, Writing – Original Draft Preparation, Writing – Review & Editing; **Chaabna K:** Data Curation, Formal Analysis, Writing – Original Draft Preparation, Writing – Review & Editing; **Mehta DH:** Writing – Original Draft Preparation, Writing – Review & Editing; **Sheikh J:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2026 Mamtani R *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Mamtani R, Abraham A, Cheema S *et al.* **Integrative Healthcare for Optimal Health [version 1; peer review: awaiting peer review]** F1000Research 2026, **15**:257 <https://doi.org/10.12688/f1000research.177886.1>

First published: 14 Feb 2026, **15**:257 <https://doi.org/10.12688/f1000research.177886.1>

1. Introduction

Over the past century, conventional medicine and public health strategies - such as vaccination programs, sanitation, and acute care interventions - have markedly reduced global mortality and morbidity. Nonetheless, the global rise of non-communicable diseases (NCDs), responsible for nearly 70% of deaths and the majority of disability-adjusted life years (DALYs),^{1,2} reveals the limits of conventional healthcare. While indispensable, conventional approaches often fail to address the multifactorial nature of NCDs, particularly in the absence of patient engagement in self-care.³

One way to address the rise in NCDs is augmenting conventional medicine with complementary and lifestyle modalities. Complementary medicine encompasses diverse practices not traditionally part of mainstream medical systems, including acupuncture, herbal and dietary supplements, meditation, massage therapy, chiropractic care, yoga, and tai chi.⁴ Lifestyle medicine, which overlaps with both conventional and complementary domains, focuses on preventing and managing NCDs through behavioral interventions including nutrition, physical activity, stress reduction, social connectivity, avoidance of harmful substances, and sleep hygiene.⁵ These approaches are not only widely used but are also supported by scientific evidence, demonstrating their potential to improve health outcomes. However they often remain unavailable through conventional healthcare practitioners, which may lead individuals to seek alternative healthcare on their own. In this context, integrative healthcare synthesizes conventional medicine with complementary and lifestyle modalities into a unified, evidence-informed framework that aims to optimize health outcomes and enhance patient well-being.⁶

Given the multifactorial nature of NCDs, the collective use of conventional, complementary, and lifestyle approaches could achieve greater impact than any of these domains in isolation. This is the central promise of integrative healthcare: an evidence-based, whole-person, and sustainable model of care that addresses the biological, psychological, social, and behavioral dimensions of health (Figure 1). Nonetheless, integrative healthcare remains overlooked as a distinct model for managing NCDs.

We argue that the primary challenge facing integrative healthcare for NCDs is not a lack of efficacy, but its marginalization within the mainstream evidence hierarchy. Integrative healthcare is rarely incorporated, despite the fact that existing trials suggest added value over conventional care alone. This commentary illustrates how integrative healthcare currently appears in the literature and outlines key steps toward a research agenda more closely aligned with the realities of integrative healthcare.

2. Evaluating the Value of Integrative Healthcare

It is helpful to review the literature in order to assess the clinical impact of integrative healthcare. Over the past five years, only 14 randomized controlled trials comparing integrative interventions with standard conventional care have been published in high-impact, peer-reviewed medical journals (*The New England Journal of Medicine*, the *Lancet*, the *BMJ*, and *JAMA*) (Table 1). These trials spanned diverse settings and involved participants of varying ages, sexes, and educational backgrounds. Across these studies, integrative healthcare approaches were generally more effective than conventional care alone. In patients with various health conditions (including Parkinson's disease, primary insomnia, and depression), integrative approaches reduced overall symptom severity. Similar added benefits were observed in trials assessing obesity, type 2 diabetes, ageing-associated decline, Crohn's disease, and pain, where integrative healthcare improved clinical outcomes beyond those achieved with conventional treatment alone. A subset of studies included follow-up beyond six and twelve months, suggesting sustained benefits over time.

Beyond clinical outcomes, integrative healthcare emphasizes shared decision-making and patient empowerment. Patients receiving integrative healthcare report higher satisfaction, greater alignment with personal values, and a stronger sense of being heard and respected by their providers.⁷⁻¹⁰ Integrative healthcare has been associated with reduced polypharmacy, improved adherence to healthy behaviors, and enhanced long-term health outcomes.¹¹ When appropriately implemented, integrative healthcare can also reduce medical costs and yield improved health outcomes.^{12,13} These observations pose an important question: if integrative healthcare approaches enhance outcomes and patient experience, why are they not more systematically implemented in NCD management?

3. Addressing challenges in implementation

Despite its promise, the widespread adoption of integrative healthcare faces several barriers. Most healthcare professionals receive limited training in complementary and lifestyle medicine, restricting their ability to implement integrative approaches effectively.^{14,15} One way to address this would be to foster interdisciplinary collaboration, allowing healthcare professionals from different fields to effectively address patient care.^{16,17} In this context, developing policies to incentivize education and research in integrative healthcare and implementing culturally-appropriate, evidence-based guidelines can streamline integrative healthcare approaches.¹⁶⁻¹⁸ The lack of reimbursement mechanisms for integrative healthcare poses a significant obstacle to access and scalability. A shift away from fee-based service




Figure 1. Integrative Health is an evidence-based, holistic model of healthcare.

toward value-based care, which links reimbursement to health outcomes and overall patient satisfaction, may result in more widespread adoption of holistic, integrative healthcare.^{16,19} By prioritizing prevention and resource optimization, integrative healthcare offers a pathway toward more sustainable and equitable health systems.^{20,21}

Table 1. Randomized control trials comparing conventional care and integrative health.

Patient symptoms and clinical diagnosis	Conventional care	Integrative health	Outcome and conclusion	Reference
Motor neuron disease (amyotrophic lateral sclerosis; progressive muscular atrophy; or primary lateral sclerosis)	Standard care, including medication for motor neuron disease and related symptoms; treatments such as non-invasive ventilation, physiotherapy, and gastrostomy; and access to other hospital-based and community-based services (such as equipment and adaptations, orthotics, respiratory, gastroenterology, clinical psychology, neuropsychology and counselling, and social care services) N=94	Along with conventional care, participants received Acceptance and Commitment Therapy (a combination of acceptance, mindfulness, motivation, and behavior change techniques to assist engagement in life-enriching activities in the presence of distressing thoughts and feelings) N=95	Integrative health demonstrated clinical effectiveness in maintaining or improving quality of life	Gould, 2024 ²²
Motor symptoms and constipation in patients with Parkinson's disease	Levodopa Combination of oral lactulose or rectal glycerin N=83	In addition to conventional care, participants received electroacupuncture. N=83	Integrative health enhanced bowel movements, reducing symptoms of constipation. Improved functional motor activity	Li, 2023 ²³
Quality of sleep in patients with Parkinson's disease	Sleep hygiene guidance from sleep clinic physicians with continued use of anti-Parkinson medication and sham acupuncture N=41	In addition to conventional care, participants received acupuncture N=42	Integrative health improved sleep quality	Yan, 202 ^{24a}
Mental disorders (depression, primary insomnia, post-traumatic stress disorder [PTSD], panic disorder, or agoraphobia)	Standard outpatient interventions, including psychological (cognitive behavioral therapy, psychoanalysis, systemic therapy) and pharmacological treatments N=201	In addition to conventional care, participants engaged in an exercise program (ImPuls) (evidence-based, moderate-to-vigorous intensity outdoor exercises lasting 30 min, combined with behavioral change techniques targeting motivational and volitional determinants of exercise behavior) N=199	Integrative health demonstrated superior efficacy in reducing global symptom severity	Wolf, 2024 ²⁵
Insomnia in patients with depression	Recommendations to engage in regular exercise, follow a healthy diet, manage stress levels, and continue regular administration of antidepressants, sedatives, or hypnotics N=90	In addition to conventional care, participants received electroacupuncture N=90	Integrative health alleviated insomnia among patients with depression	Yin, 2022 ^{26b}

Table 1. Continued

Patient symptoms and clinical diagnosis	Conventional care	Integrative health	Outcome and conclusion	Reference
Post-operative ileus secondary to resection of colorectal cancer	Multimodal analgesia, patient-controlled analgesia plus non-steroidal anti-inflammatory drugs, early oral feeds, and early mobilization N=35	In addition to conventional care, participants received electroacupuncture at one of two different locations on the body N=35 for each group	Integrative health enhanced bowel function recovery following post-operative ileus	Yang, 2022 ^{27c}
Crohn's disease	Combination of mesalazine, corticosteroids, azathioprine, or methotrexate N=33	In addition to conventional care, participants received acupuncture N=33	Integrative health induced and maintained remission of active Crohn's disease, improved intestinal flora, and lowered recurrence rates	Bao, 2022 ²⁸
Insomnia and pain in patients with chronic spinal pain	Best evidence pain management (person-centered care; pain neuroscience education and cognition-targeted exercise therapy) N=62	In addition to conventional care, participants received cognitive behavior therapy (sleep education, self-monitoring of sleep patterns, time-in-bed restriction, stimulus control, sleep hygiene, cognitive restructuring, and relaxation) N=61	Integrative health improved sleep quality and reduced the severity of insomnia, but provided no significant pain relief	Malfiet, 2024 ²⁹
Pain relief in urolithiasis	Diclofenac and sham acupuncture N=40	In addition to conventional care, participants received acupuncture N=40	Integrative health provided fast and substantial pain relief to patients with renal colic compared with sham acupuncture in the emergency setting	Tu, 2022 ^{30a}
Obesity	Low-calorie diet and liraglutide N=49	In addition to conventional care, participants engaged in an exercise routine comprising moderate to vigorous aerobic activity and strength training N=48	Integrative health preserved bone mineral density at clinically important fracture sites during weight loss	Jensen, 2024 ³¹
Obesity	Liraglutide N=41	Along with conventional care, participants engaged in an exercise routine (150 min/week of moderate-intensity aerobic physical activity, 75 min/week of vigorous-intensity aerobic physical activity, or an equivalent combination of both) N=40	Integrative health improved healthy weight loss maintenance more than liraglutide alone one year after initiation of the trial	Lundgren, 2021 ^{32d}

Table 1. Continued

Patient symptoms and clinical diagnosis	Conventional care	Integrative health	Outcome and conclusion	Reference
Obesity	Liraglutide N=49	In addition to conventional care, participants engaged in an exercise routine (150 min/week of moderate-intensity aerobic physical activity, 75 min/week of vigorous-intensity aerobic physical activity, or an equivalent combination of both) N=48	Integrative health promoted healthy weight maintenance after treatment termination compared with obesity pharmacotherapy alone and maintained body weight and composition 2 years after initiation of the trial	Jensen, 2024 ^{33d}
Type 2 diabetes	Standard care (glucose-lowering, lipid-lowering, and blood pressure-lowering medications) N=34	In addition to conventional care, participants received individualized meal plans and structured exercise plans N=64	Integrative health improved physical components of health-related quality of life, while mental components remained unchanged	MacDonald, 2021 ³⁴
Aging-associated illness	Primary health care, including health education flyers and lab test results N=199	In addition to conventional care, participants engaged in a multidomain intervention (physical exercise, cognitive training, nutrition, and disease education) N=199	Integrative health improved the quality of life	Lee, 2021 ³⁵

Note: These 14 clinical trials were published in the New England Journal of Medicine, or journals from The Lancet, BMJ, or JAMA.

N – Total number of participants in each arm of the study.

^aThese studies compared the standard of care and sham treatments to integrative health, accounting for placebo effects.

^bThis study had 3 arms: one group received conventional care, one group received conventional care and sham acupuncture and the third group received conventional care and acupuncture.

^cThis study had 3 arms: one group received conventional care, one group received conventional care in addition to electroacupuncture at one site on the body and the third received conventional care with electroacupuncture at another site on the body.

^dThe outcomes reported in these two publications are from the same trial at two different follow-up time points.

4. Conclusion

Emerging evidence supports the integration of complementary and lifestyle-based interventions into conventional care as a viable strategy for managing chronic disease and enhancing patient outcomes. Findings from recently published high-quality clinical trials presented here show the promise of integrative health. Given the widespread global use of complementary and lifestyle approaches, advancing research in integrative healthcare may support the development of more sustainable, evidence-informed health systems worldwide.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author(s) used Microsoft Copilot (Free version) in order to lightly edit the manuscript. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Data availability

The data for this article consists of bibliographic references, which are included in the References section.

Acknowledgements

We would like to acknowledge Anupama Jithesh, Projects Coordinator, Weill Cornell Medicine-Qatar, for her assistance in retrieving relevant published articles for this commentary. We also thank Philippe Piccardi, Scientific and Education Content Specialist, and Hidenori Miyagawa, Visual Design Specialist, both at Weill Cornell Medicine-Qatar, for their assistance with manuscript editing and visual asset consultation, respectively.

References

- World Health Organization (WHO): **Global Health Observatory: Global health estimates: Leading causes of DALYs**. (Accessed Apr 28, 2024). [Reference Source](#)
- Global incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years (DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries and territories and 811 subnational locations, 1990-2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet*. 2024; **403**(10440): 2133–2161.
- Safer DJ: **Overprescribed Medications for US Adults: Four Major Examples**. *J. Clin. Med. Res.* 2019; **11**(9): 617–622. [Publisher Full Text](#)
- World Health Organization (WHO): **Traditional, complementary, and integrative medicine**. (Accessed May 7, 2025). [Reference Source](#)
- John NA, John J, Tarnikanti M, et al.: **Implications of lifestyle medicine in medical practice**. *J. Family Med. Prim. Care*. 2023; **12**(2): 208–212.
- National Center for Complementary and Integrative Health: **Complementary, Alternative, or Integrative Health: What's In a Name?** 2021. (Accessed Jan 15, 2025). [Reference Source](#)
- McCaffrey AM, Pugh GF, O'Connor BB: **Understanding patient preference for integrative medical care: Results from patient focus groups**. *J. Gen. Intern. Med.* 2007; **22**(11): 1500–1505.
- Hawkes RE, Cameron E, Cotterill S, et al.: **The NHS Diabetes Prevention Programme: an observational study of service delivery and patient experience**. *BMC Health Serv. Res.* 2020; **20**(1): 1098.
- Klein J, Brauer P, Royall D, et al.: **Patient experiences of a lifestyle program for metabolic syndrome offered in family medicine clinics: a mixed methods study**. *BMC Fam. Pract.* 2018; **19**(1): 148.
- Richardson K, Petukhova R, Hughes S, et al.: **The acceptability of lifestyle medicine for the treatment of mental illness: perspectives of people with and without lived experience of mental illness**. *BMC Public Health*. 2024; **24**(1): 171.
- Srivastava SB: **Polypharmacy, Unintended Consequences, and Impact of Lifestyle Medicine**. *Am. J. Lifestyle Med.* 2023; **18**(1): 54–57.
- Herman PM, Poindexter BL, Witt CM, et al.: **Are complementary therapies and integrative care cost-effective? A systematic review of economic evaluations**. *BMJ Open*. 2012; **2**(5): e001046.
- Sommers E, D'Amico S, Goldstein L, et al.: **Integrative Approaches to Pediatric Chronic Pain in an Urban Safety-Net Hospital: Cost Savings, Clinical Benefits, and Safety**. *J. Integr. Complement. Med.* 2022; **28**(5): 445–453.
- Lippman D, Stump M, Veazey E, et al.: **Foundations of Lifestyle Medicine and its Evolution**. *Mayo Clin. Proc. Innov. Qual. Outcomes*. 2024; **8**(1): 97–111.
- Pan CX, Morrison RS, Ness J, et al.: **Complementary and alternative medicine in the management of pain, dyspnea, and nausea and vomiting near the end of Life: A systematic review**. *J. Pain Symptom Manag.* 2000; **20**(5): 374–387.
- Patel P, Madrid M: **Successfully Integrating Lifestyle Medicine Into Health Systems**. *Am. J. Lifestyle Med.* 2021; **15**(5): 510–513.
- Miller DW, Roseen EJ, Stone JAM, et al.: **Incorporating Acupuncture Into American Healthcare: Initiating a Discussion on Implementation Science, the Status of the Field, and Stakeholder Considerations**. *Glob. Adv. Health Med.* 2021; **10**: 21649561211042574.
- Jonas W, Mehta D: **Integrative Care Interest and Offerings Grow**. *NEJM Catalyst*. 4(9).
- American Medical Association: **What is value-based care?** Jul 30, 2025. Retrieved Dec 15, 2025. [Reference Source](#)
- Pathak N, Pollard KJ, McKinney A: **Lifestyle Medicine Interventions for Personal and Planetary Health: The Urgent Need for Action**. *Am. J. Lifestyle Med.* 2022; **16**(5): 589–593.
- Cramer H: **Sustainable Health Care: How Traditional, Complementary, and Integrative Medicine Could Lead the Shift Towards Eco-Friendly Practices**. *J. Integr. Complement. Med.* 2024; **30**(10): 907–909.
- Gould RL, McDermott CJ, Thompson BJ, et al.: **Acceptance and Commitment Therapy plus usual care for improving quality of life in people with motor neuron disease (COMMEND): a multicentre, parallel, randomised controlled trial in the UK**. *Lancet*. 2024; **403**(10442): 2381–2394.
- Li K, Xu S, Wang R, et al.: **Electroacupuncture for motor dysfunction and constipation in patients with Parkinson's disease: a randomised controlled multi-centre trial**. *EclinicalMedicine*. 2023; **56**: 101814. [Publisher Full Text](#)
- Yan M, Fan J, Liu X, et al.: **Acupuncture and Sleep Quality Among Patients With Parkinson Disease: A Randomized Clinical Trial**.

- JAMA Netw. Open.* 2024; **7**(6): e2417862-e.
[Publisher Full Text](#)
25. Wolf S, Seiffer B, Zeibig J-M, *et al.*: **A transdiagnostic group exercise intervention for mental health outpatients in Germany (ImPuls): results of a pragmatic, multisite, block-randomised, phase 3 controlled trial.** *Lancet Psychiatry.* 2024; **11**(6): 417–430.
 26. Yin X, Li W, Liang T, *et al.*: **Effect of Electroacupuncture on Insomnia in Patients With Depression: A Randomized Clinical Trial.** *JAMA Netw. Open.* 2022; **5**(7): e2220563.
[Publisher Full Text](#)
 27. Yang J-W, Shao J-K, Wang Y, *et al.*: **Effect of acupuncture on postoperative ileus after laparoscopic elective colorectal surgery: A prospective, randomised, controlled trial.** *EClinicalMedicine.* 2022; **49**: 101472.
 28. Bao C, Wu L, Wang D, *et al.*: **Acupuncture improves the symptoms, intestinal microbiota, and inflammation of patients with mild to moderate Crohn's disease: A randomized controlled trial.** *EClinicalMedicine.* 2022; **45**: 101300.
 29. Malfliet A, De Baets L, Bilterys T, *et al.*: **Cognitive Behavioral Therapy for Insomnia in Pain Management for Nonspecific Chronic Spinal Pain: A Randomized Clinical Trial.** *JAMA Netw. Open.* 2024; **7**(8): e2425856-e.
[Publisher Full Text](#)
 30. Tu JF, Cao Y, Wang LQ, *et al.*: **Effect of Adjunctive Acupuncture on Pain Relief Among Emergency Department Patients With Acute Renal Colic Due to Urolithiasis: A Randomized Clinical Trial.** *JAMA Netw. Open.* 2022; **5**(8): e2225735.
[Publisher Full Text](#)
 31. Jensen SBK, Sørensen V, Sandsdal RM, *et al.*: **Bone Health After Exercise Alone, GLP-1 Receptor Agonist Treatment, or Combination Treatment: A Secondary Analysis of a Randomized Clinical Trial.** *JAMA Netw. Open.* 2024; **7**(6): e2416775.
[Publisher Full Text](#)
 32. Lundgren JR, Janus C, Jensen SBK, *et al.*: **Healthy Weight Loss Maintenance with Exercise, Liraglutide, or Both Combined.** *N. Engl. J. Med.* 2021; **384**(18): 1719–1730.
 33. Jensen SBK, Blond MB, Sandsdal RM, *et al.*: **Healthy weight loss maintenance with exercise, GLP-1 receptor agonist, or both combined followed by one year without treatment: a post-treatment analysis of a randomised placebo-controlled trial.** *EClinicalMedicine.* 2024; **69**: 102475.
 34. MacDonald CS, Nielsen SM, Bjørner J, *et al.*: **One-year intensive lifestyle intervention and improvements in health-related quality of life and mental health in persons with type 2 diabetes: a secondary analysis of the U-TURN randomized controlled trial.** *BMJ Open Diabetes Res. Care.* 2021; **9**(1).
[Publisher Full Text](#)
 35. Lee W-J, Peng L-N, Lin C-H, *et al.*: **Effects of incorporating multidomain interventions into integrated primary care on quality of life: a randomised controlled trial.** *Lancet Healthy Longev.* 2021; **2**(11): e712–e723.
[Publisher Full Text](#)

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research