

Effectiveness of Resident Physicians as Triage Liaison Providers in an Academic Emergency Department



Victoria Weston MD¹, Amer Aldeen MD², Stephanie Gravenor BS¹, Sushil Jain MD MBA¹, Michael Schmidt MD¹, Sanjeev Malik MD¹

¹Northwestern University, Feinberg School of Medicine, Department of Emergency Medicine.

²Presence St. Joseph Medical Center, Emergency Medicine Physicians, Ltd.

Background

- ED overcrowding lowers patient satisfaction, door-to-physician time (DTP), and ED length of stay (LOS)
- Triage liaison providers (TLP) reduce DTP and left without being seen (LWBS)
- Prior TLP studies → advanced practice providers and attending physicians
- No published studies on resident physicians as TLP

Objective


 Resident vs. attending TLP → operational performance, patient satisfaction, and costeffectiveness


Methods

- Retrospective, observational cohort study
- Urban, academic ED, 88,000 annual visits
- Staff: 28 attending physicians, 50 resident physicians
- PGY-3/4 resident or attending MDs 11:30-19:30 weekdays
- TLP days (10/2013-1/2014) vs. baseline (10/2011-1/2012)
- Primary outcomes: differences in median ED LOS, median DTP, proportion of LWBS, proportion of "very good" overall patient satisfaction scores, and cost effectiveness
- Proportions: described with 95% confidence intervals
- Medians: measured with interquartile ranges, differences with the t-test and Mann-Whitney U test
- Cost effectiveness: calculated with annual revenue generated through LWBS capture offset by TLP cost
- Revenue: calculated using both physician-based collections and hospital-based collections


Results


	Baseline (no TLP)	Attending TLP	Resident TLP
Study Days	92	48	29
Patient Visits	19,298	10,814	6,683











Conclusions

- Resident TLP improved DTP times and LWBS > attending TLP or historical control
- Attending TLP improved patient satisfaction and the median LOS of admitted patients > resident TLP
- Physician-based collections: resident TLP > attending
 TLP
- Hospital-based collections: both resident and attending TLP are cost effective
- Residents appear to be more cost effective as TLP overall

Limitations:

- Design limitations as retrospective cohort study
- Single institution
- Academic ED with nearby pediatric ED → mainly adult patients
- PGY3 and PGY4 residents only
- Utilization limited to an academic ED with resident physicians

References:

- 1. Wiler JL, Gentle C, Halfpenny JM, et al. Optimizing emergency department front-end operations. Ann Emerg Med. 2010; 55:142–60.
- 2. Sprivulis PC, Da Silva JA, Jacobs IG, Frazer ARL, Jelinek GA. The association between hospital overcrowding and mortality among patients admitted via Western Australian emergency departments. Med J Austral. 2006; 184:208–12.
- 3. Richardson DB. Increase in patient mortality at 10 days associated with emergency department overcrowding. Med J Austral. 2006; 184:213–6.
- 4. Asplin BR, Magid DJ, Rhodes KV, Solberg LI, Lurie N, Camargo CA Jr. A conceptual model of emergency department crowding. Ann Emerg Med. 2003; 42: 173–80.
 - Asaro PV, Lewis LM, Boxerman SB. The impact of input and output factors on emergency department throughput. Acad Emerg Med. 2007; 14:235–42.