

GEDI WISE: Geriatric-Specific Assessment and Intervention in the Emergency Department is Associated with Change in Disposition

Amer Aldeen MD, Scott Dresden, MD, MS, Danielle McCarthy MD, MS, Stephanie Gravenor, D. Mark Courtney, MD, MSCI

Northwestern University, Chicago, IL
Department of Emergency Medicine



Background

- Hospitalization for geriatric patients carries risk of impaired functional status, delirium, and mortality
- To improve care and decrease cost for older adults specially trained nurse liaisons perform Geriatric ED innovation (GEDI) consultation consisting of:
 - Assessment of delirium, dementia, gait stability, functional status, and caregiver strain
 - Care coordination involving ED, social work, pharmacy, physical therapy, geriatrics and PCP
 - Referral for home health and social services

Objectives

- Describe disposition decisions following geriatric nursing assessment and care coordination for geriatric patients in the ED.
 - Admission rate for patients with uncertain disposition prior to GEDI evaluation
 - Changes in disposition after GEDI evaluation

Methods

Study design: Prospective observational cohort study in a primarily adult urban ED with 85k visits/year.

Dates: July 2013 – February 2014, Monday-Friday 9a-8p

Inclusion criteria: ED patients age 65+ who received GEDI consult triggered by Identification of Seniors at Risk (ISAR) score of >2, or ED team request.

Measurements: GEDI nurse-liaison asked ED providers their anticipated disposition for enrolled patients

As part of GEDI consult the following assessments were recorded: Timed up and go, short portable mental status questionnaire, confusion assessment method, Katz activities of daily living, Beers criteria, and modified caregiver strain index


Results

- 766 GEDI patients eligible
- 640 with anticipated dispo data
- Overall admission rate for geriatric patients 60%

Table 1: Patient Characteristics by Anticipated Disposition

Characteristic, n (%)	Likely Admit n=145	Likely Discharge N=255	Uncertain Dispo n=240
Age - Median (IQR)	79 (73-86)	78 (72-85)	80 (73-86)
Female	92 (63)	159 (62)	156 (65)
Race			
White	92 (63)	146 (57)	147 (61)
African American	36 (25)	85 (33)	65 (27)
ESI - Median (IQR)	2 (2-3)	3 (2-3)	2 (2-3)
ISAR - Median (IQR)	4 (3-4)	3 (1-3)	3 (1-3)

Figure 1: Final Disposition by Anticipated Disposition



Results

Table 2: Number and percent of positive GEDI assessments by anticipated and final disposition

+ Assessment, n (%)	Likely Admit			Likely Discharge		
	Admit n=117	DC n=28	p	Admit n=37	DC n=218	p
Final Dispo						
Delirium	9 (8%)	2 (7%)	0.89	1 (3%)	2 (1%)	0.35
Dementia	35 (31%)	8 (29%)	0.805	6 (18%)	39 (18%)	0.95
Impaired Mobility	108 (92%)	19 (68%)	<0.01	31 (84%)	158 (72%)	0.15
Intermediate-Low Independence	49 (54%)	13 (54%)	0.98	15 (50%)	80 (44%)	0.95
Pharmacy Risk	38 (60%)	7 (58%)	0.90	11 (50%)	67 (49%)	0.57
Caregiver Strain	6 (5%)	0 (0%)	0.22	5 (14%)	12 (6%)	0.07

Limitations

- Comparison of uncertain disposition patients to general Geriatric ED population may be subject to selection bias
- Unable to definitively determine roll of GEDI consult in disposition decision

Conclusions

- Lower admission rate for GEDI patients with initial uncertain disposition compared to all geriatric patients (43% vs. 60%)
- 10% of disposition decisions are changed after GEDI consult
 - Discharge may reflect benefits of care coordination
 - Admission may reflect recognition of unmet needs

Funding

The project described was supported by Grant Number 1C1CMS331055 the Department of Health and Human Services, Centers for Medicare & Medicaid Services.

Its contents are solely the responsibility of the authors and have not been approved by the Department of Health and Human Services, Centers for Medicare & Medicaid Services.