Final Exam Overview

Tiandong Wang

Department of Statistics Texas A&M University

Spring 2021

Tips for Final

To enhance **readability** of your answers:

- Try to avoid pencils; Black pen is the easiest to read.
- Write big; Scrawls are painful to read.
- Mark your response to every part of the questions clearly.
- Check your scanned copy before submission (you could try mobile apps like Adobe Scan).

Time management:

- You'll have 165 mins for the final, as the original duration was 2 hrs. With an additional 45 mins, you should be able to complete scanning, checking and uploading.
- Check out the cover page of the final exam on eCampus.
- Calculators are not necessary for the exam.



Your Feedback Matters!

Please fill out the course evaluation forms via

```
https://yourvoice.tamu.edu/Menu/
Student-Course-Evaluations/AEFIS
```

- AEFIS closes on May 2.
- Current response rate: 73%. Target rate: 70%.

Data reduction

- Sufficient statistic
- How to find a sufficient statistic
 - (Neyman-Fisher) Factorization theorem. Theorem 6.2.6
 - Exponential family (Theorem 6.2.10)
 - Transformation
- Minimal Sufficient Statistics
 - Theorem 6.2.13
 - Exponential family
- Complete Statistics
 - Definition?
 - Exponential family



Methods for finding MLEs

- Look at a graph of the likelihood function.
- Calculus method: Take partial derivative of log-likelihood with respect to each parameter and solve the likelihood equations.
- Invariance property of MLEs: MLE of a function of θ is the function applied to MLE of θ (Theorem 7.2.10)

Important Concepts in Bayes Estimation

- prior distribution of θ : $\theta \sim \pi(\theta)$. Conjugate priors: The posterior is in the same family as the prior regardless of the data observed.
- posterior distribution of θ : $\pi(\theta|\mathbf{x}) = f(\mathbf{x}|\theta)\pi(\theta)/m(\mathbf{x})$
- marginal distribution of **x**: $m(\mathbf{x}) = \int f(\mathbf{x}|\theta)\pi(\theta)d\theta$
- posterior mean of θ : $E(\theta|\mathbf{X}) = \int \theta \pi(\theta|\mathbf{x}) d\theta$ (Bayes estimator of θ)

Methods of Evaluating Estimators

- Mean squared error: MSE equals estimator variance plus estimator squared bias.
- Best Unbiased Estimators (UMVUE) (Definition 7.3.7)
- Cramér-Rao Lower Bound (Theorem 7.3.9)
- Score Function and Fisher Information (one parameter/multiple parameters), exponential family. (Lemma 7.3.11)
- Attainment of C-R Bound (Corollary 7.3.15), exponential family.
- Sufficiency and UMVUE: Rao-Blackwell theorem (Theorem 7.3.17 and Theorem 7.3.23)



Loss Function Optimality

- Loss function
- Risk function
- Bayesian risk function
- Bayes rule (squared loss and absolute loss)

Hypothesis testing

- Power function
- Type I, II error
- Size/Level of a test
- LRT; Rejection regions; p-values.
- ullet Critical value for a given size lpha/Sample Size Calculation
- Unbiased test
- Uniformly most powerful test(UMP level α test)
 - Neyman-Pearson lemma (simple test).
 - Neyman-Pearson based on sufficient statistic (simple test).
 - Neyman-Pearson lemma for randomized tests.
 - Karlin-Rubin Theorem for MLR family (composite test). Examples of MLR family: exponential with non decreasing $w(\theta)$.
- Multiple hypothesis testing.



How to construct confidence intervals

Method 1: By inverting the acceptance region of tests;

Method 2: Using pivotal quantities.

 $Q(\mathbf{X}, \theta)$ is independent of θ

- How to find pivotal quantity?
- Find a,b such that $P_{\theta}(a \leq Q(X, \theta) \leq b) = 1 \alpha$
- Pivoting the CDF of sufficient statistic



Asymptotics

- Consistency: definition, how to check consistency? (MLE is consistent)
- Asymptotic normality (CLT, Delta method)
- Asymptotic efficiency, i.e., the asymptotic variance of T_n achieves the Cramer-Rao lower bound). Review: how to calculate Cramér-Rao lower bound?
- Asymptotic relative efficiency. Definition?
- Large sample hypothesis tests; Wilks' theorem;
 Contingency tables.

- Thanks a lot for your support throughout the semester, and
- Good luck with everything in the future!!!