
A type theory for cellular spaces

Louise Leclerc
Samuel Mimram

January 13, 2026

Main Idea

- Simplicial Type Theory (*STT*)
(E. Riehl, M. Shulman, D. Gratzer, J. Weinberger, U. Buchholtz,)

Main Idea

- Simplicial Type Theory (*STT*)
(E. Riehl, M. Shulman, D. Gratzer, J. Weinberger, U. Buchholtz,)
 - $\llbracket \mathcal{U} \rrbracket = \mathbf{Psh}(\Delta)$

Main Idea

- Simplicial Type Theory (*STT*)
(E. Riehl, M. Shulman, D. Gratzer, J. Weinberger, U. Buchholtz,)
 - $\llbracket \mathcal{U} \rrbracket = \mathbf{Psh}(\Delta)$
 - $(\infty, 1)$ -categories = Complete Segal Spaces.

Main Idea

- Simplicial Type Theory (*STT*)
(E. Riehl, M. Shulman, D. Gratzer, J. Weinberger, U. Buchholtz,)
 - $\llbracket \mathcal{U} \rrbracket = \mathbf{Psh}(\Delta)$
 - $(\infty, 1)$ -categories = Complete Segal Spaces.
- Cellular Type Theory (*CellTT*)
(Using C. Rezk Θ -Spaces + F. Loubaton Thesis)

Main Idea


- Simplicial Type Theory (*STT*)
(E. Riehl, M. Shulman, D. Gratzer, J. Weinberger, U. Buchholtz,)
 - $\llbracket \mathcal{U} \rrbracket = \mathbf{Psh}(\Delta)$
 - $(\infty, 1)$ -categories = Complete Segal Spaces.
- Cellular Type Theory (*CellTT*)
(Using C. Rezk Θ -Spaces + F. Loubaton Thesis)
 - $\llbracket \mathcal{U} \rrbracket = \mathbf{Psh}(\Theta)$

Main Idea

- Simplicial Type Theory (*STT*)
(E. Riehl, M. Shulman, D. Gratzer, J. Weinberger, U. Buchholtz,)
 - $\llbracket \mathcal{U} \rrbracket = \mathbf{Psh}(\Delta)$
 - $(\infty, 1)$ -categories = Complete Segal Spaces.
- Cellular Type Theory (*CellTT*)
(Using C. Rezk Θ -Spaces + F. Loubaton Thesis)
 - $\llbracket \mathcal{U} \rrbracket = \mathbf{Psh}(\Theta)$
 - (∞, ω) -categories = Θ -Spaces.


The interval

We add to the syntax a collection of shapes



The interval

We add to the syntax a collection of shapes



Simplicial types

S shape, and X type:

$S \rightarrow X$ is a type

Simplicial types

S shape, and X type:

$S \rightarrow X$ is a type

$$\text{hom}_X(x, y) = \sum_{f:2 \rightarrow X} (f(0) = x) \times (f(1) = y)$$

Simplicial types

S shape, and X type:

$S \rightarrow X$ is a type

$$\mathrm{hom}_X(x, y) = \sum_{f:2 \rightarrow X} (f(0) = x) \times (f(1) = y)$$

$\forall n, \quad X_n := \Delta_n \rightarrow X$ is a type

Simplicial types

S shape, and X type:

$S \rightarrow X$ is a type

$$\text{hom}_X(x, y) = \sum_{f:2 \rightarrow X} (f(0) = x) \times (f(1) = y)$$

$\forall n, \quad X_n := \Delta_n \rightarrow X$ is a type

So each type give rise to a *simplicial type* !

Segal types

When does (X_n) satisfies the Segal condition ?

Segal types

When does (X_n) satisfies the Segal condition ?

$$(\Delta^n \rightarrow X) \xrightarrow{\sim} (\Lambda_{n,k} \rightarrow X)$$

Segal types

When does (X_n) satisfies the Segal condition ?

$$(\Delta^n \rightarrow X) \xrightarrow{\sim} (\Lambda_{n,k} \rightarrow X)$$

But $(\Delta^n \rightarrow X)$ is also simplicial !

Segal types

When does (X_n) satisfies the Segal condition ?

$$(\Delta^n \rightarrow X) \xrightarrow{\sim} (\Lambda_{n,k} \rightarrow X)$$

But $(\Delta^n \rightarrow X)$ is also simplicial !

$$(\Delta^2 \rightarrow X) \xrightarrow{\sim} (\Lambda_2 \rightarrow X) \quad \text{suffices !}$$

Complete types

By horn filling: A Segal admits composites $g \circ f$.

Complete types

By horn filling: A Segal admits composites $g \circ f$.

$$\text{iso}(f) :\equiv \left(\sum_{g:\text{hom}_A(y,x)} g \circ f = \text{id}_x \right) \times \left(\sum_{h:\text{hom}_A(y,x)} f \circ h = \text{id}_y \right)$$

Complete types

By horn filling: A Segal admits composites $g \circ f$.

$$\text{iso}(f) := \left(\sum_{g:\text{hom}_A(y,x)} g \circ f = \text{id}_x \right) \times \left(\sum_{h:\text{hom}_A(y,x)} f \circ h = \text{id}_y \right)$$

$$x \simeq_A y := \sum_{f:\text{hom}_A(x,y)} \text{iso}(f)$$

Complete types

By horn filling: A Segal admits composites $g \circ f$.

$$\text{isiso}(f) := \left(\sum_{g:\text{hom}_A(y,x)} g \circ f = \text{id}_x \right) \times \left(\sum_{h:\text{hom}_A(y,x)} f \circ h = \text{id}_y \right)$$

$$x \simeq_A y := \sum_{f:\text{hom}_A(x,y)} \text{isiso}(f)$$

$$\text{idtoiso} : \prod_{x,y:A} x =_A y \rightarrow x \simeq_A y$$

Complete types

By horn filling: A Segal admits composites $g \circ f$.

$$\text{isiso}(f) := \left(\sum_{g:\text{hom}_A(y,x)} g \circ f = \text{id}_x \right) \times \left(\sum_{h:\text{hom}_A(y,x)} f \circ h = \text{id}_y \right)$$


$$x \simeq_A y := \sum_{f:\text{hom}_A(x,y)} \text{isiso}(f)$$

$$\text{idtoiso} : \prod_{x,y:A} x =_A y \rightarrow x \simeq_A y$$

Reminiscent to Rezk-completes/univalent categories in HoTT.

The category Θ

- Objects: Pasting schemes.




Some objects of Θ

•

$$\mathcal{O}_0 = *$$


Some objects of Θ






$$\mathcal{O}_0 = *$$

$$\mathcal{O}_1 = [1]$$





Some objects of Θ



Some objects of Θ


 $\mathcal{O}_0 = *$

 $\mathcal{O}_1 = [1]$

 \mathcal{O}_2

 \mathcal{O}_3

Some objects of Θ


 $\mathcal{O}_0 = *$

 $\mathcal{O}_1 = [1]$

 \mathcal{O}_2

 \mathcal{O}_3

 $[n]$

A combinatorial description of Θ

- Objects are lists of objects.

$$\bullet \xrightarrow{P_1} \bullet \xrightarrow{P_2} \bullet \cdots \cdots \cdots \bullet \xrightarrow{P_n} \bullet$$

A combinatorial description of Θ

- Objects are lists of objects.

$$\bullet \xrightarrow{P_1} \bullet \xrightarrow{P_2} \bullet \cdots \cdots \cdots \bullet \xrightarrow{P_n} \bullet$$



$$* = []$$

$$\mathcal{O}_1 = [*]$$

$$\mathcal{O}_2 = [\mathcal{O}_1] = [[*]]$$

$$\bullet \longrightarrow \bullet \cdots \cdots \cdots \bullet \longrightarrow \bullet$$


$$[n] = [*, *, \dots, *]$$

A combinatorial description of Θ

- Objects are lists of objects.

$$\bullet \xrightarrow{P_1} \bullet \xrightarrow{P_2} \bullet \cdots \cdots \cdots \bullet \xrightarrow{P_n} \bullet$$

- Morphisms:



Hom Types in STT

In STT:

$$\text{hom}_A(x, y) = \sum_{f: I \rightarrow A} (f\ 0 = x) \times (f\ 1 = y)$$

$$x \text{ ————— } y$$

Hom Types in STT

In STT:

$$\text{hom}_A(x, y) = \sum_{f: I \rightarrow A} (f\ 0 = x) \times (f\ 1 = y)$$

$$x \text{ ————— } y$$

Then:

$$\text{hom}_{\text{hom}_A(x, y)}(f, g) = \sum_{H: I^2 \rightarrow A} \dots$$

$$\begin{array}{ccc} x & \text{— } f \text{ —} & y \\ \parallel & & \parallel \\ x & \text{— } g \text{ —} & y \end{array}$$

Does not generalize to Θ

$$I = \mathcal{Y}([1])$$

Does not generalize to Θ

$$I = \mathcal{Y}([1])$$

2-cells of I^2 are pairs (x, y) of 2-cells of I .

Does not generalize to Θ

$$I = \mathcal{Y}([1])$$

2-cells of I^2 are pairs (x, y) of 2-cells of I .

x, y are invertible $\Rightarrow (x, y)$ too.


Does not generalize to Θ

$$1 = \mathfrak{J}([1])$$

2-cells of \mathbf{l}^2 are pairs (x, y) of 2-cells of \mathbf{l} .

x, y are invertible $\Rightarrow (x, y)$ too.

Hence \mathcal{I}^2 is 1-categorical



Does not generalize to Θ


$$1 = \mathfrak{J}([1])$$

2-cells of \mathbf{l}^2 are pairs (x, y) of 2-cells of \mathbf{l} .

x, y are invertible $\Rightarrow (x, y)$ too.


Hence \mathcal{I}^2 is 1-categorical

$$\text{hom}_A(x, y) \neq \sum_{f: I \rightarrow A} (f\,0 = x) \times (f\,1 = y)$$




Another approach

Workaround:



Another approach

Workaround:



Two subgoals:

- Defining a suspension \$.
- Postulating the adjunction.

$(\$ \dashv \text{hom})$ adjunction

$$(A \rightarrow \text{hom}_B(x, y)) \stackrel{?}{\simeq} (\$A \rightarrow_{\bullet\bullet} (B, x, y))$$

$(\$ \dashv \text{hom})$ adjunction

$$\underbrace{(A \rightarrow \text{hom}_B(x, y))}_{\text{internal hom}} \neq \underbrace{(\$A \rightarrow_{\bullet\bullet} (B, x, y))}_{\text{internal hom}}$$

$(\$ \dashv \text{hom})$ adjunction

$$\mathfrak{b}(A \rightarrow \text{hom}_B(x, y)) = \mathfrak{b}(\$A \rightarrow \bullet\bullet (B, x, y))$$

The \flat modality


$$A :: \mathcal{U} \quad \rightsquigarrow \quad \llbracket A \rrbracket \in \mathbf{Psh}(\Theta)$$

$$\flat A : \mathcal{U} \quad \rightsquigarrow \quad \llbracket \flat A \rrbracket = \llbracket A \rrbracket(*)$$

Semantic of \flat

$$\begin{array}{ccc} & \delta & \\ \mathcal{S} & \xrightarrow{\quad} & \mathbf{Psh}(\Theta) \\ & \xleftarrow{F \mapsto F(*)} & \\ & \perp & \end{array}$$

Semantic of \flat



Crisp Type Theory

Why Crisp Type Theory ?

Crisp Type Theory

Why Crisp Type Theory ?

Because \flat is not “continuous”.

Crisp Type Theory

Why Crisp Type Theory ?

Because \flat is not “continuous”.

Two kinds of hypothesis:

continuous $X : \mathcal{U}, x : X$

Crisp Type Theory

Why Crisp Type Theory ?

Because \flat is not “continuous”.

Two kinds of hypothesis:

continuous

$X : \mathcal{U}, x : X$

crisp

$X :: \mathcal{U}, x :: X$

Crisp Type Theory

Why Crisp Type Theory ?

Because \flat is not “continuous”.

Two kinds of hypothesis:

continuous	$X : \mathcal{U}, x : X$
crisp	$X :: \mathcal{U}, x :: X$

$$\frac{\Gamma | \cdot \vdash X : \mathcal{U}}{\Gamma | \cdot \vdash \flat X : \mathcal{U}}$$

CellTT

$$\text{CellTT} = \text{HoTT} + \underbrace{\text{Idempotent comodality}}_{\text{Crisp Type Theory}} + \text{Axioms}$$

Pasting Schemes

pasting schemes: $\left\{ \begin{array}{l} \text{PS} : \text{Set} \\ [] : \text{Array PS} \rightarrow \text{PS} \end{array} \right.$

Pasting Schemes

pasting schemes: $\left\{ \begin{array}{l} \text{PS} : \text{Set} \\ [] : \text{Array PS} \rightarrow \text{PS} \end{array} \right.$

morphisms: $P \rightarrow_{\text{PS}} Q : \text{Set} \quad (P, Q : \text{PS})$

Pasting Schemes

pasting schemes: $\left\{ \begin{array}{l} \text{PS} : \text{Set} \\ [] : \text{Array PS} \rightarrow \text{PS} \end{array} \right.$

morphisms: $P \rightarrow_{\text{PS}} Q : \text{Set} \quad (P, Q : \text{PS})$

suspension: $\left\{ \begin{array}{ll} \$: \text{PS} & \rightarrow \text{PS} \\ P & \rightarrow [P] \end{array} \right.$

Pasting Schemes

pasting schemes: $\begin{cases} \text{PS} : \text{Set} \\ [] : \text{Array PS} \rightarrow \text{PS} \end{cases}$

morphisms: $P \rightarrow_{\text{PS}} Q : \text{Set} \quad (P, Q : \text{PS})$

suspension: $\begin{cases} \$: \text{PS} \rightarrow \text{PS} \\ P \rightarrow [P] \end{cases}$

$$* = [] \quad [n] = [*, \dots, *]$$

$$\mathcal{O}_n = \$^n [] = [[\dots [] \dots]]$$

Yoneda Embedding

Yoneda: $\mathbf{y} : \mathbf{PS} \rightarrow \mathcal{U}$

Yoneda Embedding

Yoneda: $\mathcal{Y} : \mathbf{PS} \rightarrow \mathcal{U}$

$$X :: \mathcal{U} \rightsquigarrow X_P \equiv \mathcal{Y}(P) \rightarrow X$$

Yoneda Embedding

Yoneda: $\mathcal{Y} : \mathbf{PS} \rightarrow \mathcal{U}$

$$X :: \mathcal{U} \rightsquigarrow X_P \equiv \mathcal{Y}(P) \rightarrow X$$

$$f :: X \rightarrow Y \rightsquigarrow f_P : X_P \rightarrow Y_P$$

Yoneda Embedding

$$\text{Yoneda: } \mathcal{Y} : \mathbf{PS} \rightarrow \mathcal{U}$$

$$X :: \mathcal{U} \rightsquigarrow X_P \equiv \mathcal{Y}(\mathcal{Y}(P) \rightarrow X)$$

$$f :: X \rightarrow Y \rightsquigarrow f_P : X_P \rightarrow Y_P$$

$$\sigma : P \rightarrow_{\mathbf{PS}} Q \rightsquigarrow \sigma^* : X_Q \rightarrow X_P$$

Some Axioms

- Equivalences are pointwise

$$\left(\prod_{P:PS} \text{is-equiv}(f_P) \right) \rightarrow \text{is-equiv}(f)$$

Some Axioms

- Equivalences are pointwise

$$\left(\prod_{P:PS} \text{is-equiv}(f_P) \right) \rightarrow \text{is-equiv}(f)$$

- \mathcal{Y} is fully faithful
- $(-)_P$ preserves colimits
- $(\sum_{P:PS} \sum_{c:X_P} \mathcal{Y}P) \rightarrow X$ is an effective epi ((-1) -truncated)

Some Axioms

- Equivalences are pointwise

$$\left(\prod_{P:\mathcal{P}S} \text{is-equiv}(f_P) \right) \rightarrow \text{is-equiv}(f)$$

- \mathcal{Y} is fully faithful
- $(-)_P$ preserves colimits
- $(\sum_{P:\mathcal{P}S} \sum_{c:X_P} \mathcal{Y}P) \rightarrow X$ is an effective epi ((-1) -truncated)

Corollary 5.1.6.11. *Let \mathcal{C} be an ∞ -category which admits small colimits. Let S be a small simplicial set and $F : \mathcal{P}(S) \rightarrow \mathcal{C}$ a colimit preserving functor. Then F is an equivalence if and only if the following conditions are satisfied:*

- (1) *The composition $f = F \circ j : S \rightarrow \mathcal{C}$ is fully faithful.*
- (2) *For every vertex $s \in S$, the object $f(s) \in \mathcal{C}$ is completely compact.*
- (3) *The set of objects $\{f(s) : s \in S_0\}$ generates \mathcal{C} under colimits.*

Some Axioms

- Equivalences are pointwise

$$\left(\prod_{P:PS} \text{is-equiv}(f_P) \right) \rightarrow \text{is-equiv}(f)$$

- \mathcal{J} is fully faithful
- $(-)_P$ preserves colimits
- $(\sum_{P:PS} \sum_{c:X_P} \mathcal{J}P) \rightarrow X$ is an effective epi ((-1) -truncated)
- \flat -discreteness is *cellular discreteness*:

$$\text{is-equiv}(\flat X \rightarrow X) \leftrightarrow \prod_{P:PS} \text{is-equiv}(X \rightarrow (\mathcal{J}(P) \rightarrow X))$$

Levelwiseness of level

Let $X :: \mathcal{U}$, then for any $n \geq -2$,

$$\text{is-}n\text{-type}(X) \leftrightarrow \prod_{P:\mathbf{PS}} \text{is-}n\text{-type}(X_P)$$

Levelwiseness of level

Let $X :: \mathcal{U}$, then for any $n \geq -2$,

$$\text{is-}n\text{-type}(X) \leftrightarrow \prod_{P:\mathbf{PS}} \text{is-}n\text{-type}(X_P)$$

- Case $n = -2$: $X \simeq \mathbb{1} \longleftrightarrow \prod_{P:\mathbf{PS}} (X_P \simeq \mathbb{1})$

Levelwiseness of level

Let $X :: \mathcal{U}$, then for any $n \geq -2$,

$$\text{is-}n\text{-type}(X) \leftrightarrow \prod_{P:\mathbf{PS}} \text{is-}n\text{-type}(X_P)$$

- Case $n = -2$: $X \simeq \mathbb{1} \iff \prod_{P:\mathbf{PS}} (X_P \simeq \mathbb{1})$
- Case $n \geq -1$: LR implication: $\|X\|_n \simeq \|bX\|_n$
HoTT book: $\text{is-}n\text{-type}(X) \leftrightarrow \text{is-equiv} \left(X^{S^{n+1}} \rightarrow X \right)$

Levelwiseness of level

Let $X :: \mathcal{U}$, then for any $n \geq -2$,

$$\text{is-}n\text{-type}(X) \leftrightarrow \prod_{P:PS} \text{is-}n\text{-type}(X_P)$$

- Case $n = -2$: $X \simeq \mathbb{1} \iff \prod_{P:PS} (X_P \simeq \mathbb{1})$
- Case $n \geq -1$: LR implication: $\|X\|_n \simeq \|bX\|_n$
HoTT book: $\text{is-}n\text{-type}(X) \leftrightarrow \text{is-equiv} \left(X^{S^{n+1}} \rightarrow X \right)$
 $\leftrightarrow \prod_{P:PS} \text{is-equiv} \left((X^{S^{n+1}})_P \rightarrow X_P \right)$

Levelwiseness of level

Let $X :: \mathcal{U}$, then for any $n \geq -2$,

$$\text{is-}n\text{-type}(X) \leftrightarrow \prod_{P:PS} \text{is-}n\text{-type}(X_P)$$

- Case $n = -2$: $X \simeq \mathbb{1} \longleftrightarrow \prod_{P:PS} (X_P \simeq \mathbb{1})$
- Case $n \geq -1$: LR implication: $\flat \|X\|_n \simeq \|\flat X\|_n$
 HoTT book: $\text{is-}n\text{-type}(X) \leftrightarrow \text{is-equiv}(X^{S^{n+1}} \rightarrow X)$
 $\leftrightarrow \prod_{P:PS} \text{is-equiv}((X^{S^{n+1}})_P \rightarrow X_P)$
 $(X^{S^{n+1}})_P \simeq \flat(\textstyle\bigwedge P \times S^{n+1} \rightarrow X) \simeq \flat(S^{n+1} \rightarrow X_P)$

Levelwiseness of level

Let $X :: \mathcal{U}$, then for any $n \geq -2$,

$$\text{is-}n\text{-type}(X) \leftrightarrow \prod_{P:PS} \text{is-}n\text{-type}(X_P)$$

- Case $n = -2$: $X \simeq \mathbb{1} \iff \prod_{P:PS} (X_P \simeq \mathbb{1})$
- Case $n \geq -1$: LR implication: $\flat \|X\|_n \simeq \|\flat X\|_n$
 HoTT book: $\text{is-}n\text{-type}(X) \leftrightarrow \text{is-equiv} (X^{\mathbb{S}^{n+1}} \rightarrow X)$
 $\iff \prod_{P:PS} \text{is-equiv} ((X^{\mathbb{S}^{n+1}})_P \rightarrow X_P)$
 $(X^{\mathbb{S}^{n+1}})_P \simeq \flat(\textstyle\bigvee P \times \mathbb{S}^{n+1} \rightarrow X) \simeq \flat(\mathbb{S}^{n+1} \rightarrow X_P)$
 $\flat(\mathbb{S}^{n+1} \rightarrow X_P) \simeq \flat X_P \simeq X_P.$

Suspension

We extend $\$$ to \mathcal{U} .


- $\downarrow(\$P) = \$(\downarrow P)$

Suspension

We extend $\$$ to \mathcal{U} .

- $\mathfrak{Y}(\$P) = \$(\mathfrak{Y}P)$
- If $P = [P_1, P_2, \dots, P_n]$:

$$(\$X)_P \cong \mathbb{1} + \left(\sum_{i: \text{Fin}_n} X_{P_i} \right) + \mathbb{1}$$



Hom Types



We postulate an adjunction

$$\flat(A \rightarrow \text{hom}_B(x, y)) \cong \flat(\$A \rightarrow_{\bullet\bullet} (B, x, y))$$

Cellular Realization

$$\langle [P_1, \dots, P_m] \rangle \equiv$$

colim



Segal Types

We define a map $\langle P \rangle \rightarrow \mathcal{L}P$

Segal Types

We define a map $\langle P \rangle \rightarrow \mathcal{L}P$

We define is-Segal:

$$\flat(\mathcal{L}P \rightarrow X) \xrightarrow{\sim} \flat(\langle P \rangle \rightarrow X)$$

Complete Types

We define a type E as the pushout

$$\begin{array}{ccc} \mathbb{N}[1] + \mathbb{N}[1] & \xrightarrow{\alpha, \beta} & \mathbb{N}[3] \\ \downarrow & & \downarrow \\ \mathbb{1} + \mathbb{1} & \xrightarrow{\quad \sqcup \quad} & E \end{array}$$

representing equivalences

Complete Types

We define a type E as the pushout

$$\begin{array}{ccc}
 \mathbb{N}[1] + \mathbb{N}[1] & \xrightarrow{\alpha, \beta} & \mathbb{N}[3] \\
 \downarrow & & \downarrow \\
 \mathbb{1} + \mathbb{1} & \xrightarrow{\quad \sqcup \quad} & E
 \end{array}$$

representing equivalences

$$E_{n+1} := \mathbb{N}^n E$$

Complete Types

We define a type E as the pushout

$$\begin{array}{ccc}
 \mathbb{N}[1] + \mathbb{N}[1] & \xrightarrow{\alpha, \beta} & \mathbb{N}[3] \\
 \downarrow & & \downarrow \\
 \mathbb{1} + \mathbb{1} & \xrightarrow{\quad \quad \quad} & E
 \end{array}
 \quad \sqcap$$

representing equivalences

$$E_{n+1} := \mathbb{N}^n E$$

We define is-complete:

$$\mathbb{N}(D_n \rightarrow X) \xrightarrow{\sim} \mathbb{N}(E_{n+1} \rightarrow X)$$

(∞, ω) -categories

As in Riehl-Shulman STT, there is a completeness condition.
 $\text{is-complete}(X)$

(∞, ω) -categories = complete Segal Types.

$$\text{Cat}_{\infty, \omega} = \sum_{X:\mathcal{U}} \text{is-Segal}(X) \times \text{is-complete}(X)$$

Some elementary results

- (∞, ω) -categories are stable by pullback and sum.

Some elementary results

- (∞, ω) -categories are stable by pullback and sum.
- hom-types of (∞, ω) -categories still are.

Some elementary results

- (∞, ω) -categories are stable by pullback and sum.
- hom-types of (∞, ω) -categories still are.
- discrete types are (∞, ω) -categories.

Some elementary results

- (∞, ω) -categories are stable by pullback and sum.
- hom-types of (∞, ω) -categories still are.
- discrete types are (∞, ω) -categories.
- representables types $\mathbb{Y}P$ are (∞, ω) -categories.

What's next ?

Main goal: Proving a Yoneda Lemma.

What's next ?

Main goal: Proving a Yoneda Lemma.

- Define a well-suited notion of fibration.

What's next ?

Main goal: Proving a Yoneda Lemma.

- Define a well-suited notion of fibration.
- Having a functorial $\text{hom}_A : A^{\text{op}} \times A \rightarrow \mathcal{U}$.

What's next ?

Main goal: Proving a Yoneda Lemma.

- Define a well-suited notion of fibration.
- Having a functorial $\text{hom}_A : A^{\text{op}} \times A \rightarrow \mathcal{U}$.
- Types of lax and oplax functors $A \multimap B$ and $A \rightharpoonup B$.

What's next ?

Main goal: Proving a Yoneda Lemma.

- Define a well-suited notion of fibration.
- Having a functorial $\text{hom}_A : A^{\text{op}} \times A \rightarrow \mathcal{U}$.
- Types of lax and oplax functors $A \multimap B$ and $A \rightharpoonup B$.
- Crans-Gray tensor product \boxtimes .

Towards a bunched-affine variant of HoTT

$$\Gamma, \Delta \quad ::= \quad \diamond \mid \Gamma * \Delta \mid \Gamma, x : A$$

Towards a bunched-affine variant of HoTT

$$\Gamma, \Delta \quad ::= \quad \diamond \mid \Gamma * \Delta \mid \Gamma, x : A$$

- Contexts are trees.

Towards a bunched-affine variant of HoTT

$$\Gamma, \Delta \quad ::= \quad \diamond \mid \Gamma * \Delta \mid \Gamma, x : A$$

- Contexts are trees.
- Weakening is admissible.

Towards a bunched-affine variant of HoTT

$$\Gamma, \Delta \quad ::= \quad \diamond \mid \Gamma * \Delta \mid \Gamma, x : A$$

- Contexts are trees.
- Weakening is admissible.
- A non-symmetric affine product \otimes .

Towards a bunched-affine variant of HoTT

$$\Gamma, \Delta \quad ::= \quad \diamond \mid \Gamma * \Delta \mid \Gamma, x : A$$

- Contexts are trees.
- Weakening is admissible.
- A non-symmetric affine product \otimes .
- New abstractions ∂ and ϱ for adjoints to \otimes .

Towards a bunched-affine variant of HoTT

$$\Gamma, \Delta \quad ::= \quad \diamond \mid \Gamma * \Delta \mid \Gamma, x : A$$

- Contexts are trees.
- Weakening is admissible.
- A non-symmetric affine product \otimes .
- New abstractions ∂ and ϱ for adjoints to \otimes .

$$\text{hom}_A(x, y) = \sum_{f: \mathbb{A}[1] \rightarrow A} (\langle 0 \mid f = x \rangle \times \langle 1 \mid f = y \rangle)$$

Some rules

$$(\otimes\text{-INTRO}) \frac{\underline{\Lambda} \mid \Gamma \vdash a : A \quad \underline{\Lambda} \mid \Delta \vdash b : B}{\underline{\Lambda} \mid \Gamma \otimes \Delta \vdash a \otimes b : A \otimes B}$$

Some rules

$$(\otimes\text{-INTRO}) \frac{\underline{\Lambda} \mid \Gamma \vdash a : A \quad \underline{\Lambda} \mid \Delta \vdash b : B}{\underline{\Lambda} \mid \Gamma \otimes \Delta \vdash a \otimes b : A \otimes B}$$

$$(\rightharpoonup\text{-INTRO}) \frac{\underline{\Lambda} \mid \diamond \vdash A : \mathcal{U} \quad \underline{\Lambda} \mid (x : A) \otimes \Gamma \vdash b : B}{\underline{\Lambda} \mid \Gamma \vdash \partial x.b : A \rightharpoonup B}$$

Some rules

$$(\otimes\text{-INTRO}) \frac{\underline{\Lambda} \mid \Gamma \vdash a : A \quad \underline{\Lambda} \mid \Delta \vdash b : B}{\underline{\Lambda} \mid \Gamma \otimes \Delta \vdash a \otimes b : A \otimes B}$$

$$(\rightarrow\text{-INTRO}) \frac{\underline{\Lambda} \mid \diamond \vdash B : \mathcal{U} \quad \underline{\Lambda} \mid \diamond \vdash A : \mathcal{U} \quad \underline{\Lambda} \mid (x : A) \otimes \Gamma \vdash b : B}{\underline{\Lambda} \mid \Gamma \vdash \partial x. b : A \rightarrow B}$$

$$(\rightarrow\text{-ELIM}) \frac{\underline{\Lambda} \mid \Gamma \vdash a : A \quad \underline{\Lambda} \mid \Delta \vdash f : A \rightarrow B}{\underline{\Lambda} \mid \Gamma \otimes \Delta \vdash \langle a \mid f \rangle : B}$$

Some rules (2)

$$(\rightarrow\text{-INTRO}) \frac{\underline{\Lambda} \mid \diamond \vdash B : \mathcal{U} \quad \underline{\Lambda} \mid \Gamma \otimes (x : A) \vdash b : B}{\underline{\Lambda} \mid \Gamma \vdash \text{ex}.b : A \rightarrow B}$$

Some rules (2)

$$(\rightarrow\text{-INTRO}) \frac{\underline{\Lambda} \mid \diamond \vdash B : \mathcal{U} \quad \underline{\Lambda} \mid \Gamma \otimes (x : A) \vdash b : B}{\underline{\Lambda} \mid \Gamma \vdash \text{ex}.b : A \rightarrow B}$$

$$(\rightarrow\text{-ELIM}) \frac{\underline{\Lambda} \mid \Gamma \vdash f : A \rightarrow B \quad \underline{\Lambda} \mid \Delta \vdash a : A}{\underline{\Lambda} \mid \Gamma \otimes \Delta \vdash f \mid a \rangle : B}$$

The end

Thank you !

Questions ?

