

# **Natural variation in teosinte at the domestication locus *teosinte branched1* (*tb1*)<sup>1</sup>**

Laura Vann<sup>2</sup>, Thomas Kono<sup>2,3</sup>, Tanja Pyhäjärvi<sup>2,4</sup>, Matthew B. Hufford<sup>2,5,7</sup>, and  
Jeffrey Ross-Ibarra<sup>2,6,7</sup>

<sup>2</sup>Department of Plant Sciences, University of California Davis

<sup>3</sup>Department of Agronomy and Plant Genetics, University of Minnesota Twin Cities

<sup>4</sup>Department of Biology, University of Oulu

<sup>5</sup>Department of Ecology, Evolution, and Organismal Biology, Iowa State University

<sup>6</sup>Center for Population Biology and Genome Center, University of California Davis

<sup>7</sup>Author for Correspondence

Manuscript received \_\_\_\_\_; revision accepted\_\_\_\_\_.

## **Acknowledgements**

Place brief acknowledgments, if desired, as a separate paragraph, using the following style: The author(s) thank(s). For brevity, do not use first names. Include grant acknowledgments here.

## 1 Abstract

2 The *teosinte branched1* (*tb1*) gene, a repressor of lateral organ growth, is a major  
3 QTL involved in branching differences between maize and its wild progenitor,  
4 teosinte. Previous work has shown that the insertion of a transposable element  
5 (*Hopscotch*) upstream of *tb1* enhances the gene's expression, causing much of the  
6 reduction in tillering observed in domesticated maize. Observations of the maize  
7 *tb1* allele in teosinte individuals, coupled with estimates of an age of insertion of  
8 the *Hopscotch* element that predates domestication, led us to investigate its  
9 prevalence and potential role in teosinte. Results from genotyping across many  
10 natural populations suggest that the *Hopscotch* element is segregating at a higher  
11 than expected frequency in a number of populations of two subspecies of teosinte,  
12 *Zea mays* ssp. *parviglumis* and *Zea mays* ssp. *mexicana*. Analysis of linkage  
13 disequilibrium between the *Hopscotch* element and variation in surrounding regions  
14 does not support a hypothesis of recent introgression from maize into teosinte.  
15 Population genetic signatures, however, are consistent with selection on this locus  
16 and suggest the *Hopscotch* insertion at *tb1* may play an ecological role in teosinte.  
17 Finally, two greenhouse experiments with *Zea mays* ssp. *parviglumis* do not suggest  
18 *tb1* controls tillering in natural populations of this subspecies. Our findings suggest  
19 that the role of the *Hopscotch* in tillering in teosinte differs from domesticated  
20 maize, and that other loci may play a role in observed variation in this trait.

# INTRODUCTION

Domesticated crops and their wild progenitors provide an excellent system in which to study adaptation and genomic changes associated with human-mediated selection (Ross-Ibarra et al., 2007). Perhaps the central focus of the study of domestication has been the identification of genetic variation underlying agronomically important traits such as fruit size and plant architecture (Olsen and Gross, 2010). Additionally, many domesticates show reduced genetic diversity when compared to their wild progenitors, and an understanding of the distribution of diversity in the wild and its phenotypic effects has become increasingly useful to crop improvement (Kovach and McCouch, 2008). But while some effort has been invested into understanding how wild alleles behave in their domesticated relatives (e.g. Bai and Lindhout, 2007), very little is known about the role that alleles found most commonly in domesticates play in natural populations of their wild progenitors (Whitton J, 1997).

Maize (*Zea mays* ssp. *mays*) was domesticated from the teosinte *Zea mays* ssp. *parviglumis* (hereafter, *parviglumis*) roughly 9,000 B.P. in southwest Mexico (Piperno et al., 2009; Matsuoka et al., 2002). Domesticated maize and the teosintes are an attractive system in which to study domestication due to the abundance of genetic tools developed for maize and well-characterized domestication loci (Hufford et al., 2012a; Doebley, 2004; Hufford et al., 2012b). Additionally, large naturally occurring populations of both *Zea mays* ssp. *parviglumis* (the wild progenitor of maize) and *Zea mays* ssp. *mexicana* (highland teosinte; hereafter *mexicana*) can be found throughout Mexico (Wilkes, 1977; Hufford et al., 2013), and genetic diversity of these taxa is estimated to be high (Ross-Ibarra et al., 2009).

Many morphological changes are associated with maize domestication, and understanding the genetic basis of these changes has been a focus of maize research

1 for a number of years (Doebley, 2004). One of the most dramatic changes is found  
2 in plant architecture: domesticated maize is characterized by a central stalk with  
3 few tillers and lateral branches terminating in a female inflorescence, while teosinte  
4 is highly tillered and bears tassels (male inflorescences) at the end of its lateral  
5 branches. The *teosinte branched1* (*tb1*) gene, a repressor of organ growth, was  
6 identified as a major QTL involved in branching (Doebley et al., 1995) and tillering  
7 (Doebley and Stec, 1991) differences between maize and teosinte. A 4.9 kb  
8 retrotransposon (*Hopscotch*) insertion into the upstream control region of *tb1* in  
9 maize acts to enhance expression of *tb1*, thus repressing lateral organ growth  
10 (Doebley et al., 1997; Studer et al., 2011). Dating of the *Hopscotch* retrotransposon  
11 suggests that its insertion predates the domestication of maize, leading to the  
12 hypothesis that it was segregating as standing variation in ancient populations of  
13 teosinte and increased to high frequency in maize due to selection during  
14 domestication (Studer et al., 2011). The effects of the *Hopscotch* insertion have  
15 been studied in maize (Studer et al., 2011), and analysis of teosinte alleles at *tb1*  
16 has identified functionally distinct allelic classes (Studer and Doebley, 2012), but  
17 little is known about the role of *tb1* or the *Hopscotch* insertion in natural  
18 populations of teosinte.

19 In teosinte and other plants that grow at high population density, individuals  
20 detect competition from neighbors via the ratio of red to far-red light. An increase  
21 in far-red relative to red light accompanies shading and triggers the shade  
22 avoidance syndrome: a suite of physiological and morphological changes such as  
23 reduced tillering, increased plant height and early flowering (Kebrom and Brutnell,  
24 2007). The *tb1* locus appears to play an important role in the shade avoidance  
25 pathway in *Zea mays* and other grasses and may therefore be crucial to the ecology  
26 of teosinte (Kebrom and Brutnell, 2007; Lukens and Doebley, 1999). In this study

1 we aim to characterize the distribution of the *Hopscotch* insertion in *parviglumis*,  
2 *mexicana*, and landrace maize, and to examine the phenotypic effects of the  
3 insertion in *parviglumis*. We use a combination of PCR genotyping for the  
4 *Hopscotch* element in our full panel and sequencing of two small regions upstream  
5 of *tb1* in a subset of teosinte populations to explore patterns of genetic variation at  
6 this locus. Finally, we test for an association between the *Hopscotch* element and  
7 tillering phenotypes in a population of *parviglumis*.

## 8 MATERIALS AND METHODS

### 9 Sampling and genotyping

10 We sampled 1,110 individuals from 350 accessions (247 maize landraces, 17  
11 *mexicana* populations, and 86 *parviglumis* populations) and assessed the presence  
12 or absence of the *Hopscotch* insertion (Table S1 and Table S2). DNA was extracted  
13 from leaf tissue using a modified CTAB approach (Doyle and Doyle, 1990; Maloof  
14 et al., 1984). We designed primers using PRIMER3 (Rozen and Skaletsky, 2000)  
15 implemented in Geneious (Kearse et al., 2012) to amplify the entire *Hopscotch*  
16 element, as well as an internal primer allowing us to simultaneously check for  
17 possible PCR bias between presence and absence of the *Hopscotch* insertion. Two  
18 PCRs were performed for each individual, one with primers flanking the *Hopscotch*  
19 (HopF/HopR) and one with a flanking primer and an internal primer  
20 (HopF/HopIntR). Primer sequences are HopF,  
21 5'-TCGTTGATGCTTTGATGGATGG-3'; HopR, 5'-AACAGTATGATTTTCATGGGACCG-3';  
22 and HopIntR, 5'-CCTCCACCTCTCATGAGATCC-3' (Fig. S1, Fig. S2) *Primers in Fig. S1*  
23 *should be labeled* . Homozygotes show a single band for absence of the element  
24 (~300bp) and two bands for presence of the element (~5kb and XX *LV, please add the*

1 *size of the second band* ), whereas heterozygotes are three-banded (Fig. S2). When only  
2 one PCR resolved well, we scored one allele for the individual. We used Phusion  
3 High Fidelity Enzyme (Thermo Fisher Scientific Inc., Waltham, Massachusetts,  
4 USA) and the following conditions for amplifications: 98°C for 3 min, 30 cycles of  
5 98°C for 15 s, 65°C for 30 s, and 72°C for 3 min 30 s, with a final extension of 72°C  
6 for 10 min. PCR products were visualized on a 1% agarose gel and scored for  
7 presence/absence of the *Hopscotch* based on band size.

## 8 Sequencing

9 In addition to genotyping, we chose a subset of *parviglumis* individuals for  
10 sequencing. We chose twelve individuals from each of four populations from Jalisco  
11 state, Mexico (San Lorenzo, La Mesa, Ejutla A, and Ejutla B). For amplification  
12 and sequencing, we selected two regions approximately 600bp in size from within  
13 the 5' UTR of *tb1* (Region 1) and from 1,235bp upstream of the start of the  
14 *Hopscotch* (66,169bp upstream from the start of the *tb1* ORF; Region 2). We  
15 designed the following primers using PRIMER3 (Rozen and Skaletsky, 2000): for  
16 the 5' UTR, 5'-GGATAATGTGCACCAGGTGT-3' and  
17 5'-GCGTGCTAGAGACACYTGTGCT-3'; for the 66kb upstream region,  
18 5'-TGTCCTCGCCGCAACTC-3' and 5'-TGTACGCCCGCCCTCATCA-3' (Fig. S1). We  
19 used Taq polymerase (New England Biolabs Inc., Ipswich, Massachusetts, USA)  
20 and the following thermal cycler conditions to amplify fragments: 94°C for 3 min,  
21 30 cycles of 92°C for 40 s, annealing for 1 min, 72°C for 40 s, and a final 10 min  
22 extension at 72°C. Annealing temperatures for Region 1 and Region 2 were 59.7°C  
23 and 58.8°C, respectively. To clean excess primer and dNTPs we added two units of  
24 Exonuclease1 and 2.5 units of Antarctic Phosphatase to 8.0  $\mu$ L of amplification  
25 product. This mix was placed on a thermal cycler with the following program:

1 37°C for 30 min, 80°C for 15 min, and a final cool-down step to 4°C.  
2 We cloned cleaned fragments into a TOPO-TA vector (Life Technologies,  
3 Grand Island, New York, USA) using OneShot TOP10 chemically competent *E.*  
4 *coli* cells, with an extended ligation time of 30 min for a complex target fragment.  
5 We plated cells on LB agar plates containing kanamycin, and screened colonies  
6 using vector primers M13 Forward and M13 Reverse under the following conditions:  
7 96°C for 5 min; then 35 cycles at 96°C for 30 s, 53°C for 30 s, 72°C for 2 min; and  
8 a final extension at 72°C for 4 min. We visualized amplification products for  
9 incorporation of our insert on a 1% agarose TAE gel.

10 Amplification products with successful incorporation of our insert were cleaned  
11 using Exonuclease 1 and Antarctic Phosphatase following the procedures detailed  
12 above, and sequenced with vector primers M13 Forward and M13 Reverse using  
13 Sanger sequencing at the College of Agriculture and Environmental Sciences  
14 (CAES) sequencing center at UC Davis. We aligned and trimmed primer sequences  
15 from resulting sequences using the software Geneious (Kearse et al., 2012).  
16 Following alignment, we verified singleton SNPs by sequencing an additional one to  
17 four colonies from each clone. If the singleton was not present in these additional  
18 sequences it was considered an amplification or cloning error, and we replaced the  
19 base with the base of the additional sequences. If the singleton appeared in at least  
20 one of the additional sequences we considered it a real variant and kept it for  
21 further analyses.

## 22 Genotyping analysis

23 We examined discrepancies between observed and expected genotype frequencies by  
24 calculating Hardy-Weinberg Equilibrium (HWE). To calculate differentiation  
25 between populations ( $F_{ST}$ ) and subspecies ( $F_{CT}$ ) we used HierFstat (Goudet,



2005). These analyses only included populations in which 8 or more individuals were sampled. To test the hypothesis that the *Hopscotch* insertion may be adaptive under certain environmental conditions, we looked for significant associations between the *Hopscotch* frequency and environmental variables using BayEnv (Coop et al., 2010). BayEnv creates a covariance matrix of relatedness between populations and then tests a null model that allele frequencies in populations are determined by the covariance matrix of relatedness alone against the alternative model that allele frequencies are determined by a combination of the covariance matrix and an environmental variable, producing a posterior probability (*i.e.*, Bayes Factor; Coop et al. 2010). We used genotyping and covariance data from Pyhäjärvi et al. (2013) for BayEnv, with the *Hopscotch* insertion coded as an additional SNP (Table S3). Environmental data were obtained from [www.worldclim.org](http://www.worldclim.org), the Harmonized World Soil Database (FAO/IIASA/ISRIC/ISSCAS/JRC, 2012) and [www.harvestchoice.org](http://www.harvestchoice.org) and summarized by principle component analysis following Pyhäjärvi et al. (2013).

## Sequence analysis

For population genetic analyses of sequenced Region 1 and sequenced Region 2 we used the Libsequence package (Thornton, 2003) to calculate pairwise  $F_{ST}$  between populations and to calculate standard diversity statistics (number of haplotypes, haplotype diversity, Watterson’s estimator  $\hat{\theta}_W$ , pairwise nucleotide diversity  $\hat{\theta}_\pi$ , and Tajima’s D). To produce a visual representation of differentiation between sequences and examine patterns in sequence clustering by *Hopscotch* genotype we used Phylip (<http://evolution.genetics.washington.edu/phylip.html>), creating neighbor-joining trees with bootstrap-supported nodes (100 repetitions). For creation of trees we also included homologous sequence data from Maize

1 HapMapV2 (Chia et al., 2012) for teosinte inbred lines (TILs), some of which are  
2 known to be homozygous for the *Hopscotch* insertion (TIL03, TIL17, TIL09), as  
3 well as 59 lines of domesticated maize.

#### 4 **Introgression analysis**

5 In order to assess patterns of linkage disequilibrium (LD) around the *Hopscotch*  
6 element in the context of chromosomal patterns of LD we used Tassel (Bradbury  
7 et al., 2007) and calculated LD between SNPs across chromosome 1 using  
8 previously published data from twelve plants each of the Ejutla A (EjuA), Ejutla B  
9 (EjuB), San Lorenzo (SLO), and La Mesa (MSA) populations (Pyhäjärvi et al.,  
10 2013). We chose these populations because we had both genotyping data for the  
11 *Hopscotch* as well as chromosome-wide SNP data for chromosome 1. For each  
12 population we filtered the initial set of 5,897 SNPs on chromosome 1 to accept only  
13 SNPs with a minor allele frequency of at least 0.1, resulting in 1,671, 3,023, 3,122,  
14 and 2,167 SNPs for SLO, EjuB, EjuA, and MSA, respectively. We then used Tassel  
15 (Bradbury et al., 2007) to calculate linkage disequilibrium ( $r^2$ ) across chromosome  
16 1 for each population.

17 We examined evidence of introgression on chromosome 1 in these same four  
18 populations (EjuA, EjuB, MSA, SLO) using STRUCTURE (Falush et al., 2003)  
19 and phased data from Pyhäjärvi et al. (2013), combined with the corresponding  
20 SNP data from a diverse panel of 282 maize lines (Cook et al., 2012). SNPs were  
21 anchored in a modified version of the IBM genetic map (Gerke et al., 2013). We  
22 created haplotype blocks using a custom Perl script that grouped SNPs separated  
23 by less than 5kb into haplotypes. We ran STRUCTURE at K=2 under the linkage  
24 model, performing 3 replicates with an MCMC burn-in of 10,000 steps and 50,000  
25 steps post burn-in.

## 1 Phenotyping of *parviglumis*

2 To investigate the phenotypic effects of the *Hopscotch* insertion in teosinte, we  
3 conducted an initial phenotyping trial (Phenotyping 1). We germinated 250 seeds  
4 of *parviglumis* collected in Jalisco state, Mexico (population San Lorenzo) (Hufford,  
5 2010) where the *Hopscotch* is segregating at highest frequency (0.44) in our initial  
6 genotyping sample set. In order to maximize the likelihood of finding the *Hopscotch*  
7 in our association population we selected seeds from sites where genotyped  
8 individuals were homozygous or heterozygous for the insertion. We chose between  
9 10-13 seeds from each of 23 sampling sites. We treated seeds with fungicide and  
10 germinated them in petri dishes with filter paper. Following germination, 206  
11 successful germinations were planted into one-gallon pots with potting soil and  
12 randomly spaced one foot apart on greenhouse benches. Plants were watered three  
13 times a day by hand and with an automatic drip containing 10-20-10 fertilizer.

14 Starting on day 15, we measured tillering index as the ratio of the sum of tiller  
15 lengths to the height of the plant (Briggs et al., 2007). Following initial  
16 measurements, we phenotyped plants for tillering index every 5 days through day  
17 40, and then on day 50 and day 60. On day 65 we measured culm diameter  
18 between the third and fourth nodes of each plant. Culm diameter is not believed to  
19 be correlated with tillering index or variation at *tb1*. Following phenotyping we  
20 extracted DNA from all plants using a modified SDS extraction protocol  
21 (<http://www.ars.usda.gov>). *what is this url?* We genotyped individuals for the  
22 *Hopscotch* insertion following the protocols listed above. Based on these initial  
23 data, we conducted a *post hoc* power analysis using data from day 40 of  
24 Phenotyping 1, indicating that a minimum of 71 individuals in each genotypic class  
25 would be needed to detect the observed effect of the *Hopscotch* on tillering index.

26 We performed a second phenotyping experiment (Phenotyping 2) in which we

1 germinated 372 seeds of *parviglumis*, choosing equally between sites previously  
2 determined to have or not have the *Hopscotch* insertion. Seeds were germinated  
3 and planted on day 7 post fruit-case removal into two gallon pots. Plants were  
4 watered twice daily, alternating between fertilized and non-fertilized water. We  
5 began phenotyping successful germinations (302 plants) for tillering index on day  
6 15 post fruit-case removal, and phenotyped every five days until day 50. At day 50  
7 we measured culm diameter between the third and fourth nodes. We extracted  
8 DNA and genotyped plants following the same guidelines as in Phenotyping 1.


9 Tillering index data for each genotypic class did not meet the criteria for a  
10 repeated measures ANOVA, so we transformed the data using a Box-Cox  
11 transformation ( $\alpha = 0$  *what is the alpha value here?* ; Car Package for R, Fox and Weisberg  
12 2011) to improve the normality and homogeneity of variance among genotype  
13 classes. We analyzed relationships between genotype and tillering index and tiller  
14 number using a repeated measures ANOVA through a general linear model function  
15 implemented in SAS v.9.3 (SAS Institute Inc., Cary, NC, USA). Additionally, in  
16 order to compare any association between *Hopscotch* genotype and tillering and  
17 associations at other presumably unrelated traits, we performed an ANOVA  
18 between culm diameter and genotype using the same general linear model in SAS.

## 19 RESULTS

### 20 Genotyping

21 Genotype of the *Hopscotch* insertion was confirmed with two PCRs for 837  
22 individuals. Among the 247 maize landrace accessions genotyped, all but eight were  
23 homozygous for the presence of the insertion (Table S1 and Table S2). Within our  
24 *parviglumis* and *mexicana* samples we found the *Hopscotch* insertion segregating in

Figure 1: Map showing the frequency of the *Hopscotch* allele in populations of *parviglumis* where we sampled more than 6 individuals. Size of circles reflects number of alleles sampled.



- 1 37 and 4 populations, respectively, and at highest frequency in the states of Jalisco,
- 2 Colima, and Michoacán in central-western Mexico (Fig. 1).
- 3 Using our *Hopscotch* genotyping, we calculated differentiation between
- 4 populations ( $F_{ST}$ ) and subspecies ( $F_{CT}$ ) for populations in which we sampled 8 or
- 5 more alleles. We found that  $F_{CT} = 0$ , and levels of  $F_{ST}$  among populations within
- 6 each subspecies (0.22) and among all populations (0.23) *are these an average of pairwise or is*
- 7 *this calculated among all pops?* are similar to those reported genome-wide in previous
- 8 studies (Pyhäjärvi et al. 2013; Table 1). Although we found large variation in
- 9 *Hopscotch* allele frequency among our populations, BayEnv analysis did not
- 10 indicate a correlation between the *Hopscotch* insertion and environmental variables
- 11 (all Bayes Factors  $< 1$ ; Table S3).

Table 1: Pairwise  $F_{ST}$  values from sequence and *Hopscotch* genotyping data

| Comparison  | Region 1 | Region 2 | <i>Hopscotch</i> |
|-------------|----------|----------|------------------|
| EjuA & EjuB | 0        | 0        | 0                |
| EjuA & MSA  | 0.326    | 0.328    | 0.186            |
| EjuA & SLO  | 0.416    | 0.258    | 0.280            |
| EjuB & MSA  | 0.397    | 0.365    | 0.188            |
| EjuB & SLO  | 0.512    | 0.290    | 0.280            |
| MSA & SLO   | 0.007    | 0        | 0.016            |

## 1 Sequencing

2 To investigate patterns of sequence diversity and linkage disequilibrium (LD) in the  
3 *tb1* region, we sequenced two small (<1kb) regions upstream of the *tb1* ORF in  
4 four populations. After alignment and singleton checking we recovered 48 and 40  
5 segregating sites for the 5' UTR region (Region 1) and the 66kb upstream region  
6 (Region 2), respectively. For Region 1, Ejutla A has the highest values of haplotype  
7 diversity, and  $\theta_\pi$ , while Ejutla B and La Mesa have comparable values of these  
8 summary statistics, and San Lorenzo has much lower values. Additionally, Tajima's  
9 D is strongly negative in the two Ejutla populations and La Mesa, but is less  
10 negative in San Lorenzo (Table 2). *need to reference Hopscotch frequencies in supplemental table*  
11 *somewhere* For Region 2, haplotype diversity and  $\theta_\pi$ , are similar for Ejutla A and  
12 Ejutla B, while La Mesa and San Lorenzo have slightly lower values for these  
13 statistics (Table 2). Tajima's D is positive in all populations except San Lorenzo, *is*  
14 *the table wrong? MSA is the only negative value in the table* indicating an excess of low frequency  
15 variants in this population (Table 2). Pairwise values of  $F_{ST}$  within population  
16 pairs Ejutla A/Ejutla B and San Lorenzo/La Mesa are 0 for both sequenced

regions as well as for the *Hopscotch* [table 1 shows 0.016 for hopscotch, not 0. which is right?](#) , while they are high for other population pairs (Table 1). Neighbor joining trees of our sequence data and data from the teosinte inbred lines (TILs; data from Maize HapMapV2, Chia et al. 2012) do not reveal any clear clustering pattern with respect to population or *Hopscotch* genotype (Figure S3); individuals within our sample that have the *Hopscotch* insertion do not group with the teosinte inbred lines or domesticated maize that have the *Hopscotch* insertion.

Table 2: Population genetic statistics from resequenced regions near the *tb1* locus

| Population                      | # Haplotypes | Hap. Diversity | $\hat{\theta}_{\pi}$ | Tajima's D |
|---------------------------------|--------------|----------------|----------------------|------------|
| <i>Region 1 (5' UTR)</i>        |              |                |                      |            |
| EJUA                            | 8            | 0.859          | 0.005                | -1.650     |
| EJUB                            | 5            | 0.709          | 0.004                | -1.831     |
| MSA                             | 6            | 0.682          | 0.004                | -1.755     |
| SLO                             | 3            | 0.318          | 0.001                | -0.729     |
| <i>Region 2 (66kb upstream)</i> |              |                |                      |            |
| EJUA                            | 8            | 0.894          | 0.018                | 0.623      |
| EJUB                            | 8            | 0.894          | 0.016                | 0.295      |
| MSA                             | 3            | 0.682          | 0.011                | -0.222     |
| SLO                             | 4            | 0.742          | 0.014                | 0.932      |

## 8 Evidence of introgression

The highest frequency of the *Hopscotch* insertion in teosinte was found in *parviglumis* sympatric with cultivated maize. Our initial hypothesis was that the high frequency of the *Hopscotch* element in these populations could be attributed

1 to introgression from maize into teosinte. To investigate this possibility we  
2 examined overall patterns of linkage disequilibrium across chromosome one and  
3 specifically in the *tb1* region. If the *Hopscotch* is found in these populations due to  
4 recent introgression we would expect to find large blocks of linked markers near this  
5 element. We find no evidence of elevated linkage disequilibrium between the  
6 *Hopscotch* and SNPs surrounding the *tb1* region in our resequenced populations  
7 (Fig. 2), and  $r^2$  in the *tb1* region does not differ significantly between populations  
8 with (average  $r^2$  of 0.085) and without (average  $r^2 = 0.082$ ) the *Hopscotch*  
9 genotype. In fact, average  $r^2$  is lower in the *tb1* region ( $r^2 = 0.056$ ) than across the  
10 rest of chromosome 1 ( $r^2 = 0.083$ ) (3). *LV, please go through and make sure the data entered into*  
11 *all the tables is correct. In Table3, both sequenced regions were labeled as "Region 1". I changed the second to*  
12 *Region 2 but don't know if the data in this column are really from Region 2*

Table 3:  $r^2$  values between SNPs on chromosome 1, in the broad *tb1* region, within the 5' UTR of *tb1* (Region 1), and 66kb upstream of *tb1* (Region 2).

| Population  | Chr. 1 | <i>tb1</i> region | Region 1 | Region 2 |
|-------------|--------|-------------------|----------|----------|
| Ejutla A    | 0.095  | 0.050             | 0.747    | 0.215    |
| Ejutla B    | 0.069  | 0.051             | 0.660    | 0.186    |
| La Mesa     | 0.070  | 0.053             | 0.914    | 0.766    |
| San Lorenzo | 0.101  | 0.067             | 0.912    | 0.636    |

13 The lack of clustering of *Hopscotch* genotypes in our NJ tree as well as the lack  
14 of LD around *tb1* does not support the hypothesis that the *Hopscotch* insertion in  
15 these populations of *parviglumis* is the result of recent introgression. However, to  
16 further explore this hypothesis we performed a STRUCTURE analysis using  
17 Illumina MaizeSNP50 data from four of our *parviglumis* populations (EjuA, EjuB,






Figure 2: Linkage disequilibrium for SNPs in Mb 261-268 on chromosome 1. The yellow rectangle indicates the location of the *Hopscotch* insertion and the green represents the *tb1* ORF. A) Ejutla A; B) Ejutla B; C) La Mesa; D). San Lorenzo

*this needs description of what upper and lower triangle are*

1 MSA, and SLO) and the maize 282 diversity panel (Cook et al., 2012; Pyhäjärvi  
2 et al., 2013). The linkage model implemented in STRUCTURE can be used to  
3 identify ancestry of blocks of linked variants, which would arise as a result of recent  
4 admixture between populations. If the *Hopscotch* insertion is present in  
5 populations of *parviglumis* as a result of recent admixture with domesticated maize,  
6 we would expect the insertion and linked variants in surrounding sites to be

1 assigned to the "maize" cluster in our STRUCTURE runs, not the "teosinte"  
2 cluster. In all runs, assignment to maize in the *tb1* region across all four  
3 *parviglumis* populations is low (average 0.017) and much below the  
4 chromosome-wide average (0.20; Table 4; Fig. 3).

Table 4: Assignments to maize and teosinte in the *tb1* and chromosome 1 regions from STRUCTURE

|             | <i>tb1</i> region |          | Chr 1 |          |
|-------------|-------------------|----------|-------|----------|
| Population  | Maize             | Teosinte | Maize | Teosinte |
| Ejutla A    | 0.022             | 0.978    | 0.203 | 0.797    |
| Ejutla B    | 0.019             | 0.981    | 0.187 | 0.813    |
| La Mesa     | 0.012             | 0.988    | 0.193 | 0.807    |
| San Lorenzo | 0.016             | 0.984    | 0.205 | 0.795    |

## 5 Phenotyping

6 To assess the contribution of *tb1* to phenotypic variation in tillering in a natural  
7 population, we grew plants from seed sampled from the San Lorenzo population of  
8 *parviglumis*, which had a high mean frequency (0.44) of the *Hopscotch* insertion  
9 based on our initial genotyping. We measured tillering index (TI), the ratio of the  
10 sum of tiller lengths to plant height, for 216 plants (Phenotyping 1) from within  
11 the San Lorenzo population, and genotyped plants for the *Hopscotch* insertion. We  
12 found the *Hopscotch* segregating at a frequency of 0.65 with no significant  
13 deviations from expected frequencies under Hardy-Weinberg equilibrium. After  
14 performing a repeated measures ANOVA between our transformed tillering index  
15 data and *Hopscotch* genotype we find a weak positive correlation between presence  
16 of the *Hopscotch* and tillering index on day 40 (p=0.0848), a result indicating the




Figure 3: STRUCTURE assignment to maize across a section of chromosome 1. The dotted lines mark the beginning of the sequenced region 50kb upstream (Sequenced region 2) and the end of the *tb1* ORF.

- 1 *Hopscotch* may actually increase tillering in *parviglumis* in contrast to its
- 2 phenotypic effect in maize. We find no correlation between tillering index and
- 3 genotype on any other day (4). Additionally we find no significant correlation
- 4 between tiller number and *Hopscotch* genotype, or culm diameter and *Hopscotch*
- 5 genotype in Phenotyping 1.
- 6 We performed a second grow-out of *parviglumis* from San Lorenzo
- 7 (Phenotyping 2) to assess whether lighting conditions or sample size may have
- 8 affected our ability to detect an effect of *tb1*. For the second grow-out we measured
- 9 tillering index every five days through day 50 for 302 plants. We found the
- 10 *Hopscotch* allele segregating at a frequency of 0.69, *is it in HWE in this pop?* with a 0.6
- 11 frequency of *Hopscotch* homozygotes, and a 0.2 frequency of both heterozygotes
- 12 and homozygotes for the teosinte allele. We found similar patterns, with a weak




Figure 4: Box-plots showing tillering index in greenhouse grow-outs for Phenotyping 1 (A) and Phenotyping 2 (B). White indicates individuals homozygous for the *Hopscotch*, light grey represents heterozygotes, and dark grey represents homozygotes for the *teosinte* (No *Hopscotch*) allele. Within boxes, dark black lines represent the median, and the edges of the boxes are the first and third quartiles. *please explain whiskers and dots on figure too.*

1 positive correlation between tillering index and *Hopscotch* genotype at day 40  
2 ( $p < 0.0611$ ), with no significant correlation on any day. Similarly, relationships  
3 between *Hopscotch* genotype and tiller number and *Hopscotch* genotype and culm  
4 diameter were not significant.

## 5 DISCUSSION

6 Adaptation occurs due to selection on standing variation or *de novo* mutations.  
7 Adaptation from standing variation has been well-described in a number of systems;  
8 for example, selection for lactose tolerance in humans (Plantinga et al., 2012;  
9 Tishkoff et al., 2007), variation at the *Eda* locus in three-spined stickleback (Kitano  
10 et al., 2008; Colosimo et al., 2005), and pupal diapause in the Apple Maggot fly  
11 (Feder et al., 2003). Although the adaptive role of standing variation has been  
12 described in many systems, its importance in domestication is not as well studied.

13 In maize, alleles at domestication loci (*RAMOSA1*, Sigmon and Vollbrecht  
14 2010; *barren stalk1*, Gallavotti et al. 2004; and *grassy tillers1*, Whipple et al. 2011)  
15 are thought to have been selected from standing variation, suggesting that diversity  
16 already present in teosinte may have played an important role in maize  
17 domestication. The *teosinte branched1* gene is one of the best characterized  
18 domestication loci, and, while previous studies have suggested that differences in  
19 plant architecture between maize and teosinte are a result of selection on standing  
20 variation at this locus, little is known about natural variation at this locus and its  
21 ecological role in teosinte (Clark et al., 2006; Studer et al., 2011). Studer et al.  
22 (2011) genotyped 90 accessions of teosinte (inbred and outbred), providing the first  
23 evidence that the *Hopscotch* insertion is segregating in teosinte (Studer et al., 2011).

24 Given that the *Hopscotch* insertion has been estimated to predate the  
25 domestication of maize, it is not surprising that it can be found segregating in

1 populations of teosinte. However, by widely sampling across teosinte populations  
2 our study provides greater insight into the distribution and prevalence of the  
3 *Hopscotch* in teosinte. While our findings are consistent with Studer et al. (2011) in  
4 that we identify the *Hopscotch* allele segregating in teosinte, we find it at higher  
5 frequency than previously suggested (Studer et al., 2011). Many of our populations  
6 with high frequency of the *Hopscotch* allele fall in the Jalisco cluster identified by  
7 Fukunaga (2005), suggesting a different history of the *tb1* locus in this region than  
8 in the Balsas River Basin where maize was domesticated (Matsuoka et al., 2002).  
9 Potential explanations for the high frequency of the *Hopscotch* element in  
10 *parviglumis* from the Jalisco cluster include gene flow from maize, genetic drift, and  
11 natural selection.

12 While gene flow from crops into their wild relatives is well-known, (Ellstrand  
13 et al., 1999; Zhang et al., 2009; Thurber et al., 2010; Baack et al., 2008; Hubner  
14 et al., 2012; Wilkes, 1977; van Heerwaarden et al., 2011; Barrett, 1983), our results  
15 are more consistent with Hufford et al. (2013) who found resistance to introgression  
16 from maize into teosinte. Furthermore, Hufford et al. (2013) showed that  
17 domestication loci, such as *tb1*, are particularly resistant to introgression in both  
18 directions of gene flow (i.e., maize to teosinte and teosinte to maize). We find no  
19 evidence of recent introgression in our analyses. Clustering patterns in our NJ trees  
20 do not reflect a pattern expected if maize alleles at the *tb1* locus had introgressed  
21 into populations of teosinte. Moreover, there is no signature of elevated LD in the  
22 *tb1* region relative to the rest of chromosome 1, and Bayesian assignment to a maize  
23 cluster in this region is lower than the chromosome-wide average (Fig. 3, Table 4).  
24 Together, these data point to an explanation other than recent introgression for the  
25 high observed frequency of *Hopscotch* in a subset of our *parviglumis* populations.

26 Although recent introgression seems unlikely, we cannot rule out ancient

1 introgression as an explanation for the presence of the *Hopscotch* in these  
2 populations. If the *Hopscotch* allele was introgressed in the distant past,  
3 recombination may have broken up LD, a process that would be consistent with  
4 our data. We find this scenario less plausible, however, as there is no reason why  
5 gene flow should have been high in the past but absent in present-day sympatric  
6 populations. In fact, early generation maize-teosinte hybrids are common in these  
7 populations today (MB Hufford, pers. observation), and genetic data support  
8 ongoing gene flow between domesticated maize and both *mexicana* and *parviglumis*  
9 in a number of sympatric populations (Hufford et al., 2013; Ellstrand et al., 2007;  
10 van Heerwaarden et al., 2011).

11       Remaining explanations for differential frequencies of the *Hopscotch* among  
12 teosinte populations include both genetic drift and natural selection. Drift may  
13 have played a role in the San Lorenzo *parviglumis* population. Previous studies  
14 using both SSRs and genome-wide SNP data have found evidence for a population  
15 bottleneck in the San Lorenzo population (Hufford, 2010; Pyhäjärvi et al., 2013),  
16 and the lower levels of sequence diversity in this population in the 5' UTR (Region  
17 1) coupled with more positive values of Tajima's D are consistent with these earlier  
18 findings suggesting a bottleneck. *deviations from HWE may be consistent too if we see excess of*  
19 *homozygotes. do we?* Such population bottlenecks can exaggerate the effects of genetic  
20 drift through which the *Hopscotch* allele may have risen to high frequency entirely  
21 by chance. This bottleneck, however, does not explain the high frequency of the  
22 *Hopscotch* in multiple populations in the Jalisco cluster. Moreover, available  
23 information on diversity and population structure among Jaliscoan populations  
24 (Hufford, 2010; Pyhäjärvi et al., 2013) is not suggestive of recent colonization or  
25 other demographic events that would predict a high frequency of the allele across  
26 populations. Finally, values of the Tajima's D statistic in the 5' UTR of *tb1* are

1 suggestive of natural selection acting upon the gene in natural population of  
2 *parviglumis*. Whereas the genome-wide average of Tajima's D in genic regions of  
3 *parviglumis* is 0.45 (Hufford et al., 2012b), the statistic is quite negative in the 5'  
4 UTR of *tb1* (Table 2). This result is consistent with repeated selective sweeps near  
5 *tb1* and a putative ecological role for the gene in *parviglumis*.

6 *do we know the Hop genotype for sequenced lines? can we separate the sequences into hop/no hop and look*  
7 *for differences? it wasn't until we did this that gt1 stuff really popped out. we should know for some of them, i*  
8 *will check I've added a few sentences on selection. Do we still want to compare sequences with and without*  
9 *Hopscotch? I agree its a good idea and could end up being really interesting. Perhaps something we could look at*  
10 *after submission and incorporate during revisions?*

11 Significant effects of the *Hopscotch* insertion on lateral branch length, number  
12 of cupules, and tillering index in domesticated maize have been well documented  
13 (Studer et al., 2011). Weber et al. (2007) have described significant phenotypic  
14 associations between markers in and around *tb1* and lateral branch length and  
15 female ear length in a sample from 74 natural populations of *parviglumis* (Weber  
16 et al., 2007); however, these data did not include markers from the *Hopscotch*  
17 region 66kb upstream of *tb1*. Our study is the first to explicitly examine the  
18 phenotypic effects of the *Hopscotch* insertion across a wide collection of individuals  
19 sampled from natural populations of teosinte. We have found no significant effect  
20 of the *Hopscotch* insertion on tillering index or tiller number, a result that is  
21 discordant with its clear phenotypic effects in maize. One interpretation of this  
22 result would be that the *Hopscotch* controls tillering in maize (Studer et al., 2011),  
23 but tillering in teosinte is affected by variation at other loci. Consistent with this  
24 interpretation, *tb1* is thought to be part of a complex pathway controlling  
25 branching, tillering and other phenotypic traits (Kebrom and Brutnell, 2007; Clark  
26 et al., 2006). A recent study by Studer and Doebley (2012) examined variation



1 across traits in a three-taxa allelic series at the *tb1* locus. Studer and Doebley  
2 (2012) introgressed nine unique teosinte *tb1* segments (one from *Zea diploperennis*,  
3 and four each from *mexicana* and *parviglumis*) into an inbred maize background  
4 and investigated phenotypic effects. Phenotypes were shown to cluster by taxon,  
5 indicating *tb1* may underlie morphological diversification of *Zea*. Additional  
6 analysis in Studer and Doebley (2012) suggested tillering index was controlled both  
7 by *tb1* and loci elsewhere in the genome. Clues to the identity of these loci may be  
8 found in QTL studies that have identified loci controlling branching architecture  
9 (*e.g.*, Doebley and Stec 1991, 1993). Many of these loci (*grassy tillers*, *gt1*;  
10 *tassel-replaces-upper-ears1*, *tru1*; *terminal ear1*, *ter1*) have been shown to interact  
11 with *tb1* (Whipple et al., 2011; Li, 2012), and both *tru1* and *ter1* affect the same  
12 phenotypic traits as *tb1* (Doebley et al., 1995). *tassel-replaces-upper-ears1* (*tru1*),  
13 for example, has been shown to act either epistatically or downstream of *tb1*,  
14 affecting both branching architecture (decreased apical dominance) and tassel  
15 phenotypes (shortened tassel and shank length and reduced tassel number; Li  
16 2012). Variation in these additional loci may have affected tillering in our  
17 collections and contributed to the lack of correlation we see between *Hopscotch*  
18 genotype and tillering.

19 In conclusion, our findings demonstrate that the *Hopscotch* allele is more  
20 widespread in populations of *parviglumis* and *mexicana* than previously thought.  
21 Analysis of linkage using SNPs from across chromosome 1 does not suggest that the  
22 *Hopscotch* allele is present in these populations due to recent introgression;  
23 however, it seems unlikely that the insertion would have drifted to high frequency  
24 in multiple populations. We do, however, find preliminary evidence of selection on  
25 the *tb1* locus in *parviglumis*; this coupled with our observation of high frequency of  
26 the *Hopscotch* insertion in a number of populations suggests that the locus plays an

1 ecological role in teosinte. In contrast to domesticated maize, the *Hopscotch*  
 2 insertion in *parviglumis* does not appear to reduce tillering. Other loci involved in  
 3 branching architecture may regulate tillering in teosinte. Future studies should  
 4 examine expression levels of *tb1* in teosinte with and without the *Hopscotch*  
 5 insertion and further characterize the effects of additional loci involved in  
 6 branching architecture (e.g. *gt1*, *tru1*, and *ter1*). These data, in conjunction with  
 7 more exhaustive phenotyping, should help reveal the ecological significance of the  
 8 domesticated *tb1* allele in natural populations of teosinte. *why not Phyb and phya? Are*  
 9 *they necessary to include? I'd had them in before in a paragraph but had been voted out I'd ditch gt1 tru1 ter1*  
 10 *and maybe just cite some people including phyb etc.*  
 11 *please check format of supp figs and tables; some are running off the page. you can use "longtable" to fix that*  
 12 *(ask Paul for example). check fig/table references, bibliography, etc. what does "rotation" mean in supp. table 3?*  
 13 *it isn't mentioned in methods. please check that all the tables and figs (including supplement) are referenced in the*  
 14 *text.*

## 1 References

- 2 BAACK, E., Y. SAPIR, M. CHAPMAN, J. BURKE, AND L. RIESEBERG. 2008.  
3 Selection on domestication traits and quantitative trait loci in crop-wild  
4 sunflower hybrids. *Mol Ecol* 17: 666–677.
- 5 BAI, Y. AND P. LINDHOUT. 2007. Domestication and breeding of tomatoes: What  
6 have we gained and what can we gain in the future? *Annals of Botany* 100:  
7 1085–1094.
- 8 BARRETT, S. 1983. Crop mimicry in weeds. *Econ Bot* 37: 255–282.
- 9 BRADBURY, P., Z. ZHANG, D. KROON, T. CASSTEVENS, Y. RAMDOSS, AND  
10 E. BUCKLER. 2007. Tassel: software for association mapping of complex traits in  
11 diverse samples. *Bioinformatics* 23: 2633–2635.
- 12 BRIGGS, W., M. McMULLEN, B. GAUT, AND J. DOEBLEY. 2007. Linkage  
13 mapping of domestication loci in a large maize-teosinte backcross resource.  
14 *Genetics* 177: 1915–1928.
- 15 CHIA, J., C. SONG, P. BRADBURY, D. COSTICH, N. DE, LEON, J. DOEBLEY,  
16 R. ELSHIRE, B. GAUT, L. GELLER, J. GLAUBITZ, M. GORE, K. GUILL,  
17 J. HOLLAND, M. HUFFORD, J. LAI, M. LI, X. LIU, Y. LU, R. McCOMBIE,  
18 R. NELSON, J. POLAND, B. PRASANNA, T. PYHÄJÄRVI, T. RONG,  
19 R. SEKHON, Q. SUN, M. TENAILLON, F. TIAN, J. WANG, X. XU,  
20 Z. ZHANG, S. KAEPPLE, J. ROSS-IBARRA, M. McMULLEN, E. BUCKLER,  
21 G. ZHANG, Y. XU, AND D. WARE. 2012. Maize hapmap2 identifies extant  
22 variation from a genome in flux. *Nat Genet* 44: 803–U238.
- 23 CLARK, R., T. WAGLER, P. QUIJADA, AND J. DOEBLEY. 2006. A distant

1 upstream enhancer at the maize domestication gene *tb1* has pleiotropic effects on  
2 plant and inflorescent architecture. *Nat Genet* 38: 594–597.

3 COLOSIMO, P., K. HOSEMAN, S. BALABHADRA, G. VILLARREAL,  
4 M. DICKSON, J. GRIMWOOD, J. SCHMUTZ, R. MYERS, D. SCHLUTER, AND  
5 D. KINGSLEY. 2005. Widespread parallel evolution in sticklebacks by repeated  
6 fixation of ectodysplasin alleles. *Science* 307: 1928–1933.

7 COOK, J., M. MCMULLEN, J. HOLLAND, F. TIAN, P. BRADBURY,  
8 J. ROSS-IBARRA, E. BUCKLER, AND S. FLINT-GARCIA. 2012. Genetic  
9 architecture of maize kernel composition in the nested association mapping and  
10 inbred association panels. *Plant Physiol* 158: 824–834.

11 COOP, G., D. WITONSKY, A. DI, RIENZO, AND J. PRITCHARD. 2010. Using  
12 environmental correlations to identify loci underlying local adaptation. *Genetics*  
13 185: 1411–1423.

14 DOEBLEY, J. 2004. The genetics of maize evolution. *Annu Rev Genet* 38: 37–59.

15 DOEBLEY, J. AND A. STEC. 1991. Genetic-analysis of the morphological  
16 differences between maize and teosinte. *Genetics* 129: 285–295.

17 DOEBLEY, J. AND A. STEC. 1993. Inheritance of the morphological differences  
18 between maize and teosinte: Comparison of results for two  $f_2$  populations.  
19 *Genetics* 134: 559–570.

20 DOEBLEY, J., A. STEC, AND C. GUSTUS. 1995. *teosinte branched1* and the origin  
21 of maize: Evidence for epistasis and the evolution of dominance. *Genetics* 141:  
22 333–346.

23 DOEBLEY, J., A. STEC, AND L. HUBBARD. 1997. The evolution of apical  
24 dominance in maize. *Nature* 386: 485–488.

- 1 DOYLE, J. AND J. DOYLE. 1990. A rapid total dna preparation procedure for  
2 small quantities of fresh tissue. *Phytochemical Bulletin* 19: 11–15.
- 3 ELLSTRAND, N., L. GARNER, S. HEDGE, R. GUADAGNUOLO, AND  
4 L. BLANCAS. 2007. Spontaneous hybridization between maize and teosinte.  
5 *Journal of Heredity* 98: 183–187.
- 6 ELLSTRAND, N., H. PRENTICE, AND J. HANCOCK. 1999. Gene flow and  
7 introgression from domesticated plants into their wild relatives. *Annu Rev Ecol*  
8 *Syst* 30: 539–563.
- 9 FALUSH, D., M. STEPHENS, AND J. PRITCHARD. 2003. Inference of population  
10 structure using multilocus genotype data: Linked loci and correlated allele  
11 frequencies. *Genetics* 164: 1567–1587.
- 12 FAO/IIASA/ISRIC/ISSCAS/JRC. 2012. Harmonized World Soil Database,  
13 version 1.2. FAO, Rome, Italy and IIASA, Laxenburg, Austria.
- 14 FEDER, J., S. BERLOCHER, J. ROETHELE, H. DAMBROSKI, J. SMITH,  
15 W. PERRY, V. GAVRILOVIC, K. FILCHAK, J. RULL, AND M. ALUJA. 2003.  
16 Allopatric genetic origins for sympatric host-plant shifts and race formation in  
17 rhagoletis. *P Natl Acad Sci Usa* 100: 10314–10319.
- 18 FOX, J. AND S. WEISBERG. 2011. An R Companion to Applied Regression, vol.  
19 Second Edition. Sage, Thousand Oaks, CA.
- 20 FUKUNAGA, N.-W. T. L. B. Z. Q. V. Y. F. M. B. K. L. L. D. J., K. 2005.  
21 Genetic diversity and population structure of teosinte. *Genetics* 169: 2241–2254.
- 22 GALLAVOTTI, A., Q. ZHAO, J. KYOZUKA, R. MEELEY, M. RITTER,  
23 J. DOEBLEY, M. PE, AND R. SCHMIDT. 2004. The role of barren stalk1 in the  
24 architecture of maize. *Nature* 432: 630–635.

- 1 GERKE, J., J. EDWARDS, G. KE, J. ROSS-IBARRA, AND M. McMULLEN. 2013.  
2 The genomic impacts of drift and selection for hybrid performance in maize.  
3 *arXiv* 1307.7313.
- 4 GOUDET, J. 2005. Hierfstat, a package for r to compute and test hierarchical  
5 f-statistics. *Mol Ecol Notes* 5: 184–186.
- 6 HUBNER, S., T. GUNTHER, A. FLAVELL, E. FRIDMAN, A. GRANER,  
7 A. KOROL, AND K. SCHMID. 2012. Islands and streams: clusters and gene flow  
8 in wild barley populations from the levant. *Mol Ecol* 21: 1115–1129.
- 9 HUFFORD, M. 2010. Genetic and ecological approaches to guide conservation of  
10 teosinte (*zea mays* ssp. *parviglumis*), the wild progenitor of maize. *PhD*  
11 *Dissertation* : 130pp.
- 12 HUFFORD, M., P. BILINSKI, T. PYHÄJÄRVI, AND J. ROSS-IBARRA. 2012a.  
13 Teosinte as a model system for population and ecological genomics. *Trends in*  
14 *Genetics* 12: 606–615.
- 15 HUFFORD, M., P. LUBINSKY, T. PYHÄJÄRVI, M. DEVENGENDO,  
16 N. ELLSTRAND, AND J. ROSS-IBARRA. 2013. The genomic signature of  
17 crop-wild introgression in maize. *PLoS Genetics* 9: e1003477.
- 18 HUFFORD, M., X. XU, J. VAN, HEERWAARDEN, T. PYHÄJÄRVI, J. CHIA,  
19 R. CARTWRIGHT, R. ELSHIRE, J. GLAUBITZ, K. GUILL, S. KAEPPLE,  
20 J. LAI, P. MORRELL, L. SHANNON, C. SONG, N. SPRINGER,  
21 R. SWANSON-WAGNER, P. TIFFIN, J. WANG, G. ZHANG, J. DOEBLEY,  
22 M. McMULLEN, D. WARE, E. BUCKLER, S. YANG, AND J. ROSS-IBARRA.  
23 2012b. Comparative population genomics of maize domestication and  
24 improvement. *Nat Genet* 44: 808–U118.

1 KEARSE, M., R. MOIR, A. WILSON, S. STONES-HAVAS, M. CHEUNG,  
2 S. STURROCK, S. BUXTON, A. COOPER, S. MARKOWITZ, C. DURAN,  
3 T. THIERER, B. ASHTON, P. MEINTJES, AND A. DRUMMOND. 2012. Geneious  
4 basic: An integrated and extendable desktop software platform for the  
5 organization and analysis of sequence data. *Bioinformatics* 28: 1647–1649.

6 KEBROM, T. AND T. BRUTNELL. 2007. The molecular analysis of the shade  
7 avoidance syndrome in the grasses has begun. *Journal of Experimental Botany*  
8 58: 3079–3089.

9 KITANO, J., D. BOLNICK, D. BEAUCHAMP, M. MAZUR, S. MORI, T. NAKANO,  
10 AND C. PEICHEL. 2008. Reverse evolution of armor plates in the threespine  
11 stickleback. *Curr Biol* 18: 769–774.

12 KOVACH, M. AND S. MCCOUCH. 2008. Leveraging natural diversity: back  
13 through the bottleneck. *Genome studies and Molecular Genetics* 11: 193–200.

14 LI, W. 2012. Tassels replace upper ears1 encodes a putative transcription factor  
15 that regulates maize shoot architecture by multiple pathways. *PhD Dissertation* :  
16 122.

17 LUKENS, L. AND J. DOEBLEY. 1999. Epistatic and environmental interactions for  
18 quantitative trait loci involved in maize evolution. *Genet Res* 74: 291–302.

19 MALOOF, M., K. SOLIMAN, R. JORGENSEN, AND R. ALLARD. 1984. Ribosomal  
20 dna spacer length polymorphisms in barley - mendelian inheritance, chromosomal  
21 location, and population dynamics. *P Natl Acad Sci Usa* 81: 8014–8018.


22 MATSUOKA, Y., Y. VIGOUROUX, M. GOODMAN, G. SANCHEZ, E. BUCKLER,  
23 AND J. DOEBLEY. 2002. A single domestication for maize shown by multilocus  
24 microsatellite genotyping. *P Natl Acad Sci Usa* 99: 6080–6084.


- 1 OLSEN, K. AND B. GROSS. 2010. Genetic perspectives on crop domestication.  
2 *Trends in Plant Science* 15: 529–537.
- 3 PIPERNO, D., A. RANERE, I. HOLST, J. IRIARTE, AND R. DICKAU. 2009.  
4 Starch grain and phytolith evidence for early ninth millennium bp maize from the  
5 central balsas river valley, mexico. *P Natl Acad Sci Usa* 106: 5019–5024.
- 6 PLANTINGA, T., S. ALONSO, N. IZAGIRRE, M. HERVELLA, R. FREGEL,  
7 J. VAN DER MEER, M. NETEA, AND C. DE LA RUA. 2012. Low prevalence of  
8 lactase persistence in neolithic south-west europe. *Eur J Hum Genet* 20: 778–782.
- 9 PYHÄJÄRVI, T., M. HUFFORD, AND J. ROSS-IBARRA. 2013. Complex patterns of  
10 local adaptation in the wild relatives of maize. *Genome Biology and Evolution* 5:  
11 1594–1609.
- 12 ROSS-IBARRA, J., P. MORRELL, AND B. GAUT. 2007. Plant domestication, a  
13 unique opportunity to identify the genetic basis of adaptation. *P Natl Acad Sci*  
14 *Usa* 104: 8641–8648.
- 15 ROSS-IBARRA, J., M. TENAILLON, AND B. GAUT. 2009. Historical divergence  
16 and gene flow in the genus *zea*. *Genetics* 181: 1399–1413.
- 17 ROZEN, S. AND H. SKALETSKY. 2000. Primer3 on the www for general users and  
18 for biologist programmers. *Methods in Molecular Biology* : 365–386.
- 19 SIGMON, B. AND E. VOLLBRECHT. 2010. Evidence of selection at the ramosal  
20 locus during maize domestication. *Mol Ecol* 19: 1296–1311.
- 21 STUDER, A. AND J. DOEBLEY. 2012. Evidence for a natural allelic series at the  
22 maize domestication locus *teosinte branched1*. *Genetics* 19: 951–958.



- 1 STUDER, A., Q. ZHAO, J. ROSS-IBARRA, AND J. DOEBLEY. 2011. Identification  
2 of a functional transposon insertion in the maize domestication gene *tb1*. *Nat*  
3 *Genet* 43: 1160–U164.
- 4 THORNTON, K. 2003. libsequence: a c++ class library for evolutionary genetic  
5 analysis. *Bioinformatics* 19: 2325–2327.
- 6 THURBER, C., M. REAGON, B. GROSS, K. OLSEN, Y. JIA, AND A. CAICEDO.  
7 2010. Molecular evolution of shattering loci in us weedy rice. *Mol Ecol* 19:  
8 3271–3284.
- 9 TISHKOFF, S., F. REED, A. RANCIARO, B. VOIGHT, C. BABBITT,  
10 J. SILVERMAN, K. POWELL, H. MORTENSEN, J. HIRBO, M. OSMAN,  
11 M. IBRAHIM, S. OMAR, G. LEMA, T. NYAMBO, J. GHORI, S. BUMPSTEAD,  
12 J. PRITCHARD, G. WRAY, AND P. DELOUKAS. 2007. Convergent adaptation of  
13 human lactase persistence in africa and europe. *Nat Genet* 39: 31–40.
- 14 VAN HEERWAARDEN, J., J. DOEBLEY, W. BRIGGS, J. GLAUBITZ,  
15 M. GOODMAN, J. GONZALEZ, AND J. ROSS-IBARRA. 2011. Genetic signals of  
16 origin, spread, and introgression in a large sample of maize landraces. *P Natl*  
17 *Acad Sci Usa* 108: 1088–1092.
- 18 WEBER, A., R. CLARK, L. VAUGHN, J. SANCHEZ-GONZALEZ, J. YU,  
19 B. YANDELL, P. BRADBURY, AND J. DOEBLEY. 2007. Major regulatory genes  
20 in maize contribute to standing variation in teosinte (*zea mays* ssp *parviglumis*).  
21 *Genetics* 177: 2349–2359.
- 22 WHIPPLE, C., T. KEBROM, A. WEBER, F. YANG, D. HALL, R. MEELEY,  
23 R. SCHMIDT, J. DOEBLEY, T. BRUTNELL, AND D. JACKSON. 2011. grassy

- 1     tillers1 promotes apical dominance in maize and responds to shade signals in the  
2     grasses. *P Natl Acad Sci Usa* 108: E506–E512.
- 3     WHITTON J, A. D. S. A. R. L., WOLF DE. 1997. The persistence of cultivar  
4     alleles in wild populations of sunflowers fiver generations after hybridization.  
5     *Theoretical and Applied Genetcs* 95: 33–40.
- 6     WILKES, H. 1977. Hybridization of maize and teosinte, in mexico and guatemala  
7     and the improvement of maize. *Economic Botany* 31: 254–293.
- 8     ZHANG, L., Q. ZHU, Z. WU, J. ROSS-IBARRA, B. GAUT, S. GE, AND  
9     T. SANG. 2009. Selection on grain shattering genes and rates of rice  
10    domestication. *New Phytol* 184: 708–720.
- 11    Supplementary Materials







| Accession | USDA Accession ID | Locality                   | Number alleles sampled | <i>Hopscotch</i> |
|-----------|-------------------|----------------------------|------------------------|------------------|
| RIHY0009  | N/A               | N/A                        | 2                      |                  |
| RIMME0006 | 566673            | Durango, Mexico            | 2                      |                  |
| RIMME0007 | 566680            | Guanajuato, Mexico         | 2                      |                  |
| RIMME0008 | 566681            | Michoacan, Mexico          | 2                      |                  |
| RIMME0009 | 566682            | Distrito Federal, Mexico   | 2                      |                  |
| RIMME0011 | 566685            | Mexico, Mexico             | 2                      |                  |
| RIMME0014 | 714151            | Breeders line; Puga: 11066 | 6                      |                  |
| RIMME0017 | 699874            | Ayotlan, Mexico            | 8                      |                  |
| RIMME0021 | N/A               | El Porvenir, Mexico        | 69                     | 0                |
| RIMME0026 | N/A               | Opopeo, Mexico             | 42                     | 0                |
| RIMME0028 | N/A               | Puruandiro, Mexico         | 28                     | 0                |
| RIMME0029 | N/A               | Ixtlan, Mexico             | 35                     |                  |
| RIMME0030 | N/A               | San Pedro, Mexico          | 27                     |                  |
| RIMME0031 | N/A               | Tenango del Aire, Mexico   | 25                     |                  |
| RIMME0032 | N/A               | Nabogame, Mexico           | 24                     |                  |
| RIMME0033 | N/A               | Puerta Encantada, Mexico   | 25                     |                  |
| RIMME0034 | N/A               | Santa Clara, Mexico        | 23                     |                  |
| RIMME0035 | N/A               | Xochimilco, Mexico         | 25                     |                  |
| RIMPA0001 | 87168             | El Salado, Mexico          | 4                      |                  |
| RIMPA0003 | 87171             | Mazatlan, Mexico           | 8                      |                  |
| RIMPA0017 | 87200             | N/A                        | 4                      |                  |
| RIMPA0019 | 87213             | El Salado, Mexico          | 2                      |                  |
| RIMPA0029 | 87244             | N/A                        | 2                      |                  |
| RIMPA0031 | 87249             | N/A                        | 2                      |                  |
| RIMPA0035 | 87288             | Jalisco, Mexico            | 4                      |                  |
| RIMPA0040 | 288185            | Mexico, Mexico             | 4                      |                  |
| RIMPA0042 | 288187            | Guerrero, Mexico           | 4                      |                  |
| RIMPA0043 | 288188            | 38 Guerrero, Mexico        | 4                      |                  |
| RIMPA0045 | 288193            | Guerrero, Mexico           | 4                      |                  |
| RIMPA0055 | 714152            | Breeders line              | 2                      |                  |
| RIMPA0056 | 714153            | Breeders line              | 2                      |                  |
| RIMPA0057 | 714154            | Breeders line              | 2                      |                  |
| RIMPA0058 | N/A               | N/A                        | 4                      |                  |

| Accession | Number of alleles sampled | <i>Hopscotch</i> Frequency |
|-----------|---------------------------|----------------------------|
| RIMMA0066 | 2                         | 1                          |
| RIMMA0075 | 2                         | 1                          |
| RIMMA0077 | 2                         | 1                          |
| RIMMA0079 | 2                         | 1                          |
| RIMMA0081 | 2                         | 1                          |
| RIMMA0084 | 2                         | 1                          |
| RIMMA0086 | 2                         | 1                          |
| RIMMA0088 | 2                         | 1                          |
| RIMMA0089 | 2                         | 1                          |
| RIMMA0090 | 2                         | 1                          |
| RIMMA0092 | 4                         | 1                          |
| RIMMA0094 | 4                         | 1                          |
| RIMMA0097 | 2                         | 1                          |
| RIMMA0099 | 2                         | 1                          |
| RIMMA0100 | 2                         | 1                          |
| RIMMA0101 | 2                         | 1                          |
| RIMMA0104 | 2                         | 1                          |
| RIMMA0108 | 2                         | 1                          |
| RIMMA0111 | 6                         | 1                          |
| RIMMA0115 | 2                         | 1                          |
| RIMMA0117 | 2                         | 1                          |
| RIMMA0130 | 2                         | 1                          |
| RIMMA0133 | 2                         | 1                          |
| RIMMA0134 | 2                         | 1                          |
| RIMMA0135 | 2                         | 1                          |
| RIMMA0142 | 2                         | 0.5                        |
| RIMMA0143 | 4                         | 1                          |
| RIMMA0146 | 4                         | 1                          |
| RIMMA0149 | 2                         | 1                          |
| RIMMA0152 | 2                         | 1                          |
| RIMMA0153 | 2                         | 1                          |
| RIMMA0154 | 2                         | 1                          |
| RIMMA0155 | 2                         | 1                          |

| PC1      |       | PC2     |       | PC3     |       | PC4     |       | PC5     |       | PC6     |       |
|----------|-------|---------|-------|---------|-------|---------|-------|---------|-------|---------|-------|
| Var      | Rot   | Var     | Rot   | Var     | Rot   | Var     | Rot   | Var     | Rot   | Var     | Rot   |
| bio1     | 0.146 | bio4    | 0.244 | prec7   | 0.287 | ts_clay | 0.41  | bio2    | 0.38  | bio10   | 0.086 |
| tmean11  | 0.146 | bio3    | 0.241 | prec8   | 0.276 | v_mod   | 0.359 | sq4     | 0.328 | x_mod   | 0.111 |
| tmean12  | 0.145 | bio7    | 0.241 | prec11  | 0.262 | ts_sand | 0.329 | ts_loam | 0.289 | sq3     | 0.207 |
| bio11    | 0.145 | prec6   | 0.237 | bio13   | 0.247 | bio15   | 0.272 | ts_sand | 0.266 | bio1    | 0.146 |
| tmax12   | 0.145 | sq7     | 0.218 | prec1   | 0.246 | prec4   | 0.259 | sq7     | 0.231 | v_mod   | 0.359 |
| tmin5    | 0.145 | prec9   | 0.217 | bio16   | 0.242 | x_mod   | 0.244 | bio18   | 0.213 | prec1   | 0.287 |
| tmean1   | 0.145 | sq3     | 0.207 | prec12  | 0.24  | prec3   | 0.226 | bio13   | 0.207 | bio1    | 0.146 |
| tmean2   | 0.145 | prec12  | 0.207 | bio19   | 0.238 | sq3     | 0.21  | prec11  | 0.183 | sq4     | 0.328 |
| tmin4    | 0.145 | bio12   | 0.204 | bio12   | 0.231 | prec5   | 0.21  | bio7    | 0.17  | sq3     | 0.207 |
| tmax1    | 0.145 | bio19   | 0.196 | prec2   | 0.222 | prec7   | 0.19  | bio16   | 0.163 | ts_sand | 0.329 |
| tmean4   | 0.145 | prec2   | 0.188 | bio18   | 0.221 | sq4     | 0.186 | bio4    | 0.157 | bio4    | 0.244 |
| tmin11   | 0.144 | prec1   | 0.185 | sq4     | 0.2   | bio3    | 0.185 | bio12   | 0.156 | prec1   | 0.287 |
| tmax11   | 0.144 | prec10  | 0.184 | prec9   | 0.18  | bio18   | 0.178 | bio3    | 0.155 | tmax1   | 0.145 |
| tmin12   | 0.144 | bio16   | 0.183 | prec10  | 0.171 | sq7     | 0.132 | prec6   | 0.154 | tmax1   | 0.145 |
| tmin2    | 0.144 | prec8   | 0.17  | prec5   | 0.161 | bio14   | 0.116 | x_mod   | 0.152 | bio1    | 0.146 |
| tmean5   | 0.144 | prec5   | 0.165 | prec4   | 0.154 | bio13   | 0.099 | prec9   | 0.144 | tmax1   | 0.145 |
| tmean10  | 0.144 | bio14   | 0.158 | sq3     | 0.147 | bio16   | 0.095 | prec8   | 0.143 | bio1    | 0.146 |
| bio6     | 0.144 | bio13   | 0.151 | bio2    | 0.143 | prec8   | 0.09  | v_mod   | 0.142 | ts_loam | 0.289 |
| tmax2    | 0.144 | bio17   | 0.149 | bio17   | 0.129 | bio7    | 0.077 | bio15   | 0.136 | ts_clay | 0.41  |
| tmean3   | 0.144 | prec3   | 0.144 | ts_loam | 0.127 | bio4    | 0.075 | prec7   | 0.112 | tmin1   | 0.143 |
| tmin1    | 0.143 | ts_clay | 0.141 | v_mod   | 0.123 | bio2    | 0.074 | prec4   | 0.108 | tmin1   | 0.143 |
| tmin10   | 0.143 | bio2    | 0.129 | prec3   | 0.113 | prec2   | 0.074 | bio14   | 0.096 | prec1   | 0.287 |
| Altitude | 0.143 | prec7   | 0.108 | x_mod   | 0.111 | bio19   | 0.068 | tmax7   | 0.093 | tmin1   | 0.143 |
| bio9     | 0.143 | tmax6   | 0.107 | bio14   | 0.099 | prec12  | 0.056 | tmax8   | 0.092 | tmin1   | 0.143 |
| tmin3    | 0.143 | x_mod   | 0.106 | bio4    | 0.07  | ts_loam | 0.053 | prec1   | 0.091 | tmin1   | 0.143 |
| bio10    | 0.142 | bio15   | 0.098 | tmax3   | 0.067 | tmax12  | 0.047 | prec2   | 0.086 | tmean1  | 0.145 |
| tmax10   | 0.142 | ts_loam | 0.088 | ts_clay | 0.065 | bio17   | 0.047 | tmin11  | 0.086 | tmax1   | 0.145 |
| tmax3    | 0.142 | tmean6  | 0.085 | bio15   | 0.056 | bio9    | 0.043 | prec5   | 0.082 | tmax1   | 0.145 |
| tmax4    | 0.142 | tmin7   | 0.082 | tmax2   | 0.055 | tmax8   | 0.042 | bio17   | 0.082 | tmean1  | 0.145 |
| tmin6    | 0.142 | bio5    | 0.082 | tmean3  | 0.052 | tmax1   | 0.041 | tmin12  | 0.08  | bio1    | 0.146 |
| tmean9   | 0.141 | tmean7  | 0.081 | ts_sand | 0.05  | tmax5   | 0.039 | prec3   | 0.078 | tmin1   | 0.143 |
| tmin9    | 0.141 | prec4   | 0.08  | prec6   | 0.048 | tmax7   | 0.039 | tmax9   | 0.078 | prec1   | 0.287 |



|             |   |         |         |         |
|-------------|---|---------|---------|---------|
| Ejutla A    | 4 | 0.15217 | 0.11902 | 0.76191 |
| Ejutla B    | 5 | 0.15258 | 0.14877 | 0.07412 |
| La Mesa     | 3 | 0.12802 | 0.08926 | 1.09209 |
| San Lorenzo | 3 | 0.09098 | 0.08926 | 0.04845 |