

Gradient term for the 4th-order symplectic integrator

似鳥啓吾

2018 年 11 月 2 日

The Poisson bracket is

$$\{A, B\} = \sum_i \left(\frac{\partial A}{\partial q_i} \frac{\partial B}{\partial p_i} - \frac{\partial A}{\partial p_i} \frac{\partial B}{\partial q_i} \right), \quad (1)$$

and for a Hamiltonian $H = T(p) + V(q)$,

$$\{X, V\} = \sum_i \left(\frac{\partial X}{\partial q_i} \cdot F_i \right), \quad (2)$$

where $F_i = -\partial V / \partial q_i$ is force. Thus,

$$\{T, V\} = \sum_i \left(\frac{p_i}{m_i} \cdot F_i \right), \quad (3)$$

and

$$\{\{T, V\}, V\} = \sum_j \left[F_j \cdot \frac{\partial}{\partial p_j} \left(\frac{p_i}{m_i} \cdot F_i \right) \right] = \sum_j \left(\frac{F_j \cdot F_j}{m_j} \right), \quad (4)$$

For the 4th-order forward symplectic integrator, we evaluate the gradient term,

$$\mathbf{G}_i = -\frac{\partial}{\partial \mathbf{r}_i} \left[\sum_{j=1}^N \frac{\mathbf{F}_j \cdot \mathbf{F}_j}{m_j} \right] = -2 \sum_{j=1}^N \left[\left(\frac{\partial}{\partial \mathbf{r}_i} \mathbf{F}_j \right) \cdot \mathbf{a}_j \right]. \quad (5)$$

Here, \mathbf{F}_i is force on particle i and \mathbf{F}_{ij} contribution from particle j , i.e.,

$$\mathbf{F}_i = \sum_{j \neq i}^N \mathbf{F}_{ij} = \sum_{j \neq i}^N \frac{G m_i m_j}{|\mathbf{r}_j - \mathbf{r}_i|^3} (\mathbf{r}_j - \mathbf{r}_i). \quad (6)$$

and \mathbf{a}_i is acceleration \mathbf{F}_i / m . Thus,

$$\frac{\partial}{\partial \mathbf{r}_i} \mathbf{F}_j = \frac{\partial}{\partial \mathbf{r}_i} \left[\sum_{k \neq j}^N \mathbf{F}_{jk} \right] = \begin{cases} \sum_{k \neq i}^N \frac{\partial}{\partial \mathbf{r}_i} \mathbf{F}_{ik} & (i = j) \\ \frac{\partial}{\partial \mathbf{r}_i} \mathbf{F}_{ji} = -\frac{\partial}{\partial \mathbf{r}_i} \mathbf{F}_{ij} & (i \neq j) \end{cases}. \quad (7)$$

The summation remains only in the diagonal term and disappears elsewhere.

$$\mathbf{G}_i = -2 \sum_{j \neq i}^N \left[\frac{\partial}{\partial \mathbf{r}_i} \mathbf{F}_{ij} \right] \cdot (\mathbf{a}_i - \mathbf{a}_j) \quad (8)$$

For the N -body system, gradient of mutual force in 3×3 matrix is given in,

$$\frac{\partial}{\partial \mathbf{r}_i} \frac{(\mathbf{r}_j - \mathbf{r}_i)}{|\mathbf{r}_j - \mathbf{r}_i|^3} = \frac{-I}{|\mathbf{r}_j - \mathbf{r}_i|^3} + \frac{3(\mathbf{r}_j - \mathbf{r}_i) \otimes (\mathbf{r}_j - \mathbf{r}_i)}{|\mathbf{r}_j - \mathbf{r}_i|^5}, \quad (9)$$

where I is a unit matrix.

Finally we have

$$\mathbf{G}_i = -2Gm_i \sum_{j \neq i}^N m_j \left[\frac{(\mathbf{a}_j - \mathbf{a}_i)}{|\mathbf{r}_j - \mathbf{r}_i|^3} - \frac{3(\mathbf{r}_j - \mathbf{r}_i) \cdot (\mathbf{a}_j - \mathbf{a}_i)}{|\mathbf{r}_j - \mathbf{r}_i|^5} (\mathbf{r}_j - \mathbf{r}_i) \right]. \quad (10)$$

One can just replace the velocity by the force in the jerk formula to compute it. Note that $\mathbf{G}_i h^2$ has a dimension of force.

1 Materials

$$\tilde{\mathbf{F}}_i = \mathbf{F}_i + \frac{h^2}{48} \frac{1}{m_i} \frac{\partial}{\partial \mathbf{r}_i} |\mathbf{F}|^2$$

$$K(\tfrac{1}{6}h)D(\tfrac{1}{2}h)\tilde{K}(\tfrac{2}{3}h)D(\tfrac{1}{2}h)K(\tfrac{1}{6}h)$$

$$\begin{aligned} & \begin{bmatrix} \frac{\partial}{\partial_1} \\ \frac{\partial}{\partial_2} \\ \frac{\partial}{\partial_3} \end{bmatrix} \begin{pmatrix} F_1 & F_2 & F_3 \end{pmatrix} \begin{bmatrix} F_1 \\ F_2 \\ F_3 \end{bmatrix} \\ &= \begin{pmatrix} \frac{\partial}{\partial_1}(F_{12} + F_{13}) & \frac{\partial}{\partial_1}F_{21} & \frac{\partial}{\partial_1}F_{31} \\ \frac{\partial}{\partial_2}F_{12} & \frac{\partial}{\partial_2}(F_{23} + F_{21}) & \frac{\partial}{\partial_2}F_{32} \\ \frac{\partial}{\partial_3}F_{13} & \frac{\partial}{\partial_3}F_{23} & \frac{\partial}{\partial_3}(F_{31} + F_{32}) \end{pmatrix} \begin{pmatrix} F_1 \\ F_2 \\ F_3 \end{pmatrix} \\ &= \begin{pmatrix} (\frac{\partial}{\partial_1}F_{12})(F_1 - F_2) + (\frac{\partial}{\partial_1}F_{13})(F_1 - F_3) \\ (\frac{\partial}{\partial_2}F_{23})(F_2 - F_3) + (\frac{\partial}{\partial_2}F_{21})(F_2 - F_1) \\ (\frac{\partial}{\partial_3}F_{31})(F_3 - F_1) + (\frac{\partial}{\partial_3}F_{32})(F_3 - F_2) \end{pmatrix} \end{aligned}$$

$$|\mathbf{F}_{\text{hard}} + \mathbf{F}_{\text{soft}}|^2 - |\mathbf{F}_{\text{hard}}|^2 = |\mathbf{F}_{\text{soft}}|^2 + 2\mathbf{F}_{\text{hard}} \cdot \mathbf{F}_{\text{soft}}$$

2 P³T

Poisson bracket is

$$\{A, B\} = \frac{\partial A}{\partial q} \frac{\partial B}{\partial p} - \frac{\partial B}{\partial q} \frac{\partial A}{\partial p}.$$

We split the Hamiltonian into a hard part and a soft part,

$$H = \underbrace{(T + V_H)}_{\text{hard}} + \underbrace{V_S}_{\text{soft}}.$$

Now,

$$\{T + V_H, V_S\} = \frac{p \cdot F_S}{m} \quad (11)$$

$$\{\{T + V_H, V_S\}, V_S\} = \frac{F_S \cdot F_S}{m} \quad (12)$$

$$\{V_S, T + V_H\} = -\frac{p \cdot F_S}{m} \quad (13)$$

$$\begin{aligned} \{\{V_S, T + V_H\}, T + V_H\} &= -\left\{ \frac{p}{m} \cdot F_S, T + V_H \right\} \\ &= -\left(\left[\frac{p}{m} \cdot \frac{\partial F_S}{\partial q} \right] \frac{p}{m} + F_S \cdot F_H \right) \end{aligned} \quad (14)$$

The leading error term is (12).