

Update on $t\bar{t}t\bar{t}$ Searches in Single Lepton/OS Dilepton Channels Using 2016 Data

Denys Lontkovskyi ¹ Freya Blekman ¹ Long Wang ²
Robert Clare ² Steve Wimpenny ²

¹Vrije Universiteit Brussel

²University of California, Riverside

March 23, 2018

Overview

1. Current status
2. Data and event selection
3. $t\bar{t}t\bar{t}$ Search Method
4. Re-blinded fit result
5. Conclusion

Current Status

- ▶ Aiming at re-preapproval, we are finalizing our documentation
 - ▶ Single-lepton: AN-17/146
 - ▶ Di-lepton: AN-17/143
- ▶ We have requested for the production of two new $t\bar{t}$ samples with 9M events each for the two channels(semi-lep and OS dilep), with dedicated cuts at generator level to increase MC stats by a factor of ~ 10 in high multiplicity/discriminant tails. Details on slide ?? to ??
- ▶ We are studying the effects of possible background from QCD multi jets with mis-identified leptons.

Data, MC and Objects

Data and MC

- ▶ Run2016 B-H, $35.9 pb^{-1}$
- ▶ Summer 16 MiniAOD MC for Morond 17
 - ▶ signal sample: $t\bar{t}t\bar{t}$ amc@NLO
 - ▶ background samples: $t\bar{t}$ (+backup, mass, width, details on slide ??), single $t(\bar{t})$, DY, W+jets, $t\bar{t} + Z/H/W/diboson$

Objects

Single Lepton

- ▶ μ : tight ID, $p_T > 26$ GeV, $|\eta| < 2.1$, $R_{\ell\ell so} < 0.15$
- ▶ e : tight ID, $p_T > 35$ GeV, $|\eta| < 1.4442$ or $1.566 < |\eta| < 2.1$
- ▶ jet : loose ID, $p_T > 30$ GeV, $|\eta| < 2.1$, $\Delta R > 0.4$

OS Dilepton

- ▶ μ : loose ID, leading(subleading) lep $p_T > 25(20)$ GeV, $|\eta| < 2.4$, $R_{\ell\ell so} < 0.15$
- ▶ e : loose ID, leading(subleading) lep $p_T > 25(20)$ GeV, $|\eta| < 1.4442$ or $1.566 < |\eta| < 2.4$
- ▶ jet : loose ID, $p_T > 30$ GeV(25 GeV if tagged as b), $|\eta| < 2.4$, $\Delta R > 0.4$

Event Selection and MC Re-weighting

Event selection

Single Lepton

- ▶ $N_l^{tight}=1$
- ▶ $N_\mu^{loose} = 0, N_e^{veto} = 0$
- ▶ $N_j \geq 8(7)$ in $e(\mu)$ channel of which $N_{tags}^M \geq 2$
- ▶ $\cancel{E}_T > 50$ GeV
- ▶ $HT \geq 500$ GeV

OS Dilepton

- ▶ Exactly 2 OS leptons
- ▶ $M_{ll} \geq 106$ GeV or $76 \geq M_{ll} \geq 20$ GeV
- ▶ $N_j \geq 4$ of which $N_{tags}^M \geq 2$
- ▶ $HT \geq 500$ GeV

MC Re-weighting

- | | | |
|-----------------|-------------------|----------------------|
| ▶ Trigger eff. | ▶ Pileup Reweight | ▶ b-tagging eff. |
| ▶ Lepton scales | ▶ JER/JEC | ▶ top p_T reweight |

$t\bar{t}t\bar{t}$ Search Method

Binned analysis fitting on event level BDT




Figure: Single μ event level BDT in ≥ 9 jet 3 btag category





Figure: μe event level BDT in CR

Event categorization in $N_j \otimes N_{tags}^M$ for limit fitting

- ▶ Single lepton channel
 - ▶ μ : N_j : 7, 8, 9, 10+; N_{tags}^M : 2, 3, 4+
 - ▶ e : N_j : 8, 9, 10+; N_{tags}^M : 2, 3, 4+
- ▶ OS Dilepton channel: N_j : 4-5, 6-7, 8+; N_{tags}^M : 2, 3+


Control Plots (BDT Input Variables)

► Single lepton channel



► OS Dilepton channel

$$N_j, BDT_{trijet1}, H_T^b, H_T^{2M}, HTH, S, H_T^{Rat}, p_T^{l1}, \eta^{l1}, \Delta R_{ll}, \Delta R_{bb}, N_{tags}^L, N_{tags}^M, p_T^{3rdjet}, p_T^{4thjet}$$



► Overall reasonable distributions agreement within uncertainties.

Sources of Systematic Uncertainties

Experimental Uncertainties

- ▶ Luminosity uncertainty
- ▶ Pileup $\pm 1\sigma$
- ▶ Lepton SFs uncertainty
- ▶ JER $\pm 1\sigma$
- ▶ JES(split)
 - ▶ SubTotalPileUp
 - ▶ SubTotalRelative
 - ▶ SubTotalPt
 - ▶ SubTotalScale
 - ▶ Jet flavor
- ▶ b-tag CSV $\pm 1\sigma$
- ▶ Heavy flavor fraction
- ▶ Top p_T reweight
- ▶ Jet normalization


Theoretical Uncertainties

- ▶ ME scale
- ▶ MC cross sections
- ▶ UE tune
- ▶ PS scale
- ▶ ME-PS matching
- ▶ PDF

Fit Strategy



- ▶ Likelihood fit using Combine Tool
- ▶ Event level BDT output discriminator distributions for fit is performed simultaneously in different $N_j \otimes N_{tags}^M$ categories.
- ▶ Blind highest jet/tag multiplicity categories.
 - ▶ single lepton: blind 10+ jets & 3+ tags category
 - ▶ OS dilepton: blind 8+ jets & 3+ tags category
- ▶ Combine results from single lepton channel and OS dilepton channel.

- ▶ Large uncertainties on MC in the signal regions.



Template Fit in Single Lepton Channel

Postfit BDT distributions



- Equiprobable binning scheme
- Blind signal rich 10+/3+ category
- Reasonable description of the data in CRs

Template Fit in Single Lepton Channel

Table: Single lepton blinded fitting results

| Channel | Expected limit $\times \sigma_{t\bar{t}t\bar{t}}^{SM}$ | Expected xsec fb | Expected significance |
|----------|---|-----------------------|--------------------------|
| e | $23.5^{+7.0}_{-6.5}$ | 216.2^{+64}_{-60} | 0.09 |
| μ | $16.0^{+7.0}_{-4.7}$ | 147.2^{+64}_{-43} | 0.12 |
| combined | $9.4^{+4.0}_{-2.7}$ | 86.5^{+37}_{-25} | 0.25 |

Fit Diagnostic in Single Lepton Channel




Figure: μ +jets channel





Figure: $e+\text{jets}$ channel

- Post-fit uncertainty reduction is under investigation

Impact of Nuisance Parameters



- ▶ Dominant sources of systematic uncertainty:
 - ▶ UE variation. Affects jet multiplicity spectrum (Sample has low statistics)
 - ▶ $t\bar{t}b\bar{b}$ normalization
 - ▶ Normalization of $t\bar{t}Z, H \rightarrow b\bar{b}$
 - ▶ Reweighting of HF component in CSV discriminant

Postfit BDT distributions

- ▶ high BDT region in signal sensitive regions is blinded
- ▶ Obvious excess in the distribution tails within all three channels, could be due to lack of MC stats




Figure: $\mu\mu$ channel




Figure: $e\mu$ channel




Figure: ee channel

Template Fit in OS Dilepton Channel

Table: OS dilepton blinded fitting results

| Channel | Expected limit $\times \sigma_{t\bar{t}t\bar{t}}^{SM}$ | Expected xsec fb | Expected significance |
|----------|---|-----------------------|--------------------------|
| $\mu\mu$ | $14.56^{+9.64}_{-5.24}$ | 134^{+89}_{-48} | 0.19 |
| $e\mu$ | $9.88^{+6.53}_{-3.53}$ | 91^{+60}_{-32} | 0.37 |
| ee | $17.56^{+11.34}_{-6.19}$ | 162^{+104}_{-57} | 0.29 |
| combined | $6.88^{+4.44}_{-2.42}$ | 63^{+41}_{-22} | 0.52 |

Fit Diagnostic in OS Dilepton Channel




Figure: nuisance pulls in $\mu\mu$




Figure: nuisance pulls in $e\mu$

- ▶ Most signal sensitive region is blinded
- ▶ No extreme pulls or constraints.
- ▶ Reasonable behavior for all the NPs
- ▶ Results are consistent between three sub-channels

Fit Diagnostic in OS Dilepton Channel




Figure: nuisance pulls in ee




Figure: nuisance pulls in OS dilep

- ▶ Mainly the sub-components of JES are constrained
 - ▶ Statistic fluctuation
 - ▶ Correlated with other NPs

Nuisance Parameters impact in OS dilep combined fit




Figure: Impact of nuisance parameters on the parameter of interest

Nuisance Parameters impact in OS dilep combined fit

- ▶ Signal systematics have the largest impacts, as we are very close to the expected signal strength.
- ▶ TTRare resembles our signal, but under control.
- ▶ Jet energy scale uncertainties and MC stats in signal enriched bins dominate.
- ▶ All nuisance parameters behave reasonably.
- ▶ Full list of nuisance impacts are on backup slides ?? to ??

Combined Results

Table: Single lepton + OS dilepton blinded combined results

| Channel | Expected limit $\times \sigma_{t\bar{t}t\bar{t}}^{SM}$ | Expected limit fb | Expected significance |
|------------|---|------------------------|--------------------------|
| l+jets | $9.4^{+4.0}_{-2.7}$ | 86.5^{+37}_{-25} | 0.25 |
| OS ll+jets | $6.9^{+4.4}_{-2.4}$ | 63^{+41}_{-22} | 0.52 |
| combined | $5.2^{+2.6}_{-1.7}$ | 48^{+24}_{-16} | 0.58 |

- Signal sensitivity is driven by the OS ll+jets channel

Conclusion

- ▶ Analysis re-blinded in order to optimize search regions/binning taking into account limited MC statistics
- ▶ Expect $\times 10$ more $t\bar{t}$ simulated events using dedicated filters in signal rich search regions that have large statistical uncertainty
 - ▶ New high jet multiplicity, high- H_T filter was optimized for the analysis
 - ▶ MC requests are added to the system

Plans for coming weeks:

- ▶ Include new MC samples when they are ready
- ▶ Finalize QCD background estimation
- ▶ Update ANs and paper draft

Backups

MC samples

Table: Simulated event samples, cross sections and event yields used in this analysis

| Sample | σ (pb) | No of events |
|---|---------------|--------------|
| /TT_TuneCUETP8M2T4_13TeV-powheg-pythia8(& backup) | 831.8 | 154652276 |
| /TT_TuneCUETP8M2T4_mtop1665_13TeV-powheg-pythia8 | 831.8 | 19379727 |
| /TT_TuneCUETP8M2T4_mtop1695_13TeV-powheg-pythia8 | 831.8 | 58540996 |
| /TT_TuneCUETP8M2T4_mtop1715_13TeV-powheg-pythia8 | 831.8 | 19578294 |
| /TT_TuneCUETP8M2T4_mtop1735_13TeV-powheg-pythia8 | 831.8 | 19323035 |
| /TT_TuneCUETP8M2T4_mtop1755_13TeV-powheg-pythia8 | 831.8 | 59252096 |
| /TT_TuneCUETP8M2T4_mtop1785_13TeV-powheg-pythia8 | 831.8 | 16376678 |
| /TT_widthx0p2_TuneCUETP8M2T4_13TeV-powheg-pythia8 | 831.8 | 19937607 |
| /TT_widthx0p5_TuneCUETP8M2T4_13TeV-powheg-pythia8 | 831.8 | 19909415 |
| /TT_widthx0p8_TuneCUETP8M2T4_13TeV-powheg-pythia8 | 831.8 | 18546839 |
| /TT_widthx2_TuneCUETP8M2T4_13TeV-powheg-pythia8 | 831.8 | 13505281 |
| /TT_widthx4_TuneCUETP8M2T4_13TeV-powheg-pythia8 | 831.8 | 19593910 |
| /TT_widthx8_TuneCUETP8M2T4_13TeV-powheg-pythia8 | 831.8 | 19524579 |
| total | | 436M |

Current status

- ▶ Existing inclusive $t\bar{t}$ sample, including mass and width variations: 436M events
 - ▶ Nevertheless, MC statistics is scarce
 - ▶ 200 MC events in 10j4b l^+l^- +jets category
 - ▶ 140 MC events in 8j4j l^+l^- +jets category
 - ▶ Need factor of 10 increase in statistics to ensure fit stability in the sensitive high multiplicity/high discriminant region
 - ▶ This requires the development of a gen level filter for efficient MC production
- ▶ Various filter configurations have been studied and we have a preferred configuration
- ▶ MC production can be started once new release is available
- ▶ Backport and genfragment are available
(Thanks Javier Fernandez!)
- ▶ CMSSW modification are signed by generator conveners
(Thanks L.Perrozzi!)
- ▶ Currently waiting for new cmssw build containing the backport
 - ▶ New patch release has to be issued by release managers

Two scenarios for new filtered samples

- ▶ Two independent samples with different lepton and jet multiplicity cuts:
 - ▶ Optimal for the two channels
 - ▶ Uses gen level cuts which are probably not useful for other analyses
 - ▶ Common sample with no fully hadronic $t\bar{t}$ decays
 - ▶ Potentially useful for other analyses
 - ▶ Worse filter efficiency
-
- ▶ Filter efficiency and acceptance loss were calculated on /MINIAODSIM level
 - ▶ Efficiency values were verified using actual filter implementation in small private /GEN-SIM production

Filters summary (l+jets and OS)

- Preferred configuration: $HT > 500$, $n\text{Jets} + n\text{Lep} \geq 8$, $n\text{Lep} \geq 1$ (26M)

| Filter cuts | Filter eff. $\times 10^{-2}$ | Acceptance loss in different jet multiplicity regions ¹ | | | | |
|---|----------------------------------|--|-----------------|--------------|--------------|---------------------|
| | | OS $N_J = 7$ | OS $N_J \geq 8$ | SL $N_J = 9$ | SL $N_J > 9$ | Ext ($\times 10$) |
| $HT > 500$ $n\text{Jets} + n\text{Lep} \geq 6$ ≥ 1 lepton | 2.3 ± 0.03 | — | — | — | — | 100 M |
| $HT > 500$ $n\text{Jets} + n\text{Lep} \geq 7$ ≥ 1 lepton | 1.4 ± 0.02 | 0.04 | 0.02 | — | — | 61 M |
| $HT > 500$ $n\text{Jets} + n\text{Lep} \geq 8$ ≥ 1 lepton | 0.6 ± 0.02 | 0.14 | 0.04 | 0.11 | 0.08 | 26 M |
| $HT > 500$ $n\text{Jets} + n\text{Lep} \geq 9$ ≥ 1 lepton | 0.2 ± 0.01 | 0.43 | 0.14 | 0.19 | 0.1 | 9M |
| $HT > 500$ $n\text{Jets} + n\text{Lep} \geq 10$ ≥ 1 lepton | 0.08 ± 0.007 | — | — | — | 0.19 | 3.5M |

¹Fraction of events passing offline cuts but rejected by gen filter

Filter optimized for l+jets channel

- Preferred configuration: $HT > 500$, $n\text{Jets} + n\text{Lep} \geq 9$, $n\text{Lep} = 1$ (9M)

| Filter cuts | Filter eff. | Acceptance loss in different jet multiplicity regions ² | | |
|---|---------------------------------|--|-------------------------------|---------------------|
| | | SL ($N_J^{\text{rec}} = 9$) | SL ($N_J^{\text{rec}} > 9$) | Ext ($\times 10$) |
| HT>500 nJets+nLep ≥ 8 1 lepton | 0.005 ± 0.0002 | 0.11 | 0.08 | 21.8 M |
| HT > 500 nJets+nLep ≥ 9 1 lepton | 0.002 ± 0.0001 | 0.19 | 0.10 | 8.7M |
| HT>500 nJets+nLep ≥ 10 1 lepton | $0.0007 \pm 6.5 \times 10^{-5}$ | — | 0.19 | 3M |

²Fraction of events passing offline cuts but rejected by gen filter

Filter optimized for OS dilepton channel

- Preferred configuration: $HT > 500$, $n\text{Jets} + n\text{Lep} \geq 7$, $n\text{Lep} = 2$ (9M)

| Filter cuts | Filter eff. | Acceptance loss in different jet multiplicity regions ³ | | |
|-------------------------------------|------------------------|--|----------------------------------|---------------------|
| | | OS ($N_j^{\text{rec}} = 7$) | OS ($N_j^{\text{rec}} \geq 8$) | Ext ($\times 10$) |
| HT>500 nJets+nLep>=5 2 lepton | $0.0046 \pm 5.5e - 05$ | 0.21 | 0.23 | 20 M |
| HT>500 nJets+nLep>=6 2 lepton | $0.0033 \pm 4.7e - 05$ | 0.21 | 0.23 | 14.4 M |
| HT>500 nJets+nLep>=7 2 lepton | $0.0020 \pm 3.6e - 05$ | 0.23 | 0.24 | 8.7 M |
| HT>500 nJets+nLep>=8 2 lepton | $0.0009 \pm 2.4e - 05$ | 0.31 | 0.25 | 3.9 M |

³Fraction of events passing offline cuts but rejected by gen filter

Summary

- ▶ There are two options to get $10\times$ MC stat extension:
 - ▶ (Preferred) Two samples with different gen level cuts
(Total of 18M events passing the filters):
 - ▶ **OS 9M events:** $HT > 500, nJets + nLep \geq 7, nLep = 2$
 - ▶ **SL 9M events:** $HT > 500, nJets + nLep \geq 9, nLep = 1$
 - ▶ Combined sample
(Total of 26M events passing the combined filter):
 - ▶ **SL+OS:** $HT > 500, nJets + nLep \geq 8, nLep \geq 1$

Definitions

- ▶ Variable definitions:

- ▶ $HT = \sum_{j: p_T > 30, |\eta| < 2.4} p_T(j)$




- ▶ nJets= number of gen jets with $p_T > 30$

- No $\eta(j)$ cut due to different CMSSW filter logic.

- ▶ Jets in HT and multiplicity definition may include jets from isolated leptons

- ▶ Lepton cuts are applied using LHE filter, similar to NoFullyHadronicDecays filter applied to single-top MC samples

impacts of nuisance parameters



impacts of nuisance parameters

