

MATH 252 - Introduction to Differential Equations Notes

Len Washington III

September 11, 2023

Contents

1	Introduction to Diff-Eq	3
1.1	Terminology and Notation	3
1.1.1	Linear vs Non-Linear DE's	3
1.2	Initial Value Problems (IVP)	4
1.2.1	Example	5
1.2.2	Example	5
1.2.3	Example	5
1.2.4	Example	6
1.2.5	Example	6
2	First-Order Differential Equations	7
2.1	Solution Curves Without a Solution	7
2.1.1	Example	7
2.1.2	Slope/Direction Fields	8
2.1.3	Example	8
2.2	Separable D.E.s	9
2.2.1	Method of Solution	10
2.2.2	Example	10
2.2.3	Example	10
2.3	First Order Linear D.E.'s	11
2.3.1	Example	12
2.3.2	Example	13
2.4	Exact Equations	13
2.4.1	Method	14
2.4.2	Example	14
2.4.3	Example	15
2.4.4	What can you do if $M_y \neq N_x$	16
2.4.5	Example	16
2.5	Substitution Methods	17
2.5.1	Example	17
2.5.2	Example	18
2.5.3	Substitution Rule	18
2.5.4	Example	18
2.5.5	Example	20
2.5.6	Example	22

3	Modeling using DE	23
3.1	Linear DE Modeling	23
3.1.1	Standard Problems	23
3.1.2	Population Model	23
3.1.3	Example	24
3.1.4	Radioactive Decay	25
3.1.5	Mixture Problems	26
4	Higher Order DEs	27
4.0.1	Example	27

Chapter 1

Introduction to Differential Equations

1.1 Terminology and Notation

Differential equation (D.E.) – An equation in which at least one derivative of an unknown function.

Order of the D.E. – The highest order of derivative in the D.E.

Example:

$$4y'' + e^x y' - 3yy' = \sin(x)$$

An example of a partial differential equation is:

$$\frac{\partial T}{\partial x} + x^2 \frac{\partial T}{\partial y} = x + y$$

however, we won't study these in this course.

1.1.1 – Linear vs Non-Linear DE's

Linear D.E. – The dependent variable and all of its derivatives in the D.E. are in separate terms to the 1st power. $y^{(n)}$ or $\frac{d^n y}{dx^n}$ where $n \neq 1$ are non-first power.

$$4y'' + e^x y' - 3yy' = \sin(x)$$

is a non-linear D.E. while

$$4y'' + e^x y' - 3y = \sin(x)$$

is linear.

The general formula of a linear D.E. would look like

$$a_n(x)y^{(n)} + a_{n-1}(x)y^{(n-1)} + \cdots + a_1(x)y' + a_0(x) = g(x)$$

Solution – a function $\phi(x)$ and an interval I for which the D.E. is satisfied when $y = \phi(x)$ for all x in I .

It may be the case that the natural domain of $\phi(x)$ is larger than I . Example: $y' = -\frac{1}{x^2}$ has a solution $\phi(x) = \frac{1}{x}$ on $I = (0, \infty)$ but the domain of $\phi(x) = (-\infty, 0) \cup (0, \infty)$. Practice:

$$\frac{d^2x}{dt^2} + 16x = 0$$

Show (Verify not derive) $x(t) = c_1 \sin(4t)$ is a solution on $(-\infty, \infty)$ where c is any real parameter.

$$\begin{aligned} x &= c_1 \sin(4t) \\ \frac{dx}{dt} &= 4c_1 \cos(4t) \\ \frac{d^2x}{dt^2} &= -16c_1 \sin(4t) \\ \text{LHS} &= \frac{d^2x}{dt^2} + 16x \\ &= -16c_1 \sin(4t) + 16(c_1 \sin(4t)) \\ &= 0 = \text{RHS} \end{aligned}$$

But the equation $x = c_2 \cos(4t)$ would also be a solution. If you have 2 equations that are both solutions, you could add them together and you would still have a solution. $x = c_1 \sin(4t) + c_2 \cos(4t)$ is a solution for all parameters c_1 and c_2 . *In fact, this is the general solution to the D.E.*

The D.E.

$$\frac{dy}{dx} = xy^{\frac{1}{2}}$$

Show $y = \left(\frac{1}{4}x^2 + C\right)^2$ is a one parameter family of solutions

$$\begin{aligned} \text{LHS} &= \frac{dy}{dx} = 2 \left(\frac{1}{4}x^2 + C\right) \times \frac{1}{2}x \\ &= x \left(\frac{1}{4}x^2 + C\right) \\ \text{RHS} &= xy^{\frac{1}{2}} = x \left(\left(\frac{1}{4}x^2 + C\right)^2\right)^{\frac{1}{2}} \\ &= x \left(\frac{1}{4}x^2 + C\right) \\ \text{LHS} &= \text{RHS} \end{aligned}$$

But there is another solution: namely $y(x) = 0$ for all x . This is called the “trivial solution”.

1.2 Initial Value Problems (IVP)

1st order IVP is a 1st order D.E. together with one extra condition:

$$\frac{dy}{dx} = f(x, y), y(x_0) = y_0$$

2nd order IVP

$$y'' = f(x, y, y')$$

Initial conditions:

- $y(x_0) = y_0$
- $y'(x_0) = y_1$

1.2.1 – Example

$$y' = y \text{ and } y(0) = 3$$

$y = ce^x$ is a one-parameter family of solutions

$$\frac{d}{dx}(ce^x) = ce^x = y$$

$$ce^1 = -2$$

$$c = -\frac{2}{e}$$

$$y = \left(-\frac{2}{e}\right) e^x$$

$$y = -2e^{x-1}$$

1.2.2 – Example

$$\text{D.E.: } y' + 2xy^2 = 0 \text{ and } y(0) = 1$$

Given that you have the solution: $y = \frac{1}{x^2+C}$, Solve:

$$-1 = \frac{1}{(0)^2 + c}$$

$$-1 = \frac{1}{c}$$

$$-1 \times c = 1$$

$$c = -1$$

$$y = \frac{1}{x^2 - 1}, I = (-1, 1)$$

1.2.3 – Example

$$\text{D.E.: } y' + 2xy^2 = 0 \text{ and } y(0) = 1$$

Example

$$x'' + 16x = 0 \text{ and } x\left(\frac{\pi}{2}\right) = 5 \text{ and } x'\left(\frac{\pi}{2}\right) = -4$$

$$x = c_1 \cos(4t) + c_2 \sin(4t)$$

$$5 = c_1 \cos(4t) + c_2 \sin(4t)$$

$$= c_1 \cos(2\pi) + c_2 \sin(2\pi)$$

$$= c_1(1) + c_2(0)$$

$$= c_1$$

$$x' = -4c_1 \sin(4t) + 4c_2 \cos(4t)$$

$$-4 = -4c_1 \sin\left(4\left(\frac{\pi}{2}\right)\right) + 4c_2 \cos\left(4\left(\frac{\pi}{2}\right)\right)$$

$$= -4c_1 \sin(2\pi) + 4c_2 \cos(2\pi)$$

$$= -4c_1(0) + 4c_2(1)$$

$$= 4c_2$$

$$-1 = c_2$$

Reasonable Question: Given a 1st order IVP, can we say whether a solution *exists* or not and, if a solution exists, is it *unique*.

Theorem: Given $y' = f(x, y)$ and $y(x_0) = y_0$, if $f(x, y)$ and $\frac{\partial f}{\partial y}$ are both continuous on a rectangle R containing (x_0, y_0) in its interior, then there exists an interval $I = (x_0 - h, x_0 + h)$ where $h > 0$ such that there exists a unique solution to IVP on I .

1.2.4 – Example

$$y' = xy^{\frac{1}{2}} \text{ and } y(1) = 2$$

- $f(x, y) = xy^{\frac{1}{2}}$ is continuous everywhere its defined $y \geq 0$
- $\frac{\partial f}{\partial y} = x \frac{1}{2} y^{-\frac{1}{2}} = \frac{x}{2y}$ is continuous everywhere its defined $y > 0$

1.2.5 – Example

$$y' = xy^{\frac{1}{2}} \text{ and } y(0) = 0$$

- $f(x, y) = xy^{\frac{1}{2}}$ is continuous for all x and $y \geq 0$
- $\frac{\partial f}{\partial y} = \frac{x}{2y}$ is continuous for all x and $y > 0$.
- **Theorem does not give any conclusion.**

Chapter 2

First-Order Differential Equations

2.1 Solution Curves Without a Solution

Given a 1st order D.E. $y' = f(x, y)$, y' is the slope of the tangent line at any point (x_0, y_0) on a solution curve

2.1.1 – Example

$$y' = f(x, y) = x + y$$

- $f(0, 0) = 0$
- $f(1, 0) = 1$

2.1.2 – Slope/Direction Fields




Figure 2.1: The direction field for the previous example

If the function $f(x, y)$ in the D.E. $y' = f(x, y)$ is reasonably simple so that we can solve $f(x, y) = 0$, we can make a “phase portrait diagram”. We will also assume $f(x, y)$ only involves the y -variable.

2.1.3 – Example

$$y' = (y + 2)(y - 3)(y - 5)$$

$$f(x, y) = (y + 2)(y - 3)(y - 5)$$

An “equilibrium solution” is a solution where y is a constant. In this example: $y = 3$, $y = 5$, $y = -2$ are each constant functions.




Figure 2.2: The equilibrium solution for the previous example.

The area around $y = 5$ is an unstable equilibrium since the solutions diverge and go in separate directions away from $y = 5$. The area around $y = 3$ is a stable equilibrium because the slopes above and below it converge to $y = 3$. The area around $y = -2$ is semi-stable, since all the slopes around it will converge in one direction, but the point isn't always $y = -2$.

2.2 Separable Differential Equations

Separable D.E.s are DE's $\frac{dy}{dx} = f(x, y)$ where $f(x, y)$ can be factored as $f(x, y) = g(x)h(y)$.

$$\frac{dy}{dx} = (1 + y^2)x^3 \text{ is separable}$$

$$\frac{dy}{dx} = \sin(xy) \text{ is not separable}$$

$$\frac{dy}{dx} = x^3y \text{ is not separable}$$

$$\frac{5}{xy} \frac{dy}{dx} = (x^2 + y) e^y$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{xy(x^2 + y)e^y}{5} \\ &= \frac{x(x^2 + y)}{5} \times ye^y \end{aligned}$$

2.2.1 – Method of Solution

“Separate the variable” to get $\frac{1}{h(y)} dy = g(x) dx$ or $p(y) dy = g(x) dx$ where $p(y) = \frac{1}{h(y)}$. **Integrate both sides**

$$\int p(y) dy = \int g(x) dx \text{ and if possible, solve for } y$$

2.2.2 – Example

$$\begin{aligned} \frac{dy}{dx} &= (1 + y^2) x^3 \\ \int \frac{1}{1 + y^2} dy &= \int x^3 dx \\ \tan^{-1}(y) + C_1 &= \frac{x^4}{4} + C_2 \\ \tan^{-1}(y) &= \frac{x^4}{4} + C_2 - C_1 \\ \tan^{-1}(y) &= \frac{x^4}{4} + C \\ y &= \tan\left(\frac{x^4}{4} + C\right) \end{aligned}$$

2.2.3 – Example

Problem 12 from the textbook.

$$\begin{aligned} \sin(3x) dx + 2y \cos^3(3x) dy &= 0 \\ \int -2y dy &= \int \frac{\sin(3x)}{\cos^3(x)} dx \\ &= \int \tan(3x) \sec^2(3x) dx \\ &= \int u \frac{1}{3} du \text{ where } u = \tan(3x), \quad du = 3 \sec^2(3x) dx \\ -2 \int y dy &= \frac{1}{3} \int u du + C \\ -y^2 &= \frac{u^2}{6} + C \\ &= \frac{\tan^2(3x)}{6} + C \\ \frac{\tan^2(3x)}{6} + y^2 &= -C \\ \frac{\tan^2(3x)}{6} + y^2 &= C \end{aligned}$$

Problem 25 from the textbook.

$$x^2 \frac{dy}{dx} = y - xy, y(-1) = -1$$

$$x^2 \frac{dy}{dx} = y - xy$$

$$x^2 \frac{dy}{dx} = y(1 - x)$$

$$\frac{dy}{y} = \frac{(1 - x)}{x^2} dx$$

$$\int \frac{dy}{y} = \int \frac{(1 - x)}{x^2} dx$$

$$\int \frac{dy}{y} = \int \frac{1}{x^2} dx - \int \frac{x}{x^2} dx$$

$$\int \frac{dy}{y} = \int \frac{1}{x^2} dx - \int \frac{1}{x} dx$$

$$\ln |y| + C_1 = -\frac{1}{x} + C_2 - \ln |x| + C_3$$

$$\ln |y| = -\frac{1}{x} - \ln |x| + C$$

$$y = e^{-\frac{1}{x}} \times e^{-\ln |x|} \times e^C$$

$$y = e^{-\frac{1}{x}} \times e^{-\ln |x|} \times e^C$$

$$y = e^{-\frac{1}{x}} \times \frac{1}{|x|} \times e^C$$

$$y = \frac{1}{|x|} e^{C - \frac{1}{x}}$$

$$-1 = \frac{1}{|-1|} e^{C - \frac{1}{-1}}$$

$$-1 = \frac{1}{1} e^{C - (-1)}$$

$$-1 = e^{C+1}$$

2.3 First Order Linear Differential Equations

$$a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

$$\frac{dy}{dx} + \frac{a_0(x)}{a_1(x)}y = \frac{g(x)}{a_1(x)}$$

$$\left. \frac{dy}{dx} + P(x)y = f(x) \right\} \text{ Standard form of a 1st-order linear DE}$$

We will try to find a function $\mu(x)$ such that by multiplying the D.E. by an integrating factor (I.F.) $\mu(x)$:

$$\mu(x) \frac{dy}{dx} + \mu(x)P(x)y = \mu(x)f(x)$$

such that the LHS is an exact derivative, Observe:

$$\frac{d}{dx} (\mu(x)y) = \mu(x) \frac{dy}{dx} + \frac{d\mu}{dx} y$$

from which we see

$$\mu(x)P(x) = \frac{d\mu}{dx}$$

$$P(x)dx = \frac{d\mu}{\mu(x)}$$

$$\int P(x)dx = \int \frac{d\mu}{\mu}$$

$$\int P(x)dx = \ln \mu$$

$$\ln \mu = \int P(x)dx$$

$$\mu = e^{\int P(x)dx}$$

2.3.1 – Example

$$x \frac{dy}{dx} - 4y = x^6 e^x$$

Standard form: $\frac{dy}{dx} - \frac{4}{x}y = x^5 e^x$

$$P(x) = -\frac{4}{x}$$

$$\mu = e^{\int \frac{-4}{x} dx}$$

$$= e^{-4 \ln x}$$

$$= e^{\ln x^{-4}}$$

$$= x^{-4}$$

$$\text{I.F.} = \mu = x^{-4}$$

Now multiply the standard form of the given D.E. by x^{-4} .

$$x^{-4} \frac{dy}{dx} - x^{-4} \frac{4}{x} y = x^{-4} x^5 e^x$$

$$x^{-4} \frac{dy}{dx} - x^{-4} \frac{4}{x} y = x e^x$$

$$\int \frac{d}{dx} (x^{-4} y) = \int x e^x$$

$$x^{-4} y = \int x e^x$$

2.3.2 – Example

$$\begin{aligned}
 (x^2 - 9) \frac{dy}{dx} + xy &= 0 \\
 (x^2 - 9) \frac{dy}{dx} + xy &= 0 \\
 \frac{dy}{dx} + \frac{x}{x^2 - 9}y &= 0 \\
 P(x) &= \frac{x}{x^2 - 9} \\
 \int P(x)dx &= \int \frac{x}{x^2 - 9}dx \\
 \int P(x)dx &= \int \frac{1}{u - 9} \frac{du}{2} \\
 \int P(x)dx &= \frac{1}{2} \int \frac{1}{u - 9} du \\
 \int P(x)dx &= \frac{1}{2} \ln |u - 9| \\
 \int P(x)dx &= \frac{1}{2} \ln |x^2 - 9| \\
 \mu &= e^{\frac{1}{2} \ln |x^2 - 9|} \\
 \mu &= e^{\ln |(x^2 - 9)^{\frac{1}{2}}|} \\
 \mu &= (x^2 - 9)^{\frac{1}{2}} \\
 \mu &= \sqrt{x^2 - 9} \\
 \sqrt{x^2 - 9} \left(\frac{dy}{dx} + \frac{x}{x^2 - 9}y \right) &= \sqrt{x^2 - 9}(0) \\
 \sqrt{x^2 - 9} \frac{dy}{dx} + \frac{x}{\sqrt{x^2 - 9}}y &= 0 \\
 \int \frac{d}{dx} (y\sqrt{x^2 - 9}) &= \int 0 \\
 y\sqrt{x^2 - 9} &= C \\
 y &= \frac{C}{\sqrt{x^2 - 9}}
 \end{aligned}$$

2.4 Exact Equations

1st Order D.E. in differential form

$$M(x, y)dx + N(x, y)dy = 0$$

Given a function

$$z = f(x, y)$$

, the total differential, dz , is defined as

$$dz = \frac{\partial f}{\partial x}dx + \frac{\partial f}{\partial y}dy$$

2.4.1 – Method

See if we can find a function $f(x, y)$ such that

$$\frac{\partial f}{\partial x} = M, \frac{\partial f}{\partial y} = N$$

If we can do this, then the D.E. is equivalent to

$$df = 0 \Rightarrow f(x, y) = c$$

is an implicit solution of D.E. Assume that M and N have continuous 1st order partials (assuming f exists)

$$\left. \begin{array}{l} My = \frac{\partial}{\partial y} \frac{\partial f}{\partial x} dy = f_{xy} \\ Nx = \frac{\partial}{\partial x} \frac{\partial f}{\partial y} dy = f_{yx} \end{array} \right\} \text{Theorem tells us these are equal}$$

This provides a quick test to check if the D.E. is exact or not.

2.4.2 – Example

$$\begin{aligned} 2xydx + (x^2 - 1)dy &= 0 \\ M(x, y) = 2xy, N(x, y) &= x^2 - 1 \end{aligned}$$

To check if the D.E. is exact

$$M_y = 2x = N_x$$

We now know there exists a function $f(x, y)$ with

$$\begin{aligned} \frac{\partial f}{\partial x} &= M = 2xy \\ \frac{\partial f}{\partial y} &= N = x^2 - 1 \end{aligned}$$

$$\begin{aligned}
 f_M(x, y) &= \int \frac{\partial f}{\partial x} dx \\
 &= \int 2xy dx \\
 &= x^2 y + \phi(y) \\
 \frac{\partial f}{\partial y} (x^2 y + \phi(y)) &= x^2 - 1 \text{ required to equal } N \\
 x^2 + \phi'(y) &= x^2 - 1 \\
 \phi'(y) &= -1 \\
 \phi(y) &= \int -1 dy \\
 &= -y \\
 f(x, y) &= x^2 y - y \\
 d(f(x, y)) &= 0 \\
 f(x, y) &= c \\
 x^2 y - y &= c \text{ is an implicit solution of the D.E.}
 \end{aligned}$$

Note: the f_M format is just there to show which partial equation was integrated. It was made by me and, as far as I know, not standardly known.

2.4.3 – Example

$$\begin{aligned}
 (e^{2y} - y \cos(xy)) dx + (2xe^{2y} - x \cos(xy) + 2y) dy &= 0 \\
 M_y &= N_x \\
 \frac{\partial}{\partial y} (e^{2y} - y \cos(xy)) &= \frac{\partial}{\partial x} (2xe^{2y} - x \cos(xy) + 2y) \\
 2e^{2y} - [\cos(xy) - y \sin(xy) \times x] &= 2e^{2y} - (\cos(xy) - x \sin(xy) \times y) + 0 \\
 2e^{2y} - \cos(xy) + xy \sin(xy) &= 2e^{2y} - \cos(xy) + xy \sin(xy) \\
 \frac{\partial f}{\partial x} = M &= e^{2y} - y \cos(xy) \\
 \frac{\partial f}{\partial y} = N &= 2xe^{2y} - x \cos(xy) + 2y \\
 f_N(x, y) &= \int \frac{\partial f}{\partial y} dy \\
 &= \int (2xe^{2y} - x \cos(xy) + 2y) dy \\
 &= \frac{2xe^{2y}}{2} - \frac{x \sin(xy)}{x} + 2 \times \frac{y^2}{2} + \phi(x) \\
 &= xe^{2y} - \sin(xy) + y^2 + \phi(x)
 \end{aligned}$$

Take the ∂x of this and equate with M :

$$\begin{aligned} M &= \frac{\partial}{\partial x} (xe^{2y} - \sin(xy) + y^2 + \phi(x)) \\ e^{2y} - y \cos(xy) &= e^{2y} - y \cos(xy) + 0 + \phi'(x) \\ 0 &= \phi'(x) \\ \phi(x) &= c \end{aligned}$$

So $f(x, y) = c_2$ is the solution

$$xe^{2y} - \sin(xy) + y^2 = c$$

$$dx = \frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy$$

2.4.4 – What can you do if $M_y \neq N_x$

Sometimes you can multiply the DE by an integrating factor $\mu(x, y)$ to get an exact DE. If

$$\frac{M_y - N_x}{N}$$

is a function of only x , then

$$\mu = e^{\int \frac{M_y - N_x}{N} dx}$$

will be an I.F.

If

$$\frac{N_x - M_y}{M}$$

is a function of only y , then

$$\mu = e^{\int \frac{N_x - M_y}{M} dy}$$

will be an I.F.

2.4.5 – Example

$$xydx + (2x^2 + 3y^2 - 20) dy = 0$$

$$M_y = x$$

$$N_x = 4x$$

$$M_y \neq N_x \quad \frac{N_x - M_y}{M} = \frac{4x - x}{xy}$$

$$= \frac{3x}{xy}$$

$$= \frac{3}{y} \text{ is a function of just } y$$

So:

$$\begin{aligned}
 \mu &= e^{\int \frac{3}{y} dy} \\
 &= e^{3 \ln y} \\
 &= y^3 \\
 xy^4 dx + y^3 (2x^2 + 3y^2 - 20) dy &= 0(y^3) \\
 xy^4 dx + (2x^2 y^3 + 3y^5 - 20y^3) dy &= \\
 M_y &= N_x \\
 4xy^3 &= 4xy^3 \frac{\partial f}{\partial x}
 \end{aligned}$$

2.5 Substitution Methods

Taking a D.E. that's not:

- Separable
- 1st Order Linear
- Exact

and making a substitution to turn the new D.E. into one of these **Theorem:** Given a D.E.

$$M(x, y)dx + N(x, y)dy = 0$$

A function $f(x, y)$ is said to be homogenous of order α if $f(tx, ty) = t^\alpha f(x, y)$.

2.5.1 – Example

Given:

$$f(x, y) = x^3 + 5xy^2 - y^3$$

Then:

$$\begin{aligned}
 f(tx, ty) &= (tx)^3 + 5(tx)(ty)^2 - (ty)^3 \\
 &= t^3 x^3 + 5t^3 xy^2 - t^3 y^3 \\
 &= t^3 (x^3 + 5xy^2 - y^3) \\
 &= t^3 f(x, y)
 \end{aligned}$$

2.5.2 – Example

$$\begin{aligned}
 f(x, y) &= \frac{x + y}{x^2 + y^2} \\
 f(tx, ty) &= \frac{tx + ty}{(tx)^2 + (ty)^2} \\
 f(tx, ty) &= \frac{tx + ty}{x^2t^2 + y^2t^2} \\
 f(tx, ty) &= \frac{t}{t^2} \times \frac{x + y}{x^2 + y^2} \\
 f(tx, ty) &= \frac{t}{t^2} f(x, y) \\
 f(tx, ty) &= \frac{1}{t} f(x, y)
 \end{aligned}$$

$f(x, y) = \frac{x+y}{x^2+y^2}$ is homogenous of order $\alpha = -1$

2.5.3 – Substitution Rule

If $M(x, y)$ and $N(x, y)$ are homogenous, each of the same order, then $u = \frac{y}{x}$ i.e., $y = ux$ or $v = \frac{x}{y}$ (i.e. $x = vy$) will produce a separable D.E.

2.5.4 – Example

Solve the separable D.E. and then back-substitute

$$(x^2 + y^2)dx + (x^2 - xy)dy = 0$$

$$M(x, y) = x^2 + y^2 \quad N = x^2 - xy$$

$$M_y = 2y \quad N_x = 2x - y$$

$$M_y \neq N_x$$

$$M(tx, ty) = (tx)^2 + (ty)^2$$

$$= t^2x^2 + t^2y^2$$

$$= t^2(x^2 + y^2)$$

$$= t^2M(x, y) \quad M \text{ is homogeneous of order 2 and so is } N$$

$$u = \frac{y}{x}$$

$$y = ux$$

$$dy = udx + xdu$$

$$(x^2 + (ux)^2)dx + (x^2 - x(ux))(udx + xdu) = 0$$

$$(x^2 + u^2x^2)dx + (x^2 - ux^2)(udx + xdu) = 0$$

$$(1 + u^2)x^2dx + x^2(1 - u)(udx + xdu) = 0$$

$$(1 + u^2)x^2dx + x^2(udx + xdu - u^2dx - uxdu) = 0$$

$$x^2(1dx + u^2dx + udx + xdu - u^2dx - uxdu) = 0$$

$$x^2(1dx + u^2dx - u^2dx + udx + xdu - uxdu) = 0$$

$$x^2(1dx + udx + xdu - uxdu) = 0$$

$$x^2(1 + u)dx + x^3(1 - u)du = 0$$

$$\int \frac{1}{x} dx = \int -\frac{1 - u}{1 + u} du$$

$$= \int \frac{u - 1}{u + 1} du$$

$$= \int \frac{u + (1 - 2)}{u + 1} du$$

$$= \int \left(\frac{u + 1}{u + 1} - \frac{2}{u + 1} \right) du$$

$$= \int \left(1 - \frac{2}{u + 1} \right) du$$

$$\ln|x| = \int \left(1 - \frac{2}{u + 1} \right) du$$

$$= u - 2 \ln|u + 1| + C$$

$$\ln|x| = \frac{y}{x} - 2 \ln \left| \frac{y}{x} + 1 \right| + C$$

Theorem: An equation of the form

$$\frac{dy}{dx} + P(x)y = f(x)y^n$$

where $n \neq 0, 1$ is called a Bernoulli Equation. The substitution

$$u = y^{1-n}$$

will transform the D.E. into a 1st order linear.

2.5.5 – Example

$$x \frac{dy}{dx} + y = x^2 y^2$$
$$\frac{dy}{dx} + \frac{y}{x} = xy^2$$

is a **Bernoulli equation** with $n = 2$.

$$\begin{aligned}
 u &= y^{1-2} \\
 &= y^{-1} \\
 &= \frac{1}{y} \\
 \frac{du}{dx} &= \frac{du}{dy} \times \frac{dy}{dx} \\
 &= -1y^{-2} \frac{dy}{dx} \\
 &= -\frac{1}{y^2} \frac{dy}{dx} \\
 -y^{-2} \frac{dy}{dx} + -y^{-2} \times \frac{y}{x} &= -y^{-2} \times xy^2 \\
 -y^{-2} \frac{dy}{dx} + -\frac{1}{x} y^{-1} &= -x \\
 \frac{du}{dx} - \frac{1}{x} u &= -x \\
 \text{I.F.} = \mu &= e^{\int P(x) dx} \\
 &= e^{-\int \frac{1}{x} dx} \\
 &= e^{-\ln|x|} \\
 &= e^{\ln|x^{-1}|} \\
 &= x^{-1} \\
 \frac{1}{x} \frac{du}{dx} - \frac{1}{x^2} u &= -1 \\
 \frac{d}{dx} \left(\frac{1}{x} u \right) &= -1 \\
 \int \frac{d}{dx} \left(\frac{1}{x} u \right) &= \int -1 dx \\
 \frac{1}{x} u &= \int -1 dx \\
 \frac{1}{x} u &= -x + C \\
 \frac{1}{x} \times 1y &= -x + C \\
 \frac{1}{x(-x + C)} &= y \\
 y &= \frac{1}{Cx - x^2}
 \end{aligned}$$

Theorem: If the D.E. can be expressed as

$$\frac{dy}{dx} = f(Ax + by + C)$$

for particular numbers A , B , C , then let

$$u = Ax + By + C$$

to get a separable D.E.

2.5.6 – Example

$$\frac{dy}{dx} = (-2x + y)^2 - 7, y(0) = 0$$

$$u = -2x + y$$

$$\frac{du}{dx} = \frac{dy}{dx} \times \frac{du}{dy}$$

$$= -2 + \frac{dy}{dx}$$

$$\frac{du}{dx} + 2 = \frac{dy}{dx}$$

$$\frac{du}{dx} + 2 = u^2 - 7$$

$$\frac{du}{dx} = u^2 - 9$$

$$\frac{du}{u^2 - 9} = dx$$

$$\int \frac{du}{u^2 - 9} = \int dx$$

$$\int \frac{du}{(u+3)(u+9)} = x + C$$

$$\int \frac{du}{(u+3)(u+9)} = x + C$$

Chapter 3

Modeling using DE

3.1 Linear DE Modeling

3.1.1 – Standard Problems

- 1) Population Growth (or decline)
- 2) Radioactive Decay
- 3) Newton's Law of Cooling
- 4) Mixture Problems

3.1.2 – Population Model

Assume the rate of population change is proportional to the size of the population

$P(t)$ = population at time t

$$\frac{dP}{dt} = kP$$

$\frac{\frac{dP}{dt}}{P} = k$ is the relative growth rate of the population

$$\begin{aligned}
\frac{dP}{dt} &= kP \\
\frac{dP}{P} &= kdt \\
\int \frac{dP}{P} &= \int kdt \\
\ln|P| &= kt + C \\
|P| &= e^{kt+C} \\
|P| &= e^{kt}e^C \\
|P| &= Ae^{kt} \text{ where } A > 0 \\
P &= \pm Ae^{kt} \\
P &= Be^{kt} \text{ where } B \neq 0 \\
P &= De^{kt} \text{ where } D \text{ can be any real number}
\end{aligned}$$

The constant can become any number because 0 would be a valid rate of population change, it means that the population size isn't changing.

3.1.3 – Example

If, initially at 2 p.m., there are 1,000 bacteria on a petri dish and at 4 p.m., there are 2,000 bacteria. Assuming constant relative growth rate, how many bacteria are there at 5 p.m.?

$P(t)$ = population t hours after 2 p.m.

$$P(t) = Ae^{kt}$$

$$1000 = Ae^{(0)k}$$

$$1000 = Ae^0$$

$$1000 = A(1)$$

$$A = 1000$$

$$P(2) = 2000$$

$$P(2) = 1000e^{2k}$$

$$2000 = 1000e^{2k}$$

$$2 = e^{2k}$$

$$\ln(2) = 2k$$

$$k = \frac{\ln(2)}{2}$$

$$P(t) = 1000e^{\frac{\ln(2)}{2}t}$$

$$P(3) = 1000e^{\frac{\ln(2)}{2}(3)}$$

$$= 1000e^{1.5 \ln(2)}$$

$$= 1000e^{\ln(2^{1.5})}$$

$$= 1000(2^{1.5})$$

$$= 2000(\sqrt{2})$$

$$P(3) \approx 2828.427(\sqrt{2})$$

$$P(t) = 1000e^{\frac{t}{2} \ln(2)}$$

$$= 1000e^{\frac{t}{2} \ln(2)}$$

$$= 1000e^{\ln(2^{\frac{t}{2}})}$$

$$= 1000 \times 2^{\frac{t}{2}}$$

3.1.4 – Radioactive Decay

$$m(t) = m_0e^{kt} \text{ where } k < 0$$

The Half-Life is the amount of time it takes for half of the original amount to remain:

$$\frac{1}{2}A_0 = A_0e^{kt} \Rightarrow \frac{1}{2} = e^{kt}$$

3.1.5 – Mixture Problems

Setup

Initially, the container has 200 gallons of brine solution (salt-water) of concentration $\frac{10 \text{ lbs}}{200 \text{ gallons}} = 0.05 \frac{\text{lbs}}{\text{gallon}}$. A solution of $\frac{5 \text{ lbs}}{200 \text{ gallons}} 0.025 \frac{\text{lbs}}{\text{gallon}}$ is poured into the initial container at a rate of $\frac{4 \text{ gallons}}{\text{min}}$. How many pounds of salt are there in the container after 2 hours.

Let $y(t) = \# \text{ lbs of salt } t \text{ minutes after the process starts}$
 $\frac{dy}{dt} =$ The rate of change of $\# \text{ lbs of salt}$

$$\begin{aligned} \frac{dy}{dt} &= 0.025 \frac{\text{lbs}}{\text{gal}} \times 4 \frac{\text{gal}}{\text{min}} \left\{ \text{rate in} \right. \\ &\quad \left. - \frac{y(t) \text{ lbs}}{200 \text{ gal}} \times 4 \frac{\text{gal}}{\text{min}} \right\} \text{rate out} \\ &= (0.025) 4 \frac{\text{lbs}}{\text{min}} - \frac{4y(t)}{200} \frac{\text{lbs}}{\text{min}} \\ &= 0.1 \frac{\text{lbs}}{\text{min}} - \frac{y(t)}{50} \frac{\text{lbs}}{\text{min}} \\ &= 0.1 - \frac{y(t)}{50} \end{aligned}$$

$$\begin{aligned} \frac{dy}{dt} + \frac{1}{50}y &= 0.1 \\ \mu &= e^{\int P(t)dt} \\ &= e^{\int \frac{1}{50}dt} \\ &= e^{\frac{t}{50}} \end{aligned}$$

$$e^{\frac{t}{50}} \left(\frac{dy}{dt} \right) + e^{\frac{t}{50}} \left(\frac{1}{50}y \right) = e^{\frac{t}{50}} (0.1)$$

$$\frac{d}{dt} \left(e^{\frac{t}{50}} y \right) = e^{\frac{t}{50}} (0.1)$$

$$\int \frac{d}{dt} \left(e^{\frac{t}{50}} y \right) = \int \frac{1}{10} e^{\frac{t}{50}}$$

$$e^{\frac{t}{50}} y = \frac{1}{10} \times \frac{e^{\frac{t}{50}}}{\frac{1}{50}} + C$$

$$e^{\frac{t}{50}} y = 5e^{\frac{t}{50}} + C$$

$$\begin{aligned} y &= 5 + Ce^{-\frac{t}{50}} \\ &= 5 + Ce^{-0.02t} \end{aligned}$$

$$\begin{aligned} y(120) &= 5 + Ce^{-0.02(120)} \\ &= 5 + Ce^{-2.4} \end{aligned}$$

Chapter 4

Higher Order Differential Equations

An n th order DE is linear if it had the form

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + a_{n-2}(x) \frac{d^{n-2} y}{dx^{n-2}} + \cdots + a_1(x) \frac{dy}{dx} + a_0 y = g(x)$$

Theorem: If all the coefficient functions are continuous and $a_n(x)$ is not 0 on an interval I and $g(x)$ is continuous, then any initial value problem

$$DE + y(x_0) = y_0$$

has a unique solution on the interval I if $g(x) = 0$. i.e.

$$a_n(x)y^{(n)} + \cdots + a_0(x)y = 0$$

then the DE is said to be homogeneous.

4.0.1 – Example

$$y'' - 3y' - 4y = 0$$

Show $y_1 = e^{4x}$ is a solution and $y_2 = e^{-x}$ is a solution.

$$y_1 = e^{4x}$$

$$y_1' = 4e^{4x}$$

$$y_1'' = 16e^{4x}$$

$$16e^{4x} - 3(4e^{4x}) - 4e^{4x} = 0$$

$$16e^{4x} - 12e^{4x} - 4e^{4x} = 0$$

$$e^{4x}(16 - 12 - 4) = 0$$

$$e^{4x}(0) = 0$$

$$0 = 0$$

$$y_3 = 6y_1 = 6e^{4x}$$

$$y'_3 = 6y'_1 = 24e^{4x}$$

$$y''_3 = 6y''_1 = 96e^{4x}$$

$$96e^{4x} - 3(24e^{4x}) - 4(6e^{4x}) = 0$$

$$96e^{4x} - 72e^{4x} - 24e^{4x} = 0$$

$$e^{4x}(96 - 72 - 24) = 0$$

$$e^{4x}(0) = 0$$

$$0 = 0$$

Theorem: Superposition Principle: if y_1, y_2, \dots, y_m are each solutions of an n th order Linear, homogenous DE, then $c_1y_1 + c_2y_2 + \dots + c_my_m$ will also be a solution for any constants c_1, c_2, \dots, c_m .

Our goal is to express the general solution in as concise a way as possible.

Linear combination – a collection of solutions y_1, y_2, \dots, y_m is linearly independent is if the only way $c_1y_1 + c_2y_2 + \dots + c_my_m = 0$ is iff (if and only if) all of the constants $c_1, c_2, \dots, c_m = 0$. Otherwise we say y_1, y_2, \dots, y_m are linearly dependent. **Theorem:** If the DE is an n th order Linear Homogeneous equation then there will exist a collection of n linearly independent solutions y_1, y_2, \dots, y_n and the general solution will be $y_c = c_1y_1 + c_2y_2 + \dots + c_ny_n$. One way to check for linear independence is to compile the Wronskian

$$W(y_1, y_2, \dots, y_n) = \det \begin{bmatrix} y_1 & y_2 & \cdots & y_n \\ y'_1 & y'_2 & \cdots & y'_n \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)} & y_2^{(n-1)} & \cdots & y_n^{(n-1)} \end{bmatrix}$$