Contents

0.1		lose Model
0.2		odel
0.3	Example $2 \dots$	
0.4	Lead Uptake Model .	
Exam	ple 1 Multiple Dose I	Model
Two C	Compartment Model	
		$input \to x(t) \to^{k_1} y(t) \to^{k_2}$
System	n of ODE's	
		$\frac{dx}{dt} = -k_1 x + I(t), x(0) = 0$
		$\frac{dy}{dt} = k_1 x - k_2 y, y(0) = 0$
0.2.1	Methods of Sol	ution for Systems of Equations
(1) E	Elimination	
(2) L	aplace Transforms	
(3) N	Matrix methods	
	• eigenvalue problem	
(4) N	Numerical methods	

 $\bullet\,$ Mathematica, Matlab and Python

Example 2

$$\frac{dx}{dt} = x_1 + 3x_2, \quad x_1(0) = 5$$

$$\frac{dy}{dt} = 5x_1 + 3x_2, \quad x_2(0) = 1$$

$$sX_1(s) - x_1(0) = \mathcal{L}\{x_1(t)\} \qquad sX_2(s) - x_2(0) = \mathcal{L}\{x_2(t)\}$$

$$sX_1(s) - x_1(0) = \mathcal{L}\{x_1\} + \mathcal{L}\{3x_2\} \qquad sX_2(s) - x_2(0) = \mathcal{L}\{5x_1\} + \mathcal{L}\{3x_2\}$$

$$sX_1(s) - 5 = X_1(s) + 3X_2(s) \qquad sX_2(s) - 1 = 5X_1(s) + 3X_2(s)$$

$$sX_1(s) - 5 - X_1(s) = 3X_2(s) \qquad sX_2(s) - 1 - 5X_1(s) = 3X_2(s)$$

$$sX_1(s) - 5 - X_1(s) = sX_2(s) - 1 - 5X_1(s) = 3X_2(s)$$

$$sX_1(s) + 4X_1(s) - sX_2(s) = 4$$

$$(s + 4)X_1(s) - sX_2(s) = 4$$

$$(s + 4)X_1(s) - sX_2(s) = 4$$

$$\frac{dx}{dt} = -k_1x + D\delta(t), \quad x(0) = 0$$

$$\frac{dy}{dt} = k_1x - k_2y, \quad y(0) = 0$$

$$(1)$$

Use Laplace Transforms

$$X(s) = \mathcal{L}\left\{x(t)\right\}$$
 $Y(s) = \mathcal{L}\left\{y(t)\right\}$

 $\mathscr{L}\left\{ \left(\mathbf{1}\right) \right\} :$

$$sX(s) - x(0) = \mathcal{L}\left\{-k_1x\right\} + \mathcal{L}\left\{D\delta(t)\right\}$$

$$sX(s) - 0 = -k_1X(s) + D(1)$$

$$sX(s) + k_1X(s) = D$$

$$(s + k_1)X(s) = D$$

$$X(s) = \frac{D}{s + k_1}$$

$$X(s) = \frac{D}{s + k_1}$$

$$X(s) = \frac{D}{s + k_1}$$

$$Y(s) = \frac{Dk_1}{(s + k_2)(s + k_1)}$$

$$s = -k_{1} s = -k_{2}$$

$$Dk_{1} = A(-k_{1} + k_{1}) + B(-k_{1} + k_{2}) Dk_{1} = A(-k_{2} + k_{1}) + B(-k_{2} + k_{2})$$

$$Dk_{1} = A(0) + B(k_{2} - k_{1}) Dk_{1} = A(k_{1} - k_{2}) + B(0)$$

$$Dk_{1} = B(k_{2} - k_{1}) Dk_{1} = A(k_{1} - k_{2})$$

$$B = \frac{Dk_{1}}{k_{2} - k_{1}} A = \frac{Dk_{1}}{k_{1} - k_{2}}$$

$$X(s) = \frac{D}{s + k_{1}} Y(s) = \frac{Dk_{1}}{k_{2} - k_{1}} + \frac{Dk_{1}}{k_{1} - k_{2}}$$

$$x(t) = \mathcal{L}^{-1} \left\{ \frac{D}{s + k_{1}} \right\} y(t) = \mathcal{L}^{-1} \left\{ \frac{Dk_{1}}{k_{2} - k_{1}} \right\} + \mathcal{L}^{-1} \left\{ \frac{Dk_{1}}{k_{1} - k_{2}} \right\}$$

Lead Uptake Model

- Excess of lead in body leads to lead (Pb) poisoning.
- Symptoms of Pb poisoning

Adults GI tract – vomiting pain

Children Serious – central nervous system disorders, anemia

• Pb in the environment:

Natural small

Human-made

- gasoline
- paint
- waterpipes
- make-up
- plastics
- How does Pb enter the body?
 - Inhalation
 - Eat or drinking
- How is Pb removed from the body?
 - urine
 - hair, nails, sweat