

Example 1 Pendulum

$\phi(t)$ = angular displacement.

$\phi(0) = \theta$ = initial angular displacement.

Let t_ϕ = period of the pendulum (time for 1 complete cycle)

What factors determine t_ϕ ($[t_\phi = T]$)?

1. g

$$\bullet [g] = LT^{-2}$$

2. m – mass

$$\bullet [m] = M$$

3. r – length

$$\bullet [r] = L$$

4. θ = initial displacement

$$\bullet [\theta] = 1$$

5. air resistance (drag)

6. friction in hinge

Assume $t_\phi = f(m, g, r, \theta)$ Dimension reduction:

$$\begin{aligned} [t_\phi] &= [m^a g^b r^c \theta^d] \\ [t_\phi] &= [m]^a [g]^b [r]^c [\theta]^d \\ M^0 L^0 T^1 &= M^a (LT^2)^b L^c (M^0 L^0 T^0)^d \\ &= M^a L^b T^{2b} L^c M^{0d} L^{0d} T^{0d} \\ &= M^a L^b T^{2b} L^c (1) \\ &= M^a L^{b+c} T^{2b} \end{aligned}$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & -2 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\begin{aligned}
\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & -2 & 0 & 0 & 1 \end{array} \right] &= (r_2 \leftarrow 2r_2 + r_3) \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & -2 & 0 & 0 & 1 \end{array} \right] \\
&= (r_2 \leftrightarrow r_3) \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 & 1 \\ 0 & 0 & 2 & 0 & 0 \end{array} \right] \\
&= \left(r_2 \leftarrow -\frac{r_2}{2} \right) \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -\frac{1}{2} \\ 0 & 0 & 2 & 0 & 0 \end{array} \right] \\
&= \left(r_3 \leftarrow \frac{r_3}{2} \right) \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -\frac{1}{2} \\ 0 & 0 & 1 & 0 & 0 \end{array} \right] \\
a = 0 \quad b = -\frac{1}{2} \quad c = 0 \quad d = k \\
t_\phi = m^0 g^{-\frac{1}{2}} r^c \theta^d
\end{aligned}$$

Buckingham's Pi Theorem

Let q = physical quantity (measurable). q depends on n parameters p_1, \dots, p_n

$$q = f(p_1, p_2, \dots, p_n)$$

Assume fundamental dimensions h, T, M .

$$[q] = L^{l_0} T^{t_0} M^{m_0}$$

$$[p_i] = L^{l_i} T^{t_i} M^{m_i}$$

$$[q] = [p_1]^{a_1} [p_2]^{a_2} \dots [p_n]^{a_n}$$

Objective: find a_j where $j = 1, 2, \dots, n$

$$L^{l_0} T^{t_0} M^{m_0} = (L^{l_1} T^{t_1} M^{m_1})^{a_1} \times (L^{l_2} T^{t_2} M^{m_2})^{a_2} \times \dots \times (L^{l_n} T^{t_n} M^{m_n})^{a_n}$$

$$L : l_0 =$$

$A\vec{x} = \vec{b}$ has a solution of form

$$\vec{x} = \vec{x}_\phi + \gamma_1 \vec{x}_1 + \gamma_2 \vec{x}_2 + \dots + \gamma_n \vec{x}_n$$

where γ is arbitrary, \vec{x}_ϕ is the particular solution, and $\{\vec{x}_j\}$ is the null space of A .

Example 2

Drag on a sphere within a fluid. The sphere has radius R moving at velocity v with a resistant drag force D_F . What factors play a role?

R = radius $[L]$

v = velocity $[LT^{-1}]$

ρ = density $[ML^{-3}]$

μ = dynamic viscosity $[ML^{-1}T^{-1}]$

$$D_F = f(R, v, \rho, \mu)$$

$$[D_F] = [R]^a [v]^b [\rho]^c [\mu]^d$$

$$MLT^{-2} = L^a (LT^{-1})^b (ML^{-3})^c (ML^{-1}T^{-1})^d$$

$$MLT^{-2} = L^a \times L^b T^{-b} \times M^c L^{-3c} \times M^d L^{-d} T^{-d}$$

$$MLT^{-2} = M^{c+d} L^{a+b-3c-d} T^{-b-d}$$

$$\begin{bmatrix} 1 & 1 & -3 & -1 \\ 0 & -1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

$$\begin{aligned} \left[\begin{array}{cccc|c} 1 & 1 & -3 & -1 & 1 \\ 0 & -1 & 0 & -1 & -2 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right] &= (r_2 \leftarrow -r_2) \left[\begin{array}{cccc|c} 1 & 1 & -3 & -1 & 1 \\ 0 & 1 & 0 & 1 & 2 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right] \\ &= (r_1 \leftarrow r_1 - r_2) \left[\begin{array}{cccc|c} 1 & 0 & -3 & -2 & 0 \\ 0 & 1 & 0 & 1 & 2 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right] \\ &= (r_1 \leftarrow r_1 + 3r_3) \left[\begin{array}{cccc|c} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 2 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right] \end{aligned}$$

$$\begin{aligned} a + d &= 0 & b + d &= 2 & c + d &= 1 \\ a &= -d & b &= -d + 2 & c &= -d + 1 \end{aligned}$$

Example 3 T-Rex Top Speed

How fast can a Tyrannosaurus Rex (T-Rex) walk or run?

Develop a model!

Find v = velocity $[v] = \left[\frac{L}{T}\right]$

What variables are important?

m mass, $[M]$

g gravity, $[LT^{-2}]$

h hip length, $[L]$

s stride length – distance between 2 steps, $[s] = L$

$$v = f(s, h, g, m)$$

$$[v] = [s^a h^b g^c m^d]$$

$$LT^{-1} = L^a L^b (LT^{-2})^c M^d$$

$$LT^{-1} = L^{a+b} L^c T^{-2c} M^d$$

$$LT^{-1} = L^{a+b+c} T^{-2c} M^d$$

Equate exponents:

$$L : \quad 1 = a + b + c$$

$$T : \quad -1 = -2c$$

$$M : \quad 0 = d$$