
bassinat

Lindsey Weyant

12/2/2021

```
inat%>%
  ggplot(aes(x = taxon_kingdom_name)) +
  geom_bar() +
  labs(x = "Kingdom Name", y = "Count", title = "Kingdoms of the ECWA Watershed")
```

Kingdoms of the ECWA Watershed




```
inat%>%
count(scientific_name, sort = TRUE)
```

```
## 5 Cardinalis cardinalis
                                 75
## 6 Phytolacca americana
                                 63
## 7 Liriodendron tulipifera
                                 62
## 8 Hedera helix
                                 61
## 9 Ligustrum sinense
                                 61
## 10 Turdus migratorius
                                 58
## # ... with 1,642 more rows
inatfungi <- inat %>%
  filter(taxon_kingdom_name == "Fungi")
inatfungi %>%
  count(scientific_name)
## # A tibble: 68 x 2
     scientific_name
                                    n
##
      <chr>
                                <int>
## 1 Amanita flavoconia
## 2 Amanita flavorubens
## 3 Amanita parcivolvata
                                    1
## 4 Amanita persicina
                                    1
## 5 Amanita rubescens
                                    1
## 6 Apioperdon pyriforme
## 7 Aureoboletus betula
                                    3
## 8 Auricularia americana
                                    1
## 9 Baeospora myosura
                                    1
## 10 Cantharellus cinnabarinus
## # ... with 58 more rows
inatprotozoa <- inat %>%
 filter(taxon_kingdom_name == "Protozoa")
inatprotozoa %>%
  count(scientific_name)
## # A tibble: 3 x 2
##
     scientific_name
                             n
     <chr>>
                         <int>
## 1 Diachea leucopodia
                             1
## 2 Fuligo septica
## 3 Lycogala epidendrum
inatanimalia <- inat %>%
  filter(taxon_kingdom_name == "Animalia")
inatanimalia %>%
    count(scientific_name, sort=TRUE)
## # A tibble: 974 x 2
##
      scientific_name
                                   n
##
      <chr>>
                               <int>
## 1 Cardinalis cardinalis
                                  75
                                  58
## 2 Turdus migratorius
## 3 Buteo lineatus
                                  56
## 4 Thryothorus ludovicianus
                                  50
## 5 Papilio polyxenes
                                  45
## 6 Danaus plexippus
                                  43
```

```
## 7 Sciurus carolinensis
                                  43
## 8 Storeria dekayi
                                  43
## 9 Xylocopa virginica
                                  43
## 10 Argiope aurantia
                                  39
## # ... with 964 more rows
inatplantae <- inat %>%
  filter(taxon_kingdom_name == "Plantae")
inatplantae %>%
  count(scientific_name)
## # A tibble: 607 x 2
##
      scientific_name
                                 n
##
      <chr>
                             <int>
## 1 Abelmoschus esculentus
                                 1
## 2 Acalypha hispida
## 3 Acalypha rhomboidea
                                 1
## 4 Acer buergerianum
                                 1
## 5 Acer floridanum
                                 9
## 6 Acer negundo
                                24
## 7 Acer palmatum
                                 2
## 8 Acer rubrum
                                12
## 9 Acer saccharinum
                                 2
## 10 Acer saccharum
                                 1
## # ... with 597 more rows
2\ \mathrm{most} common kingdoms: animalia and plantae
inatanimalia %>%
  ggplot(aes(x=taxon_class_name)) +
  geom_bar() +
  scale_x_discrete(guide = guide_axis(n.dodge=3)) +
  labs(x= "Class Name", y = "Count", title = "Animalia Observations by Class")
```

Animalia Observations by Class




```
inatanimalia %>%
 filter(taxon_class_name == "Insecta") %>%
count(scientific_name, sort = TRUE)
## # A tibble: 719 x 2
##
      scientific_name
                             n
##
      <chr>
                         <int>
##
  1 Papilio polyxenes
## 2 Danaus plexippus
                            43
## 3 Xylocopa virginica
## 4 Harmonia axyridis
                            39
## 5 Bombus impatiens
                            34
## 6 Papilio glaucus
                            34
## 7 Plathemis lydia
                            33
## 8 Junonia coenia
                            28
  9 Phyciodes tharos
                            28
## 10 Alaus oculatus
                            26
## # ... with 709 more rows
inatanimalia %>%
  filter(taxon_class_name == "Aves") %>%
count(scientific_name, sort = TRUE)
## # A tibble: 120 x 2
##
      scientific_name
                                   n
##
      <chr>
                               <int>
```

75

1 Cardinalis cardinalis

```
## 2 Turdus migratorius
                                  58
## 3 Buteo lineatus
                                  56
## 4 Thryothorus ludovicianus
                                  50
## 5 Setophaga coronata
                                  39
## 6 Spinus tristis
                                  39
## 7 Ardea herodias
                                  36
## 8 Sayornis phoebe
                                  35
## 9 Dryobates pubescens
                                  32
## 10 Mimus polyglottos
                                  30
## # ... with 110 more rows
inatanimalia %>%
  filter(taxon_class_name == "Amphibia") %>%
  count(scientific_name, sort = TRUE)
## # A tibble: 14 x 2
##
     scientific name
##
      <chr>>
                                <int>
## 1 Lithobates palustris
                                   23
## 2 Hyla chrysoscelis
                                   20
## 3 Acris crepitans
                                   17
## 4 Hyla cinerea
                                   16
## 5 Lithobates catesbeianus
                                   13
## 6 Lithobates clamitans
                                    9
## 7 Anaxyrus americanus
                                    7
## 8 Ambystoma opacum
                                    4
                                    3
## 9 Eurycea cirrigera
                                    2
## 10 Anaxyrus fowleri
## 11 Lithobates sphenocephalus
                                    2
## 12 Notophthalmus viridescens
## 13 Pseudacris crucifer
## 14 Pseudacris feriarum
                                    1
inatanimalia %>%
  filter(taxon_class_name == "Reptilia") %>%
count(scientific_name, sort = TRUE)
## # A tibble: 29 x 2
##
      scientific name
##
      <chr>
                                  <int>
## 1 Storeria dekayi
## 2 Anolis carolinensis
                                     25
## 3 Haldea striatula
                                     22
## 4 Pantherophis alleghaniensis
                                     18
## 5 Agkistrodon contortrix
                                     14
## 6 Scincella lateralis
                                    13
## 7 Trachemys scripta
                                    13
## 8 Chelydra serpentina
                                    8
                                     7
## 9 Thamnophis sirtalis
## 10 Carphophis amoenus
## # ... with 19 more rows
inatplantae %>%
  ggplot(aes(x=taxon_class_name)) +
  geom_bar() +
  scale_x_discrete(guide = guide_axis(n.dodge=3)) +
```

Plantae Observations by Class



```
inatplantae %>%
  filter(taxon_class_name == "Magnoliopsida") %>%
count(scientific_name, sort = TRUE)
```

```
## # A tibble: 489 x 2
      scientific_name
##
                                  n
##
      <chr>
                              <int>
## 1 Alliaria petiolata
                                111
## 2 Ficaria verna
                                107
## 3 Impatiens capensis
                                 91
## 4 Phytolacca americana
                                 63
## 5 Liriodendron tulipifera
## 6 Hedera helix
                                 61
## 7 Ligustrum sinense
                                 61
## 8 Toxicodendron radicans
                                 54
## 9 Liquidambar styraciflua
                                 52
## 10 Sassafras albidum
                                 51
## # ... with 479 more rows
```

```
inatplantae %>%
  filter(taxon_class_name == "Liliopsida") %>%
count(scientific_name, sort = TRUE)
```

```
## # A tibble: 95 x 2
## scientific_name r
```

##		<chr></chr>	<int></int>
##	1	Microstegium vimineum	139
##	2	Arisaema triphyllum	28
##	3	Smilax rotundifolia	23
##	4	Commelina communis	18
##	5	Allium vineale	16
##	6	Dioscorea polystachya	16
##	7	Chasmanthium latifolium	14
##	8	Tipularia discolor	11
##	9	Lycoris radiata	10
##	10	Iris pseudacorus	9
##	#	with 85 more rows	