

**Serum Phospholipid (PL) Fatty Acid Composition
Predicts Declines in Insulin Sensitivity (IS) and Beta-Cell
Function Over 6-years in the Prospective Metabolism
and Islet Cell Evaluation (PROMISE) Cohort**

Luke W. Johnston, MSc, PhD (c)

Disclosures

Disclosures for all co-authors for consulting, advisory boards, and research support from:

- Merck, Novo Nordisk, Sanofi, Takeda, AztraZeneca, BI/Lilly, Janssen, Abbott, Health Canada, Lawson Foundation, Canadian Institutes for Health Research, Canadian Diabetes Association, Banting and Best Diabetes Association

No conflicts of interest pertaining to the study being presented.

Fatty acids as risk factor for diabetes and it's pathogenesis

¹Giacca et al. (2011); Xiao, Giacca, and Lewis (2009)

Fatty acids as risk factor for diabetes and it's pathogenesis

- Fatty acids either gotten from diet or *de novo*
- Harm or benefit to metabolism depending on chain length and unsaturation
- Example: higher palmitic acid (16:0) experimentally shown to be lipotoxic to beta-cells *in vivo* and *in vitro*¹

¹Giacca et al. (2011); Xiao, Giacca, and Lewis (2009)

²Yang, Kang, and Guan (2013); Newsholme et al. (2007); Giacca et al. (2011); Gehrman, Elsner, and Lenzen (2010); Ebbert and Jensen (2013)

Fatty acids as risk factor for diabetes and it's pathogenesis

- Fatty acids either gotten from diet or *de novo*
- Harm or benefit to metabolism depending on chain length and unsaturation
- Example: higher palmitic acid (16:0) experimentally shown to be lipotoxic to beta-cells *in vivo* and *in vitro*¹

Potential mechanisms from fatty acid composition²

- Fluidity of cell membrane (lipid bilayer)
- Inflammation (eg. via eicosanoids)
- Lipotoxicity (by-product accumulation)

¹Giacca et al. (2011); Xiao, Giacca, and Lewis (2009)

²Yang, Kang, and Guan (2013); Newsholme et al. (2007); Giacca et al. (2011); Gehrmann, Elsner, and Lenzen (2010); Ebbert and Jensen (2013)

Key findings from large cohorts on phospholipid (PL) fatty acid composition and incident diabetes:

- CHS (USA)³:
 - 16:0 and 18:0 higher risk for DM
 - 18:1n-7 lower risk for DM
 - 18:3n-3 slight lower risk for DM
- EPIC (Europe)⁴:
 - 16:0 higher risk for DM
 - 18:3n-6 higher risk for DM
- ARIC (USA)⁵:
 - 18:1n-9 slight lower risk for DM
 - 18:0 slight higher risk for DM

³Ma et al. (2015); Djoussé et al. (2011)

⁴Forouhi et al. (2014); Kröger et al. (2011)

⁵L. Wang et al. (2003)

Limitations: Few longitudinal designs or beta-cell function measures

- Few longitudinal studies
- Few studies on pathophysiology
 - No longitudinal data on beta-cell function

Limitations: Few longitudinal designs or beta-cell function measures

- Few longitudinal studies
- Few studies on pathophysiology
 - No longitudinal data on beta-cell function

Objective

To determine whether individual PL fatty acids associate longitudinally with insulin sensitivity (IS) and beta-cell function over a 6 year period.

Methods: Prospective Metabolism and Islet cell Evaluation (PROMISE) Cohort

Longitudinal observational cohort

- At-risk for diabetes:
 - Central obesity
 - Hypertension
 - Family history
- London and Toronto, Canada
- Visits every 3-yrs; 6-yrs of follow-up (3 time points)
- Follow-up rate of 79.6% over 6-yrs
- OGTT (3 samples) at each visit
- 22 PL fatty acids at baseline visit (**n=477**)



Variables of interest

Outcomes

Outcome	Measure	Marker of:
Insulin sensitivity ⁶	HOMA-IS ⁷ ISI	Hepatic IS Whole body IS
Beta-cell function ⁸	IGI/IR ISSI-2	1st phase response Disposition index

⁶Matthews, Hosker, and Rudenski (1985); Matsuda and DeFronzo (1999)

⁷Inverted HOMA-IR

⁸N. Wareham et al. (1995); Retnakaran et al. (2009)

Variables of interest

Outcomes

Outcome	Measure	Marker of:
Insulin sensitivity ⁶	HOMA-IS ⁷ ISI	Hepatic IS Whole body IS
Beta-cell function ⁸	IGI/IR ISSI-2	1st phase response Disposition index

Over the 6-yrs, median declines of **8.7% to 19.5%**.

⁶Matthews, Hosker, and Rudenski (1985); Matsuda and DeFronzo (1999)

⁷Inverted HOMA-IR

⁸N. Wareham et al. (1995); Retnakaran et al. (2009)

Variables of interest

Outcomes

Outcome	Measure	Marker of:
Insulin sensitivity ⁶	HOMA-IS ⁷ ISI	Hepatic IS Whole body IS
Beta-cell function ⁸	IGI/IR ISSI-2	1st phase response Disposition index

Over the 6-yrs, median declines of **8.7% to 19.5%**.

Predictors

22 phospholipid (PL) fatty acids quantified using thin-layer chromatography and gas chromatography, using an internal standard.

⁶Matthews, Hosker, and Rudenski (1985); Matsuda and DeFronzo (1999)

⁷Inverted HOMA-IR

⁸N. Wareham et al. (1995); Retnakaran et al. (2009)

Statistical analysis techniques

Generalized estimating equations (GEE)

- Robust longitudinal technique
- Fatty acids: *Time-independent* (only at baseline)
- Outcomes: *Time-dependent*
- P-values adjusted using False Discovery Rate (multiple testing)

Statistical analysis techniques


Generalized estimating equations (GEE)

- Robust longitudinal technique
- Fatty acids: *Time-independent* (only at baseline)
- Outcomes: *Time-dependent*
- P-values adjusted using False Discovery Rate (multiple testing)


Covariate selection

- Literature + causal directed acyclic graphs + information criteria:
 - Final model: Visit number, sex, ethnicity, baseline age, waist circumference, total free fatty acids, ALT, family history of diabetes


Adjusted GEE results: Several PL fatty acids (mol%) associate with IS and beta-cell function



PL 16:0 and 20:2n-6 (mol%) have significant interactions



Lower PL 20:2n-6 and higher 16:0 associate with steeper declines in beta-cell function



Previous literature is fairly consistent with PL results⁹

- 16:0, 18:0 have consistent harmful associations
- 18:1n-9, 18:1n-7 have consistent positive associations
- Inconsistent for 20:2n-6
- *However, no previous report has been on longitudinal outcomes nor interactions over time*

⁹Forouhi et al. (2014); Kröger et al. (2011); Ma et al. (2015); Mahendran et al. (2014); L. Wang et al. (2003)

Acknowledgements

- **Supervisor:** Dr. Anthony Hanley
- **Co-Supervisor:** Dr. Richard Bazinet
- **Committee Member:** Dr. Adria Giacca
- **Hanley Lab:** Ingrid Santaren, Robin Glicksman, Zhila Semnani-Azad, Windy Wang
- **Bazinet Lab:** Katie Hopperton, Marco Trepanier, Alex Kitson, Lin Lin, Zhen Liu, Shoug Alashmali
- **Research Nurses:** Jan Neuman, Paula Van Nostrand, Stella Kink, Annette Barnie, Sheila Porter, Mauricio Marin
- **Funding:** CDA, CIHR, BBDC



References

- Djoussé, Luc, Mary L Biggs, Rozenn N Lemaitre, Irena B King, Xiaoling Song, Joachim H Ix, Kenneth J Mukamal, David S Siscovick, and Dariush Mozaffarian. 2011. "Plasma Omega-3 Fatty Acids and Incident Diabetes in Older Adults." *Am J Clin Nutr* 94 (2): 527–33. doi:10.3945/ajcn.111.013334.
- Ebbert, Jon O., and Michael D. Jensen. 2013. "Fat Depots, Free Fatty Acids, and Dyslipidemia." *Nutrients* 5 (2): 498–508. doi:10.3390/nu5020498.
- Forouhi, Nita G., Albert Koulman, Stephen J. Sharp, Fumiaki Imamura, Janine Kröger, Matthias B. Schulze, Francesca L. Crowe, et al. 2014. "Differences in the Prospective Association Between Individual Plasma Phospholipid Saturated Fatty Acids and Incident Type 2 Diabetes: The EPIC-InterAct Case-Cohort Study." *Lancet Diabetes Endocrinol* 2 (10): 810–18. doi:10.1016/S2213-8587(14)70146-9.
- Gehrmann, W. M. Elsner, and S. Lenzen. 2010. "Role of Metabolically Generated Reactive Oxygen Species for Lipotoxicity in Pancreatic Beta-Cells." *Diabetes Obes Metab* 12 Suppl 2: 149–58. doi:10.1111/j.1463-1326.2010.01265.x.
- Giacca, Adria, Changting Xiao, Andrei I. Oprescu, Andre C. Carpentier, and Gary F. Lewis. 2011. "Lipid-Induced Pancreatic Beta-Cell Dysfunction: Focus on in Vivo Studies." *Am J Physiol Endocrinol Metab* 300 (2): E255–62. doi:10.1152/ajpendo.00416.2010.
- Kröger, Janine, Vera Zietemann, Cornelia Enzenbach, Cornelia Weikert, Eugène Hjm Jansen, Frank Döring, Hans-Georg Joost, Heiner Boeing, and Matthias B. Schulze. 2011. "Erythrocyte Membrane Phospholipid Fatty Acids, Desaturase Activity, and Dietary Fatty Acids in Relation to Risk of Type 2 Diabetes in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study." *Am J Clin Nutr* 93 (1): 127–42. doi:10.3945/ajcn.110.005447.
- Ma, Wenjie, Jason H. Y. Wu, Qianyi Wang, Rozenn N. Lemaitre, Kenneth J. Mukamal, Luc Djoussé, Irena B. King, et al. 2015. "Prospective Association of Fatty Acids in the de Novo Lipogenesis Pathway with Risk of Type 2 Diabetes: The Cardiovascular Health Study." *Am J Clin Nutr* 101 (1): 153–63. doi:10.3945/ajcn.114.092601.
- Mahendran, Yuvaraj, Jyrki Ågren, Matti Uusitupa, Henna Cederberg, Jagadish Vangipurapu, Alena Stančáková, Ursula Schwab, Johanna Kuusisto, and Markku Laakso. 2014. "Association of Erythrocyte Membrane Fatty Acids with Changes in Glycemia and Risk of Type 2 Diabetes." *Am J Clin Nutr* 99 (1): 79–85. doi:10.3945/ajcn.113.069740.
- Matsuda, M., and R.A. DeFronzo. 1999. "Insulin Sensitivity Indices Obtained from Oral Glucose Tolerance Testing: Comparison with the Euglycemic Insulin Clamp." *Diabetes Care* 22 (9): 1462–70.
- Matthews, D.R., J.P. Hosker, and A.S. Rudenski. 1985. "Homeostasis Model Assessment: Insulin Resistance and Beta-Cell Function from Fasting Plasma Glucose and Insulin Concentrations in Man." *Diabetologia* 28 (7): 412–19.
- Newsholme, Philip, Deirdre Keane, Hannah J. Welters, and Noel G. Morgan. 2007. "Life and Death Decisions of the Pancreatic Beta-Cell: The Role of Fatty Acids." *Clin Sci* 112 (1): 27–42. doi:10.1042/CS20060115.
- Retnakaran, R., Y. Qi, M.I. Goran, and J.K. Hamilton. 2009. "Evaluation of Proposed Oral Disposition Index Measures in Relation to the Actual Disposition Index." *Diabet Med* 26 (12): 1198–1203.