

# The role of the growth hormone–insulin-like growth factor axis in glucose homeostasis

R. I. G. Holt, H. L. Simpson\* and P. H. Sönksen\*

Endocrinology Sub-Division, Fetal Origins of Adult Disease Division, University of Southampton, Southampton General Hospital, Southampton, and  
\*Department of Medicine, King's College School of Medicine & Dentistry, St Thomas Hospital Campus, London, UK

Accepted 31 May 2002

## Abstract

Homeostatic mechanisms normally maintain the plasma glucose concentration within narrow limits despite major fluctuations in supply and demand. There is increasing evidence that the growth hormone (GH)–insulin-like growth factor (IGF) axis may play an important role in glucose metabolism. GH has potent effects on intermediary metabolism, some of which antagonize the actions of insulin. In contrast, IGF-I has insulin-like actions, which are, in the case of glucose metabolism, opposite to those of GH. There is often deranged glucose metabolism in situations where GH is deficient or in excess. The clinical administration of GH or IGF-I results in altered glucose metabolism and changes in insulin resistance. Despite these observations, the precise role of GH and IGF-I and their interactions with insulin in controlling normal glucose homeostasis are unknown. In diabetes, GH secretion is abnormally increased as a result of reduced portal insulin resulting in impaired hepatic IGF-I generation. Evidence suggests that this may contribute to the development of diabetic microvascular complications. IGF-I ‘replacement’ in diabetes is under investigation and new methods of delivering IGF-I as a complex with IGFBP-3 offer exciting new prospects.

*Diabet. Med.* 20, 3–15 (2003)

**Keywords** growth hormone, insulin-like growth factor, glucose metabolism, growth hormone deficiency, insulin resistance


## Introduction

Normally plasma glucose concentration is maintained within a narrow range despite wide fluctuations in supply and demand. Although under normal physiological conditions insulin is the prime regulator of glucose metabolism, there is increasing evidence that the growth hormone (GH)–insulin-like growth factor-I (IGF-I) axis plays an important contributory role. GH has potent effects on intermediary metabolism, amongst which is the ability to antagonize the actions of insulin. Although GH and IGF-I exert opposite effects on glucose metabolism, the precise role of these two hormones in controlling normal glucose homeostasis is unknown. The purpose of this review is to examine the available evidence to address this question. The data for this review reflect our own clinical and academic interests and experience. We have also undertaken a comprehensive search using multiple electronic databases and hand searches of reference lists of articles.

## Growth hormone and glucose homeostasis

In the 1930s, Houssay and Biasotti observed that hypophysectomy reduced the hyperglycaemia of experimental diabetes in dogs [1]. The diabetogenic factor isolated from pituitaries was found to have growth-promoting activity and was named ‘growth hormone’ [2]. Confirmation of the diabetogenic properties of GH was made after it was administered in excess to experimental animals and man [3,4]. Following the development of reliable GH assays, human studies showed the importance of GH in glucose metabolism [5]. In both healthy subjects and those with Type 1 diabetes, a pulse of GH increases fasting hepatic glucose output, by increasing hepatic gluconeogenesis and glycogenolysis, and decreases peripheral glucose utilization through the inhibition of glycogen synthesis and glucose oxidation [6–10]. Growth hormone stimulates lipolysis with the release of glycerol and non-esterified fatty acids (NEFA). This provides a further mechanism for the diabetogenic properties of GH through the effect of NEFA to increase hepatic glucose output and decrease peripheral glucose oxidation according to the glucose–fatty acid cycle [11–13].

Correspondence to: Dr Richard I. G. Holt, Senior Lecturer in Endocrinology & Metabolism, Level D (MP811) South Academic Block, Southampton General Hospital, Tremona Road, Southampton SO16 6YD, UK. E-mail: righ@soton.ac.uk



**Figure 1** Growth hormone–insulin-like growth factor-I axis in Type 1 diabetes.

## GH and diabetes

Growth hormone concentrations are up to two to three times higher in individuals with diabetes compared with healthy subjects (Fig. 1) [14]. This reflects an increase in pituitary secretion, as the metabolic clearance rate of GH in diabetes is normal [15]. A series of studies by Wurzbürger *et al.* has shown the importance of portal insulin concentrations in regulating hepatic IGF-I generation and for the normal functioning of the GH-IGF-I axis. GH administration in Type 1 diabetes has little effect on serum IGF-I levels, while residual insulin secretion, as reflected by plasma C-peptide levels, determines the degree of GH hypersecretion [16–19].

Although there is dispute about the relative importance of different hormones in the causation of the ‘dawn phenomenon’, there is evidence that nocturnal GH hypersecretion plays an important role in the early morning reduction of insulin sensitivity and rise in plasma glucose [20,21]. Patients with Type 1 diabetes who are GH-deficient do not exhibit the ‘dawn phenomenon’, and a single bolus of GH to these individuals decreases morning insulin sensitivity. The mechanism is probably not simply secondary to an increase in NEFA, because insulin sensitivity decreases even if the lipid concentrations are maintained at a constant level with a heparin infusion [22].

## Growth hormone and the microvascular complications of diabetes

In 1953, Poulsen presented the case of a woman with Type 1 diabetes and background diabetic retinopathy, which regressed after she developed panhypopituitarism after post-partum pituitary necrosis [23]. Subsequently in the 1960s, pituitary ablation was shown to result in a regression of diabetic retinopathy, which could be related to the degree of GH deficiency [24]. These observations led to the hypothesis that GH may be involved in the development on angiopathic

diabetic complications [25,26]. Theoretically, this susceptibility may be further augmented in Type 1 diabetic patients by the supra-physiological levels of insulin in the retina, nerve and kidney produced by insulin injections into the systemic rather than portal circulation. Although effective, pituitary ablation was associated with significant morbidity and mortality and was superseded by retinal photocoagulation [27]. Patients with Type 1 diabetes and GH deficiency had decreased rates of retinopathy, supporting a role for GH development of microvascular complications [28]. There is, however, no evidence that GH replacement therapy causes an increased incidence of retinopathy in GH-deficient patients with or without diabetes [29]. Furthermore, many other growth factors, including IGF-I, have been implicated in the development of microangiopathic complications [30–32]. This remains a controversial area, as some advocate a protective role for IGF-I in the development of microangiopathic complications of diabetes [33]. According to this hypothesis, IGF-I replacement therapy in patients with diabetes will restore the normal feedback control on GH hypersecretion and thus reduce the risk of developing microvascular complications as well as glycaemic control.

## Growth hormone excess

## Animal models

Transgenic animals, which over-express GH, have added to our understanding of the role of the GH-IGF axis in glucose metabolism [34,35]. These animals develop insulin resistance, marked hyperinsulinaemia, hyperglycaemia, and hypertriglyceridaemia in association with a number of molecular abnormalities. The number of hepatic insulin receptors is reduced to around 50% of normal in transgenic animals, although the affinity of the receptor is unaffected [36,37]. The phosphorylation status of the insulin receptor and the insulin receptor substrate-1 (IRS-1) is increased so that the insulin receptor and IRS-1 are maximally activated in the resting state [38,39]. The basal association of phosphatidylinositol 3-kinase (PI3-kinase) with IRS-1, as well as PI3-kinase activity, are also increased to maximal levels. This maximal activation of the insulin receptor under basal conditions has the effect of making the tissue insensitive to further stimulation by insulin *in vivo*.

The transgenic animals have modified expression of several genes coding for proteins regulating carbohydrate metabolism. Glycogen synthase and glycogen phosphorylase activity in muscle and liver of transgenic mice is reduced and remains unchanged in liver after feeding [34]. There are also decreases in glucokinase and GLUT-2 mRNA concentrations [34].

The effects of GH on insulin signalling are directly related to the hyperinsulinaemia. If this is normalized by the administration of streptozotocin, the number of insulin receptors and the insulin-stimulated auto-phosphorylation activity of the insulin receptor return to control values [40].

## Human studies

### Acromegaly

Patients with acromegaly develop insulin resistance and hyperinsulinaemia, while up to 40% become diabetic [41,42]. Glucose intolerance usually resolves rapidly after normalization of GH concentrations with surgery or radiotherapy [43], but remission of diabetes may take several years [44]. The effect of somatostatin analogue therapy on glucose metabolism is more inconsistent [45]. Although somatostatin analogues effectively reduce GH secretion and IGF-I production, insulin secretion is also reduced and so therapy may be accompanied by either improvements or deterioration in glucose tolerance, depending on the relative effect on each of these hormones. The GH receptor antagonist, pegvisomant, results in an improvement in insulin resistance in acromegaly [46]. As this compound results in a decrease in IGF-I levels, it is possible that pegvisomant may worsen insulin resistance in patients with diabetes who already have low serum IGF-I. Antagonizing GH action in Type 1 diabetes may also result in more hypoglycaemic episodes, with increased hypoglycaemia unawareness, as occurs in patients with Type 1 diabetes and GH deficiency [17].


### Adult growth hormone deficiency

There has been a considerable advance in the understanding of glucose metabolism in adults with GH deficiency (GHDA). GHDA is commonly associated with a decrease in serum IGF-I and so it is unclear whether the insulin resistance of GHDA is secondary to decreased IGF-I concentrations or is a direct consequence of reduced GH action. However, serum IGF-I concentrations in GHDA may overlap with the age-matched normal range for IGF-I. A further understanding of the relative contributions of GH and IGF-I to glucose metabolism could be gained by assessing whether insulin sensitivity in GHDA is related to the pretreatment IGF-I concentration.

There is, however, evidence to suggest that acute decreases in GH and IGF-I concentrations are not associated with insulin resistance. If the GH receptor antagonist, pegvisomant, is administered, GH action is blocked and serum IGF-I concentrations are rapidly reduced. Under these circumstances, insulin sensitivity remains unchanged [47].

GHDA is associated with changes in body composition which lead to a reduction in lean body mass and increase in fat mass, predominantly central adiposity. It appears that the central obesity has an aetiological role in the development of insulin resistance and hyperinsulinaemia in GHDA, as fasting insulin and C-peptide concentrations correlate positively with body mass index, waist to hip ratio and central fat mass [48]. A significant difference in insulin sensitivity remains between GHDA and normal controls, however, even after corrections are made for body fat.

The molecular mechanisms underlying the insulin resistance of GHDA remain unclear. GHDA has decreased glycogen stores and a 50–64% reduction in insulin-mediated glucose utilization (Fig. 2) [49–51]. Glycolytic flux and glycogen



**Figure 2** Abnormalities in insulin-stimulated glucose metabolism in growth hormone deficiency.

synthesis are normal at baseline, but insulin activation of these pathways is reduced [50]. The decreased activation of muscle glycogen synthase by insulin accounts for the reduced muscle glycogen content. Insulin-induced muscle hexokinase activity is attenuated in GHDA, leading to normal or reduced concentrations of glucose-6-phosphate despite increased intramuscular cellular glucose levels [50]. In addition to these peripheral abnormalities, there are defects in pancreatic  $\beta$ -cell function; given the degree of insulin resistance, insulin and C-peptide secretion is often inappropriately low [52].

### Effect of GH replacement therapy in GHDA

The acute and chronic effects of rhGH replacement on glucose metabolism in adults have been reported in a number of GH trials (Table 1) [52–60]. In the short term, there is a reduction in insulin sensitivity in association with an increase in plasma glucose and insulin concentrations. Long-term, insulin sensitivity gradually improves in association with an increase in lean body mass and reduction in fat mass. There are, however, discrepancies between studies, but these may reflect duration of GHDA, differences in body composition and the effect of ageing on insulin resistance [61].

There are data from post-marketing surveillance studies looking at any adverse effects of long-term rhGH replacement. The two largest are the KIMS (Pharmacia and Upjohn International Metabolic Database) database and HypoCCS (The hypopituitary control and complications study) run by Eli Lilly. KIMS is a pharmaco-epidemiological survey of rhGH replacement in GHDA and currently comprises over 6000 patients, with a total of 10 000 patient years follow-up (mean duration of replacement 2.8 years). The majority of the 32 individuals, who have developed diabetes in the KIMS database, were obese or had a family history of diabetes, while one had impaired glucose tolerance prior to starting rhGH replacement therapy. The number developing diabetes is in line with the expected increase in diabetes that would be seen with increasing age in the general population and, considering the

**Table 1** Growth hormone (GH) trials examining the effect of rhGH on glucose metabolism

| Trial                | rhGH dose                   | Method                | Duration   | Effect of GH on glucose metabolism  | Study |
|----------------------|-----------------------------|-----------------------|------------|---|-------|
| DBPC crossover study | 0.25 IU/kg per week         | Clamp                 | 6 weeks    | Glu and Ins ↑ and Si ↓  | [53]  |
| Open study           | 0.25 IU/kg per week         | IVGTT                 | 6 months   | Glu and Ins, Si →   |       |
|                      |                             |                       | 4 weeks    | Glu, Ins and C-pep, 1st and 2nd Ins secretion after 1 week ↑ Ins-mediated Glu disposal and Glu decay ↓. Fractional Ins MCR →                        | [54]  |
|                      |                             |                       | 3 months   | Glu, Ins and C-pep ↑. All other variables →   |       |
| DBPC                 | 2 IU/m <sup>2</sup> per day | OGTT                  | 4 months   | Glu, Ins and C-pep ↑, Si & Sg ↓   | [52]  |
| Open label           | 1.6 IU/day                  | IVGTT and HOMA        | 30 months  | 3/11 developed IGT. Si ↓, Ins ↑, Ins and C-pep MCR ↑  | [55]  |
| DBPC                 | 0.07 IU/kg per day          | Fasting isotope study | 1 month    | Glu, Ins and C-pep ↑, Glucose Ra ↑, glucose MCR ↑   | [56]  |
|                      |                             |                       | 6 months   | Glu, Ins and C-pep ↑, Glucose Ra unchanged, glucose MCR ↑   |       |
| Open                 | 0.125 IU/kg per week        | IVGTT                 | 12 months  | HbA <sub>1c</sub> ↑   | [57]  |
|                      |                             |                       | 18 months  | Glu and Ins, HbA <sub>1c</sub> : no change from baseline values. Sg ↑ and Si ↓  |       |
| Open                 | 0.01–0.05 IU/kg per day     | OGTT                  | 12 months  | Glu and Ins ↑. AUC for Glu and Ins ↑  | [58]  |
|                      |                             |                       | 48 months  | Glu N, Ins ↑ (but lower than 1 year value) AUC for Glu and Ins N  |       |
| Open                 | 0.025 IU/kg per day         | Fasting data          | 120 months | Glu, Ins and C-pep the same compared with untreated GHDA. No healthy control group  | [60]  |
| Open                 |                             | OGTT                  | 84 months  | Glu and Ins → in rhGH-treated group. Si ↓ and Ins AUC ↑ in untreated patients. Comparison with GHDA not treated with GH or only treated for < 18/12 | [59]  |

GHDA, GH-deficient adults; DBPC, double-blind placebo-controlled study; OS, open study; clamp, hyperinsulinaemic-euglycaemic clamp; IVGTT, intravenous glucose tolerance test; OGTT, oral glucose tolerance test; Glu, fasting plasma glucose; Ins, fasting plasma insulin; C-pep, C-peptide; AUC, area under the curve; Si, insulin sensitivity; Sg, glucose effectiveness; MCR, metabolic clearance rate; HOMA, homeostasis model assessment of insulin sensitivity; Ra, glucose appearance.

concomitant risk factors, the incidence may be less than would be expected for an age and weight-matched population [62,63]. HypoCCS consists of 1881 rhGH-replaced patients (mean duration of replacement 2 years) and a control group of 352 GHDA patients not receiving GH, and has shown no increase in the incidence of diabetes compared with the control group [64].

### Metabolic syndrome

The 'metabolic syndrome' is a cluster of cardiovascular risk factors including central obesity, dyslipidaemia, hypertension, insulin resistance and Type 2 diabetes [65]. There are similarities between the metabolic syndrome and untreated GHDA, including central obesity and insulin resistance, which suggest that abnormalities of GH secretion may be important in the metabolic aberrations of the metabolic syndrome [66].

Central adiposity is associated with a blunted secretion of GH [67]. Although this could be a consequence of obesity, the reduction in GH secretion may have causal effects. We have shown that GH secretion is related more to age and fitness than adiposity itself, suggesting that the changes in body composition with age may be a consequence of the GH hyposecretion rather than *vice versa* [68]. In a further study, the administration of GH (9.5 µg/kg) to 30 centrally obese men for 9 months led to a reduction in total body fat, in particular abdominal subcutaneous and visceral adipose tissue, in association with an improvement in insulin sensitivity, through an increase in the glucose disposal rate [69].

## Insulin-like growth factors and glucose homeostasis

Insulin-like growth factor-I is a single chain peptide, which has structural homology with pro-insulin. Although IGF-I is synthesized widely [70], the majority of circulating IGF-I is derived from the liver [71]. Its synthesis is regulated by GH, insulin and nutritional intake [72–74]. It has profound effects on the regulation of proliferation and differentiation of many cell types, as well as metabolic effects, which are similar to those of insulin, including actions on glucose metabolism (Table 2).

### Animal studies

#### IGF-I transgenic animals

The issue of possible functional redundancies in the insulin and IGF systems *in vivo* has been addressed using knockout mice for genes encoding IGF-I or the IGF-I receptor [75–77]. The severity of the phenotypes, however, has prevented in depth examination of glucose metabolism in these mice, as 95% die. If the mice, which lack IGF-I, IGF-II or the IGF-I receptor, survive, they do not develop diabetes or ketoacidosis [78].

There have been further insights by specifically manipulating the IGF-I gene in different tissues [79,80]. If the gene for IGF-I is specifically knocked out in the liver, serum IGF-I concentrations fall to 15–25% that of wild-type animals, whilst GH increases six-fold. The liver IGF-I-deficient animals

**Table 2** Comparison between insulin and insulin-like growth factor (IGF)-I

|                          |                            | Insulin                    | IGF-I  |
|--------------------------|----------------------------|----------------------------|--|
| Structure                |                            | Single-chain polypeptide   | Single-chain polypeptide                                 |
| Weight, kD               |                            | 5807                       | 7500   |
| Chromosome               |                            | 19                         | 12   |
| Binding proteins         |                            | None                       | 6 high affinity<br>4 low affinity IGFBP related peptides |
| Receptor                 |                            | Transmembrane glycoprotein | Transmembrane glycoprotein                               |
| Post receptor Signalling |                            | IRS-1<br>Tyrosine kinase   | IRS-1<br>Tyrosine kinase                                 |
| Metabolic actions        |                            |                            |  |
| Glucose                  | Hepatic glucose production | ↓↓↓                        | ↓  |
|                          | Peripheral utilization     | ↑ (only at high doses)     | ↑  |
| Protein                  | Protein synthesis          | →                          | ↑  |
|                          | Protein degradation        | ↓                          | →  |
| Free fatty acids         | Lipolysis                  | ↓                          | ? ↓  |

have similar glucose concentrations during fasting and following glucose challenge to normal animals but are hyperinsulinaemic and insulin-resistant. Interestingly, the mice lacking liver IGF-I were leaner than wild-type mice [79]. Although these data show a role for IGF-I in glucose homeostasis, it remains unclear whether the insulin resistance results directly from a reduction in serum IGF-I, indirectly through GH hypersecretion or via changes in body composition.

#### Effects of IGF-I administration in animals

When IGF-I is given to rats as an intravenous infusion, it causes hypoglycaemia by stimulating peripheral glucose uptake, glycolysis and glycogen synthesis, but has only a minimal effect on hepatic glucose production [81]. Using a hyperinsulinaemic euglycaemic clamp, IGF-I has been shown to suppress hepatic glucose output, although to a lesser degree than insulin at the doses used, and increase peripheral glucose utilization in dogs [82]. As hepatic expression of the IGF-I receptor is low, it is likely that this effect of IGF-I on hepatic glucose output is by an indirect mechanism.

#### Effects of IGF-I administration in humans

In 1963, Froesch *et al.* observed that serum had more insulin-like activity than could be explained by the insulin concentration alone [83]. They termed this phenomenon ‘non-suppressible insulin-like activity’ (NSILA) and went on to describe that, compared with insulin, the hypoglycaemic action of NSILA was more pronounced in muscle than adipose tissue [84]. NSILA was isolated and fully characterized in 1978, shown to be identical to Somatomedin C and termed ‘insulin-like growth factor-I’ [85].

The effects of an intravenous IGF-I infusion are similar to those described in animals and lead to hypoglycaemia [86]. A single intravenous dose of 100 µg/kg results in the rapid onset of symptomatic hypoglycaemia and is equipotent to 0.15 IU/kg of insulin [87]. A continuous intravenous infusion of IGF-I causes a 50% fall in C-peptide levels despite the maintenance

of euglycaemia [88,89]. Insulin sensitivity is increased with respect to glucose by IGF-I through increased peripheral glucose uptake and decreased hepatic glucose production [90,91]. A subcutaneous infusion of IGF-I also causes hypoglycaemia, although the effect is slower in onset than insulin and decreases more slowly after the infusion is stopped because of the presence of the IGF binding proteins. The IGF-I infusion results in lower GH levels, which may also contribute to the improvements in insulin sensitivity.

## IGF-I and diabetes

### Type 1 diabetes

Insulin regulates hepatic IGF-I production and has an independent effect as well as an additive effect to GH [92,93]. Insulin-deficient diabetic rats have decreased hepatic GH binding whilst *in vitro* insulin has been shown to increase rat hepatocyte GH receptor expression [92]. However, in recent onset or mild streptozotocin-induced diabetes in rats, GH binding may be normal, implying that there may be additional post-receptor defects [94]. Insulin also increases IGF-I mRNA concentrations directly by increasing transcript stability rather than by increasing transcription [95].

Serum IGF-I concentrations are reduced in patients with Type 1 diabetes and are not normalized by intensive insulin treatment [94,96]. As the major source of circulating IGF-I is the liver and IGF-I levels are inappropriately low for the higher GH levels seen in Type 1 diabetes, it is thought that there is an acquired state of hepatic GH resistance. Seven days of rhGH treatment has little effect on IGF-I levels in patients with Type 1 diabetes. This points to the importance of portal insulin for hepatic IGF-I secretion [26]. The route of insulin administration in Type 1 diabetes appears important, as insulin delivered by the hepatic portal route (as occurs normally *in vivo*) is better at normalizing IGF-I levels and hepatic IGF-I expression in diabetic rats than subcutaneous insulin despite there being no difference in glycaemic control [73]. When insulin is given to

patients with Type 1 diabetes by a continuous intraperitoneal infusion using an implantable pump (CPII), portal insulin levels increase and there is near-normalization of IGF-I levels [97]. Low circulating IGF-I may explain the increased GH secretion seen in Type 1 diabetes through reduced negative feedback control. rhIGF-I replacement therapy in adolescents and adults with Type 1 diabetes results in a reduction in overnight GH secretion, indicating restoration of the normal negative feedback on pituitary GH secretion [98,99]. Thus IGF-I may exert its effect on glucose metabolism directly or indirectly by inhibiting GH secretion.

### IGF-I treatment in Type 1 diabetes

There has been interest in treating patients with Type 1 diabetes with IGF-I at a replacement dose to correct the derangements of the GH-IGF axis and to exploit its hypoglycaemic actions. When a single dose of IGF-I (40 µg/kg) was administered to adolescents with Type 1 diabetes, hepatic insulin sensitivity increased and the glucose production rate fell [99,100]. Over 7 days, IGF-I (40 µg/kg bd) increased peripheral glucose uptake and reduced proteolysis, despite a reduction in insulin requirement to maintain euglycaemia [98,101].

After 3 months treatment with 40 µg/kg of rhIGF-I at night, insulin sensitivity and HbA<sub>1c</sub> improved in adolescents with Type 1 diabetes. This was associated with a reduction in the insulin dose required to maintain glycaemic control [102]. Although these improvements were not maintained over the 6 months of the study, it was thought that the deterioration in control was related to poor compliance with the multi-injection regime rather than a reduction in the biological effect of IGF-I. More recently, the administration of a combination of IGF-I and IGFBP-3 over 2 weeks resulted in improvements in insulin sensitivity together with a reduction in the insulin requirements, without the side-effects that are associated with IGF-I alone [103].

It is unclear whether the improvement in metabolic control is achieved through a reduction in GH secretion or through a direct hypoglycaemic action of IGF-I. In an attempt to address this issue, Simpson *et al.* investigated the effects of IGF-I using an octreotide and glucagon infusion to suppress endogenous GH secretion in patients with Type 1 diabetes. IGF-I administration led to a decrease in hepatic glucose production, an increase in peripheral glucose utilization and an increase in glucose metabolic clearance rate when compared with both GH and placebo, suggesting that IGF-I has an independent effect on glucose metabolism [104].

### IGF-I treatment in Type 2 diabetes

Similar trials have been performed to those involving Type 1 diabetic patients, but the doses of rhIGF-I used have been larger. rhIGF-I (80–120 µg/kg bd) results in an improvement in glycaemic control with reduced fasting plasma glucose, insulin and C-peptide concentrations in association with a

sustained rise in serum IGF-I to values often above the normal physiological range [105,106]. rhIGF-I therapy was associated with a reduction in fat mass, which may partly explain the improved insulin sensitivity.

The use of high doses of IGF-I has been limited by a significant number of side-effects, particularly jaw ache, optic disc swelling, arthralgia, myalgia and headache [106]. The exact cause of these side-effects is unknown but it has been hypothesized that they result from the high concentrations of free IGF-I that occur after the supra-physiological IGF-I doses [107]. IGF-I was subsequently withdrawn from development as an adjunct to treatment in diabetes. Recently, 2 weeks treatment with combined IGF-I/IGFBP3 in Type 2 diabetes resulted in a reduction in insulin doses and fasting blood glucose concentrations, and an increase in free and total IGF-I concentrations. Notably different from previous studies with IGF-I alone, this was achieved with minimal side-effects [108].

### IGF-I treatment in growth hormone deficiency

The metabolic actions of IGF-I have been studied by administering IGF-I to severely GH-deficient young adults [109,110]. IGF-I and GH both increased lean body mass and decreased the percent fat mass, by promoting lipolysis and lipid oxidation. Seven days' treatment with rhIGF-I had no effect on hepatic glucose output, while rhGH increased hepatic glucose output. Both hormones increased lipid oxidation. Interestingly, after 8 weeks of treatment, both hormones increased hepatic glucose output, but in contrast to rhGH treatment, rhIGF-I was not associated with decreased carbohydrate oxidation or increased glucose and insulin concentrations. It is not clear why rhIGF-I should have increased hepatic glucose output in these studies, as the majority of published studies show IGF-I to have either decreased or had no effect on hepatic glucose output.

### IGF-I treatment in severe insulin resistance syndromes

Extreme insulin resistance occurs in patients who have primary defects in insulin action at the receptor or post-receptor levels, and may lead to frank diabetes mellitus that does not respond to insulin therapy.

The first report of the use rhIGF-I to treat insulin resistance was in a girl with Type 1 diabetes who required insulin doses of > 2000 IU/h [111]. During an episode of diabetic ketoacidosis, she was treated with intravenous rhIGF-I, which resulted in a rapid lowering of her plasma glucose concentration. After the acute episode, she was treated with weekly subcutaneous rhIGF-I (500 µg/kg) maintaining acceptable glycaemic control. She did not respond to a lower dose, and therapy had to be discontinued because of side-effects. Further studies have corroborated these findings in syndromes of severe insulin resistance with and without receptor abnormalities (Table 3) [112–117]. Although these studies show an improvement in glycaemic control and a reduction in the dose of insulin

**Table 3** Studies involving the use of rhIGF-I in the treatment of severe insulin resistance

| Diagnosis                                      | IGF-I dose             | Duration                             | Effect   | Reference |
|--|------------------------|--------------------------------------|--|-----------|
| Rabden–Mendenhall syndrome                     | 100 µg/kg bd           |                                      | Glu ↓ acutely but not chronically  | [112]     |
| Type A insulin resistance                      |                        |                                      | Glu, Ins and C-pep ↓   | [113]     |
| Severe insulin resistance                      | sc rhIGF-I (150 µg/kg) | 5 days                               | Glu, Ins and C-pep ↓   | [114]     |
|  |                        |                                      | Lipids →   |           |
| Type A insulin resistance without IR mutations | 100 µg/kg              | 4 weeks                              | Glu, Ins, fructosamine ↓. Si ↓ (shown by IVGTT in one patient)   | [115]     |
| Severe insulin resistance without IR mutations | 100 µg/kg bd           | 1 month                              | Diabetic patients ( <i>n</i> = 4): HbA <sub>1c</sub> and ins requirement ↓. Ins ↓ in non-diabetic patients | [116]     |
| 11 subjects IR and post receptor defects       | 100–400 µg/kg bd       | Mean 9 months<br>Range 1.5–16 months | Glu and Ins ↓ (acutely and chronic)  | [117]     |

Glu, Fasting plasma glucose; Ins, fasting plasma insulin; C-pep, C-peptide; Si, insulin sensitivity; IR, insulin receptor.

required, the doses of rhIGF-I used were high, as was the rate of side-effects.

The mechanisms by which this fall in glucose is achieved have not been studied in humans, but mice, which are genetically deficient for the insulin receptor, provide one possible model [118]. In this model, IGF-I led to phosphorylation of IGF-I receptors and increased PI3-kinase p85 subunit concentrations in the liver and skeletal muscle, suggesting that IGF-I stimulated glucose uptake via the IGF-I receptor in a PI3-kinase-dependent manner. The failure of IGF-I to prevent the death of these animals suggests that IGF-I receptors cannot effectively mediate all the metabolic actions of insulin receptors.

#### IGF-I treatment and GH insensitivity (Laron) syndromes

The relative contributions of low GH and IGF-I concentrations to the alterations in insulin sensitivity in GH deficiency are difficult to dissect. There are, however, insights from patients with primary GH resistance, secondary to inherited molecular defects of the GH receptor or post-receptor pathways [119]. The clinical course is characterized by progressive obesity, insulin resistance, and a tendency for hypoglycaemia, followed later in life by hyperinsulinaemia, glucose intolerance and even diabetes. In adulthood, β-cell failure may ensue, resulting in insulinopenia and Type 2 diabetes [119].

Circulating GH concentrations are increased and GH therapy produces no clinical benefit. In contrast, IGF-I treatment of children with Laron syndrome can normalize many of these biochemical abnormalities, including glucose intolerance and hyperinsulinaemia. IGF-I treatment may even be complicated by hypoglycaemia [120].

#### Gene deletion of IGF-I

Woods *et al.* reported a patient with a homozygous partial deletion of the IGF-I gene, resulting in IGF-I deficiency [121]. This child had insulin resistance, hyperinsulinaemia and short stature. IGF-I therapy resulted in a dose-dependent improvement of insulin sensitivity into the normal range and a reduction in basal and stimulated insulin concentrations [122].

### IGF binding proteins and glucose metabolism

The IGFs are present in the circulation and throughout the extracellular space almost entirely bound to members of a family of at least six high-affinity IGF binding proteins (IGFBP-1 to -6) [123,124]. Less than 1% of the IGF in the circulation is free, with 90% bound in a stable 150-kD ternary complex comprising IGF-I or IGF-II, IGFBP-3 and an acid-labile subunit. IGFBP-1, -2, -4 and -6 bind most of the remaining IGF in a binary complex of 30–40 kD, while IGFBP-5 is also capable of forming a ternary complex with IGF-I and the acid-labile subunit. All six IGFBPs have negligible affinity for insulin. The IGFBPs co-ordinate and regulate the biological functions of the IGFs, although we have little knowledge about how this works. They transport IGFs and control their efflux from the circulation. The IGFBPs prolong the half-lives of IGFs and regulate their metabolic clearance. Alteration in the quantity and function of the IGFBPs may therefore affect the hypoglycaemic effect of the IGFs.

#### IGFBP-1

Although not the most predominant IGF-I binding protein, IGFBP-1 is the only binding protein to show rapid regulation of concentration *in vivo*. Under many circumstances, IGFBP-1 is inversely related to insulin concentration [125] and has a diurnal variation, with the highest concentrations being overnight when insulin levels are lowest [126]. IGFBP-1 concentrations are high in Type 1 diabetes and correlate with glycosylated haemoglobin [127]. A bolus of intravenous IGFBP-1 causes hyperglycaemia and prevents exogenous IGF-I from having a hypoglycaemic action in rats [128]. In normal human subjects there is an inverse relationship between IGFBP-1 and free IGF-I levels. Although a single intravenous bolus of IGF-I acutely stimulates IGFBP-1 production [129], continuous subcutaneous administration of IGF-I over 3 days leads to an increase in free and total IGF-I concentrations with a decrease in IGFBP-1 concentrations [130]. These data suggest that IGFBP-1 can acutely regulate bioactivity of IGF-I concentrations and suggest a role for IGF-I in glucose regulation.

There are situations where serum IGFBP-1 is not inversely associated with insulin. One example of this is chronic liver disease, where both insulin and IGFBP-1 concentrations are higher than in normal controls [131]. This may reflect differences in the importance of portal vs. peripheral insulin concentrations. In chronic liver disease, porto-systemic shunts develop, which result in increased peripheral insulin concentrations at the expense of hepatic hypo-insulinaemia. This may explain the observation that IGFBP-1 concentrations are higher in children with portal hypertension compared with those children without portal hypertension [132].

### IGFBP-1 transgenic animals

Over-expression of IGFBP-1 in transgenic mice has provided insight into the physiological role of this binding protein in modulating the metabolic effects of the IGFs. In most strains of IGFBP-1 transgenic mice, the animals develop fasting hyperglycaemia, impaired glucose tolerance, and modest insulin resistance in skeletal muscle and hepatic tissue [133,134].

The effect of IGFBP-1 is to attenuate the hypoglycaemic action of IGF-I. If insulin or des (1–3) IGF-I, an analogue of IGF-I that does not bind IGFBP-1, is infused, there are no differences in the hypoglycaemic response between normal and animals transgenic for IGFBP-1 [133]. Similarly, the glucose response to subcutaneous insulin was similar in transgenic and wild-type mice, suggesting that constitutive over-expression of IGFBP-1 results in impaired glucose tolerance despite normal insulin sensitivity [135]. There is sexual dimorphism in at least one strain of transgenic mice, with male mice having higher stimulated glucose and insulin levels than females [134].


The transgenic animals have increased gluconeogenesis and hepatic insulin resistance, which contributes to the fasting hyperglycaemia [136]. In isolated hepatocytes, the inhibition of glucose production from pyruvate by insulin is decreased in transgenic mice compared with wild-type animals [136]. Furthermore, serum from transgenic mice results in more glucose production by hepatocytes than serum from wild-type mice [136].

### IGFBP-2 transgenic animals

Transgenic mice, which over-express IGFBP-2, have reduced fasting serum glucose and insulin concentrations, but post-prandial concentrations are unaffected by the genotype [137].

### Non-islet cell tumour hypoglycaemia

Non-islet cell tumour hypoglycaemia (NICTH) is a rare endocrine paraneoplastic syndrome, which is characterized by recurrent fasting hypoglycaemia in the absence of insulin and C-peptide [138,139]. The precise cause of the hypoglycaemia in NICTH remains unclear (Fig. 3). IGF-II is normally synthesized as pro-IGF-II and this is processed in a stepwise fashion to form the mature protein. In NICTH, there is abnormal processing of



**Figure 3** Mechanism for hypoglycaemia in non-islet cell tumour hypoglycaemia.

IGF-II and larger isoforms of IGF-II, known as 'big IGF-II', are invariably released into the circulation. Big IGF-II binds with lower affinity to IGFBP-3 and the acid-labile subunit and consequently has higher bioavailability and bioactivity [140,141]. The switch from IGF-II to big IGF-II appears to be a key step in causing hypoglycaemia as IGF-II is over-expressed in a number of tumours, such as the embryonic Wilm's tumour, without the development of hypoglycaemia [142]. The hypoglycaemia may result through a direct hypoglycaemic action of big IGF-II and indirectly through suppression of GH secretion [143–146]. The reduction in GH secretion results in a reduction of IGF-I, IGFBP-3 and acid-labile subunit (ALS). The reduction in IGFBP-3 and ALS is associated with increased free IGF-I concentrations, which may also contribute to the hypoglycaemia [147]. GH administration in NICTH leads to increased IGFBP-3 and ALS production and has been used to treat the hypoglycaemia of NICTH successfully both in the short term [148] and long term [145]. The massive over-production of big IGF-II may also lead to saturation of IGF-I binding sites of the IGFBPs with a consequential further increase in free IGF-I.

### Conclusions

Although the relative contributions of GH and IGF-I to normal glucose homeostasis are difficult to study, there is evidence from patients, animal models and cell cultures to indicate a role for both GH and IGF-I in normal glucose homeostasis. The biological reason for the need for GH and IGF-I may be to allow partitioning of glucose to metabolically active tissues at a time of need and to organize energy storage at times of substrate excess. This is of particular importance in skeletal muscle, which is responsible for 80% of post-prandial glucose transport. Furthermore, as GH has powerful lipolytic properties, this provides glycerol as a gluconeogenic substrate and NEFA as an alternative substrate to glucose at times when glucose or glycogen may be in short supply, such as during fasting or exercise.



The GH-IGF axis is deranged in patients with diabetes and this probably contributes to both the metabolic disturbance and the susceptibility to microvascular complications. Treatment with IGF-I results in improvements in metabolic control but its use has been limited by side-effects. More recently, the combination of IGF-I and IGFBP-3 has been shown to improve metabolic control but without these side-effects. Further work will be needed to confirm the safety and efficacy of this compound.

While insulin may have the overall responsibility to control glucose concentrations, primarily through its effects on hepatic glucose output, IGF-I acting through both endocrine and local mechanisms may serve to fine tune insulin sensitivity. Finally, the contribution of GH and IGF-I to glucose homeostasis may increase at times where insulin action is reduced, either through insulin deficiency or insulin resistance.

## References

- Houssay BA, Biasotti A. La diabetes pancreatica de los perros hipofiseoprivos. *Rev Soc Argent Biol* 1930; 6: 251–296.
- Young FG. Anterior pituitary fraction and carbohydrate metabolism I; the preparation and properties of diabetogenic extracts. *J Endocrinol* 1939; 1: 339–354.
- Young FG. 'Growth' and the diabetogenic action of anterior pituitary preparations. *BMJ* 1941; 2: 897–901.
- Luft R, Ikkos D, Gemzell CA, Olivercrona H, Sjogren B. The effects of growth hormone in man. *Lancet* 1958; 1: 720–721.
- Zierler K, Rabinowitz R. Roles of insulin and growth hormone based on studies of forearm metabolism in man. *Medicine* 1963; 42: 385–395.
- Tamborlane WV, Sherwin RS, Koivisto V, Hendler R, Genel M, Felig P. Normalization of the growth hormone and catecholamine response to exercise in juvenile-onset diabetic subjects treated with a portable insulin infusion pump. *Diabetes* 1979; 28: 785–788.
- Bak JF, Moller N, Schmitz O. Effects of growth hormone on fuel utilization and muscle glycogen synthase activity in normal humans. *Am J Physiol* 1991; 260: E736–742.
- Press M, Tamborlane WV, Sherwin RS. Importance of raised growth hormone levels in mediating the metabolic derangements of diabetes. *N Eng J Med* 1984; 310: 810–815.
- Fowelin J, Attvall S, von Schenck H, Smith U, Lager I. Characterization of the insulin-antagonistic effect of growth hormone in insulin-dependent diabetes mellitus. *Diabet Med* 1995; 12: 990–996.
- Fowelin J, Attvall S, von Schenck H, Smith U, Lager I. Characterization of the insulin-antagonistic effect of growth hormone in man. *Diabetologia* 1991; 34: 500–506.
- Gerich JE, Lorenzi M, Bier DM, Tsalikian E, Schneider V, Karam JH *et al.* Effects of physiologic levels of glucagon and growth hormone on human carbohydrate and lipid metabolism. Studies involving administration of exogenous hormone during suppression of endogenous hormone secretion with somatostatin. *J Clin Invest* 1976; 57: 875–884.
- Randle PJ, Hales CN, Garland PB, Newsholme EA. The glucose fatty acid cycle: its role in insulin sensitivity and the metabolic disturbance of diabetes mellitus. *Lancet* 1963; i: 785–789.
- Ferrannini E, Barrett EJ, Bevilacqua S, DeFronzo RA. Effect of fatty acids on glucose production and utilization in man. *J Clin Invest* 1983; 72: 1737–1747.
- Hansen AP, Johansen K. Diurnal patterns of blood glucose, serum free fatty acids, insulin, glucagon and growth hormone in normals and juvenile diabetics. *Diabetologia* 1970; 6: 27–33.
- Navalesi R, Pilo A, Vigneri R. Growth hormone kinetics in diabetic patients. *Diabetes* 1975; 24: 317–327.
- Wurzbürger MI, Prelevic GM, Sonksen PH, Peric LA, Till S, Morris RW. The effects of improved blood glucose on growth hormone and cortisol secretion in insulin-dependent diabetes mellitus. *Clin Endocrinol (Oxf)* 1990; 32: 787–797.
- Wurzbürger MI, Prelevic GM, Sonksen PH. Restoration of hypoglycaemia awareness by human recombinant growth hormone. *Lancet* 1992; 339: 496–497.
- Wurzbürger MI, Prelevic GM, Sonksen PH, Balint-Peric LA, Wheeler M. The effect of recombinant human growth hormone on regulation of growth hormone secretion and blood glucose in insulin-dependent diabetes. *J Clin Endocrinol Metab* 1993; 77: 267–272.
- Wurzbürger MI, Prelevic GM, Sonksen PH, Wheeler M, Balint-Peric L. Effect of recombinant human growth hormone treatment on insulin-like growth factor (IGF-I) levels in insulin-dependent diabetic patients. *Acta Diabetologica* 1995; 32: 131–134.
- Campbell PJ, Bolli GB, Cryer PE, Gerich JE. Pathogenesis of the dawn phenomenon in patients with insulin-dependent diabetes mellitus. Accelerated glucose production and impaired glucose utilization due to nocturnal surges in growth hormone secretion. *N Eng J Med* 1985; 312: 1473–1479.
- Perriello G, De Feo P, Torlone E, Fanelli C, Santeusano F, Brunetti P *et al.* The dawn phenomenon in type 1 (insulin-dependent) diabetes mellitus: magnitude, frequency, variability, and dependency on glucose counterregulation and insulin sensitivity. *Diabetologia* 1991; 34: 21–28.
- Boyle PJ, Avogaro A, Smith L, Shah SD, Cryer PE, Santiago JV. Absence of the dawn phenomenon and abnormal lipolysis in type 1 (insulin-dependent) diabetic patients with chronic growth hormone deficiency. *Diabetologia* 1992; 35: 372–379.
- Poulsen JE. Houssay phenomenon in man: recovery from retinopathy in a case of diabetes with Simmonds disease. *Diabetes* 1953; 2: 7–12.
- Wright AD, Kohner EM, Oakley NW, Hartog M, Joplin GF, Fraser TR. Serum growth hormone levels and the response of diabetic retinopathy to pituitary ablation. *BMJ* 1969; 2: 346–348.
- Lundbaek K, Christensen NJ, Jensen VA, Johansen K, Olsen TS, Hansen AP *et al.* Diabetes, diabetic angiopathy, and growth hormone. *Lancet* 1970; 2: 131–133.
- Sonksen PH, Russell-Jones D, Jones RH. Growth hormone and diabetes mellitus. A review of sixty-three years of medical research and a glimpse into the future? *Hormone Res* 1993; 40: 68–79.
- Sharp PS, Fallon TJ, Brazier OJ, Sandler L, Joplin GF, Kohner EM. Long-term follow-up of patients who underwent yttrium-90 pituitary implantation for treatment of proliferative diabetic retinopathy. *Diabetologia* 1987; 30: 199–207.
- Alzaid AA, Dinneen SF, Melton LJ, Rizza RA. The role of growth hormone in the development of diabetic retinopathy. *Diabetes Care* 1994; 17: 531–534.
- Blank D, Riedl M, Reitner A, Schnack C, Scherthaner G, Clodi M *et al.* Growth hormone replacement therapy is not associated with retinal changes. *J Clin Endocrinol Metab* 2000; 85: 634–636.
- Hyer SL, Sharp PS, Brooks RA, Burrin JM, Kohner EM. A two-year follow-up study of serum insulinlike growth factor-I in diabetics with retinopathy. *Metabolism* 1989; 38: 586–589.
- Flyvbjerg A. Growth factors and diabetic complications. *Diabet Med* 1990; 7: 387–399.
- Chiarelli F, Santilli F, Mohn A. Role of growth factors in the development of diabetic complications. *Hormone Res* 2000; 53: 53–67.
- Janssen JA, Lamberts SW. Circulating IGF-I and its protective role in the pathogenesis of diabetic angiopathy. *Clin Endocrinol (Oxf)* 2000; 52: 1–9.

- 34 Valera A, Rodriguez-Gil JE, Yun JS, McGrane MM, Hanson RW, Bosch F. Glucose metabolism in transgenic mice containing a chimeric P-enolpyruvate carboxykinase/bovine growth hormone gene. *FASEB J* 1993; 7: 791–800.
- 35 Ikeda A, Chang KT, Matsumoto Y, Furuhashi Y, Nishihara M, Sasaki F *et al.* Obesity and insulin resistance in human growth hormone transgenic rats. *Endocrinology* 1998; 139: 3057–3063.
- 36 Balbis A, Dellacha JM, Calandra RS, Bartke A, Turyn D. Down regulation of masked and unmasked insulin receptors in the liver of transgenic mice expressing bovine growth hormone gene. *Life Sci* 1992; 51: 771–778.
- 37 Balbis A, Bartke A, Turyn D. Overexpression of bovine growth hormone in transgenic mice is associated with changes in hepatic insulin receptors and in their kinase activity. *Life Sci* 1996; 59: 1363–1371.
- 38 Dominici FP, Cifone D, Bartke A, Turyn D. Alterations in the early steps of the insulin-signaling system in skeletal muscle of GH-transgenic mice. *Am J Physiol* 1999; 277: E447–454.
- 39 Dominici FP, Cifone D, Bartke A, Turyn D. Loss of sensitivity to insulin at early events of the insulin signaling pathway in the liver of growth hormone-transgenic mice. *J Endocrinol* 1999; 161: 383–392.
- 40 Dominici FP, Balbis A, Bartke A, Turyn D. Role of hyperinsulinemia on hepatic insulin receptor concentration and autophosphorylation in the presence of high growth hormone levels in transgenic mice overexpressing growth hormone gene. *J Endocrinol* 1998; 159: 15–25.
- 41 Ezzat S, Forster MJ, Berchtold P, Redelmeier DA, Boerlin V, Harris AG. Acromegaly. Clinical and biochemical features in 500 patients. *Medicine (Baltimore)* 1994; 73: 233–240.
- 42 Colao A, Baldelli R, Marzullo P, Ferretti E, Ferone D, Gargiulo P *et al.* Systemic hypertension and impaired glucose tolerance are independently correlated to the severity of the acromegalic cardiomyopathy. *J Clin Endocrinol Metab* 2000; 85: 193–199.
- 43 Sonksen PH, Greenwood FC, Ellis JP, Lowy C, Rutherford A, Nabarro JD. Changes of carbohydrate tolerance in acromegaly with progress of the disease and in response to treatment. *J Clin Endocrinol Metab* 1967; 27: 1418–1430.
- 44 Sonksen PH. Aetiology and epidemiology of diabetes. *Br J Hosp Med* 1972; 7: 151–156.
- 45 Koop BL, Harris AG, Ezzat S. Effect of octreotide on glucose tolerance in acromegaly. *Eur J Endocrinol* 1994; 130: 581–586.
- 46 Parkinson C, Trainer PJ. GH receptor antagonists in acromegaly. *Proc Am Endocrine Soc Conference* 2001: 58.1.
- 47 Muller AF, Leebeek FW, Janssen JA, Lamberts SW, Hofland L, van der Lely AJ. Acute effect of pegvisomant on cardiovascular risk markers in healthy men: implications for the pathogenesis of atherosclerosis in GH deficiency. *J Clin Endocrinol Metab* 2001; 86: 5165–5171.
- 48 Salomon F, Cuneo RC, Umpleby AM, Sonksen PH. Interactions of body fat and muscle mass with substrate concentrations and fasting insulin levels in adults with growth hormone deficiency. *Clin Sci (Lond)* 1994; 87: 201–206.
- 49 Salomon F, Cuneo RC, Umpleby AM, Sonksen PH. Glucose and fat metabolism in adults with growth hormone deficiency. *Clin Endocrinol (Oxf)* 1994; 41: 315–322.
- 50 Alford FP, Hew FL, Christopher MC, Rantza C. Insulin sensitivity in growth hormone (GH)-deficient adults and effect of GH replacement therapy. *J Endocrinol Invest* 1999; 22: 28–32.
- 51 Hew FL, Koschmann M, Christopher M, Rantza C, Vaag A, Ward G *et al.* Insulin resistance in growth hormone-deficient adults: defects in glucose utilization and glycogen synthase activity. *J Clin Endocrinol Metab* 1996; 81: 555–564.
- 52 Rosenfalck AM, Fisker S, Hilsted J, Dinesen B, Volund A, Jorgensen JO *et al.* The effect of the deterioration of insulin sensitivity on beta-cell function in growth-hormone-deficient adults following 4-month growth hormone replacement therapy. *Growth Horm IGF Res* 1999; 9: 96–105.
- 53 Fowelin J, Attvall S, Lager I, Bengtsson BA. Effects of treatment with recombinant human growth hormone on insulin sensitivity and glucose metabolism in adults with growth hormone deficiency. *Metabolism* 1993; 42: 1443–1447.
- 54 O'Neal DN, Kalfas A, Dunning PL, Christopher MJ, Sawyer SD, Ward GM *et al.* The effect of 3 months of recombinant human growth hormone (GH) therapy on insulin and glucose-mediated glucose disposal and insulin secretion in GH-deficient adults: a minimal model analysis. *J Clin Endocrinol Metab* 1994; 79: 975–983.
- 55 Rosenfalck AM, Maghsoudi S, Fisker S, Jorgensen JO, Christiansen JS, Hilsted J *et al.* The effect of 30 months of low-dose replacement therapy with recombinant human growth hormone (rhGH) on insulin and C-peptide kinetics, insulin secretion, insulin sensitivity, glucose effectiveness, and body composition in GH-deficient adults. *J Clin Endocrinol Metab* 2000; 85: 4173–4181.
- 56 Salomon F, Umpleby AM, Cuneo RC, Sonksen PH. Protein, fat and glucose metabolism during treatment with recombinant human growth hormone in adults with growth hormone deficiency. Short and long term effects. *Endocrinol Metab* 1997; 4: 121–128.
- 57 Riedl M, Ludvik B, Pacini G, Clodi M, Kotzmann H, Wagner O *et al.* The increased insulin sensitivity in growth hormone-deficient adults is reduced by growth hormone replacement therapy. *Eur J Clin Invest* 2000; 30: 771–778.
- 58 al Shoumer KA, Gray R, Anyaoku V, Hughes C, Beshyah S, Richmond W *et al.* Effects of four years' treatment with biosynthetic human growth hormone (GH) on glucose homeostasis, insulin secretion and lipid metabolism in GH-deficient adults. *Clin Endocrinol (Oxf)* 1998; 48: 795–802.
- 59 Chrisoulidou A, Beshyah SA, Rutherford O, Spinks TJ, Mayet J, Kyd P *et al.* Effects of 7 years of growth hormone replacement therapy in hypopituitary adults. *J Clin Endocrinol Metab* 2000; 85: 3762–3769.
- 60 Gibney J, Wallace JD, Spinks T, Schnorr L, Ranicar A, Cuneo RC *et al.* The effects of 10 years of recombinant human growth hormone (GH) in adult GH-deficient patients. *J Clin Endocrinol Metab* 1999; 84: 2596–2602.
- 61 Fink RI, Kolterman OG, Griffin J, Olefsky JM. Mechanisms of insulin resistance in aging. *J Clin Invest* 1983; 71: 1523–1535.
- 62 Monson JP, Bengtsson BA, Abs R, Feldt-Rasmussen U, Wuster C. Can growth hormone therapy cause diabetes? KIMS Strategic Committee. *Lancet* 2000; 355: 1728–1729.
- 63 Wilton P. Review of KIMS data. *Proceedings of the Am Endo Soc Conference* 2001.
- 64 Hartman ML. Review of HypoCCS data. *Proceedings of the Am Endo Soc Conference* 2001.
- 65 Reaven GM. Banting lecture 1988. Role of insulin resistance in human disease. *Diabetes* 1988; 37: 1595–1607.
- 66 Johannsson G, Bengtsson BA. Growth hormone and the metabolic syndrome. *J Endocrinol Invest* 1999; 22: 41–46.
- 67 Veldhuis JD, Iranmanesh A, Ho KK, Waters MJ, Johnson ML, Lizarralde G. Dual defects in pulsatile growth hormone secretion and clearance subserve the hypsomatotropism of obesity in man. *J Clin Endocrinol Metab* 1991; 72: 51–59.
- 68 Holt RIG, Webb E, Pentecost C, Sonksen PH. Ageing and physical fitness are more important than obesity in determining exercise induced generation of growth hormone. *J Clin Endocrinol Metab* 2001; 86: 5715–5720.
- 69 Johannsson G, Marin P, Lonn L, Ottosson M, Stenlof K, Bjorntorp P *et al.* Growth hormone treatment of abdominally obese men reduces abdominal fat mass, improves glucose and lipoprotein metabolism, and reduces diastolic blood pressure. *J Clin Endocrinol Metab* 1997; 82: 727–734.
- 70 D'Ercole AJ, Stiles AD, Underwood LE. Tissue concentrations of somatomedin C: further evidence for multiple sites of synthesis and

- paracrine or autocrine mechanisms of action. *Proc Natl Acad Sci USA* 1984; **81**: 935–939.
- 71 Yakar S, Liu JL, Stannard B, Butler A, Accili D, Sauer B *et al*. Normal growth and development in the absence of hepatic insulin-like growth factor I. *Proc Natl Acad Sci USA* 1999; **96**: 7324–7329.
- 72 Mathews LS, Hammer RE, Brinster RL, Palmiter RD. Expression of insulin-like growth factor I in transgenic mice with elevated levels of growth hormone is correlated with growth. *Endocrinology* 1988; **123**: 433–437.
- 73 Russell-Jones DL, Rattray M, Wilson VJ, Jones RH, Sonksen PH, Thomas CR. Intraperitoneal insulin is more potent than subcutaneous insulin at restoring hepatic insulin-like growth factor-I mRNA levels in the diabetic rat: a functional role for the portal vascular link. *J Mol Endocrinol* 1992; **9**: 257–263.
- 74 Thissen JP, Ketelslegers JM, Underwood LE. Nutritional regulation of the insulin-like growth factors. *Endocr Rev* 1994; **15**: 80–101.
- 75 Liu JP, Baker J, Perkins AS, Robertson EJ, Efstratiadis A. Mice carrying null mutations of the genes encoding insulin-like growth factor I (Igf-1) and type 1 IGF receptor (Igf1r). *Cell* 1993; **75**: 59–72.
- 76 DeChiara TM, Efstratiadis A, Robertson EJ. A growth-deficiency phenotype in heterozygous mice carrying an insulin-like growth factor II gene disrupted by targeting. *Nature* 1990; **345**: 78–80.
- 77 Baker J, Liu JP, Robertson EJ, Efstratiadis A. Role of insulin-like growth factors in embryonic and postnatal growth. *Cell* 1993; **75**: 73–82.
- 78 Lamothe B, Baudry A, Desbois P, Lamotte L, Bucchini D, De Meyts P *et al*. Genetic engineering in mice: impact on insulin signalling and action. *Biochem J* 1998; **335**: 193–204.
- 79 Sjogren K, Wallenius K, Liu JL, Bohlooly YM, Pacini G, Svensson L *et al*. Liver-derived IGF-I is of importance for normal carbohydrate and lipid metabolism. *Diabetes* 2001; **50**: 1539–1545.
- 80 Yakar S, Liu JL, Fernandez AM, Wu Y, Schally AV, Frystyk J *et al*. Liver-specific igf-1 gene deletion leads to muscle insulin insensitivity. *Diabetes* 2001; **50**: 1110–1118.
- 81 Jacob R, Barrett E, Plewe G, Fagin KD, Sherwin RS. Acute effects of insulin-like growth factor I on glucose and amino acid metabolism in the awake fasted rat. Comparison with insulin. *J Clin Invest* 1989; **83**: 1717–1723.
- 82 Shojaae-Moradie F, Umpleby AM, Thomason MJ, Jackson NC, Boroujerdi MA, Sonksen PH *et al*. A comparison of the effects of insulin-like growth factor-I, insulin and combined infusions of insulin and insulin-like growth factor-I on glucose metabolism in dogs. *Eur J Clin Invest* 1995; **25**: 920–928.
- 83 Froesch ER, Burgi H, Ramseier EB, Bally P, Labhart A. Antibody suppressible and non-suppressible insulin-like activities in human serum and their physiological significance. An insulin assay with adipose tissue of increased precision and specificity. *J Clin Invest* 1963; **42**: 1816–1834.
- 84 Froesch ER. The physiology and pharmacology of adipose tissue lipolysis: its inhibition and implications for the treatment of diabetes. *Diabetologia* 1967; **3**: 475–487.
- 85 Rinderknecht E, Humbel RE. The amino acid sequence of human insulin-like growth factor I and its structural homology with proinsulin. *J Biol Chem* 1978; **253**: 2769–2776.
- 86 Zenobi PD, Graf S, Ursprung H, Froesch ER. Effects of insulin-like growth factor-I on glucose tolerance, insulin levels, and insulin secretion. *J Clin Invest* 1992; **89**: 1908–1913.
- 87 Guler HP, Zapf J, Froesch ER. Short-term metabolic effects of recombinant human insulin-like growth factor I in healthy adults. *N Eng J Med* 1987; **317**: 137–140.
- 88 Boulware SD, Tamborlane WV, Matthews LS, Sherwin RS. Diverse effects of insulin-like growth factor I on glucose, lipid, and amino acid metabolism. *Am J Physiol* 1992; **262**: E130–133.
- 89 Turkalj I, Keller U, Ninnis R, Vosmeer S, Stauffacher W. Effect of increasing doses of recombinant human insulin-like growth factor-I on glucose, lipid, and leucine metabolism in man. *J Clin Endocrinol Metab* 1992; **75**: 1186–1191.
- 90 Boulware SD, Tamborlane WV, Rennert NJ, Gesundheit N, Sherwin RS. Comparison of the metabolic effects of recombinant human insulin-like growth factor-I and insulin. Dose-response relationships in healthy young and middle-aged adults. *J Clin Invest* 1994; **93**: 1131–1139.
- 91 Russell-Jones DL, Bates AT, Umpleby AM, Hennessy TR, Bowes SB, Hopkins KD *et al*. A comparison of the effects of IGF-I and insulin on glucose metabolism, fat metabolism and the cardiovascular system in normal human volunteers. *Eur J Clin Invest* 1995; **25**: 403–411.
- 92 Scott CD, Baxter RC. Production of insulin-like growth factor I and its binding protein in rat hepatocytes cultured from diabetic and insulin-treated diabetic rats. *Endocrinology* 1986; **119**: 2346–2352.
- 93 Boni-Schnetzler M, Schmid C, Meier PJ, Froesch ER. Insulin regulates insulin-like growth factor I mRNA in rat hepatocytes. *Am J Physiol* 1991; **260**: E846–851.
- 94 Maes M, Underwood LE, Ketelslegers JM. Low serum somatomedin-C in insulin-dependent diabetes: evidence for a postreceptor mechanism. *Endocrinology* 1986; **118**: 377–382.
- 95 Goya L, de la Puente A, Ramos S, Martin MA, Escrivá F, Alvarez C *et al*. Regulation of IGF-I and -II by insulin in primary cultures of fetal rat hepatocytes. *Endocrinology* 2001; **142**: 5089–5096.
- 96 Amiel SA, Sherwin RS, Hintz RL, Gertner JM, Press CM, Tamborlane WV. Effect of diabetes and its control on insulin-like growth factors in the young subject with type I diabetes. *Diabetes* 1984; **33**: 1175–1179.
- 97 Hanaire-Broutin H, Sallerin-Caute B, Poncet MF, Tauber M, Bastide R, Chale JJ *et al*. Effect of intraperitoneal insulin delivery on growth hormone binding protein, insulin-like growth factor (IGF)-I, and IGF-binding protein-3 in IDDM. *Diabetologia* 1996; **39**: 1498–1504.
- 98 Carroll PV, Umpleby M, Ward GS, Imuere S, Alexander E, Dunger D *et al*. rhIGF-I administration reduces insulin requirements, decreases growth hormone secretion, and improves the lipid profile in adults with IDDM. *Diabetes* 1997; **46**: 1453–1458.
- 99 Cheetham TD, Jones J, Taylor AM, Holly J, Matthews DR, Dunger DB. The effects of recombinant insulin-like growth factor I administration on growth hormone levels and insulin requirements in adolescents with type 1 (insulin-dependent) diabetes mellitus. *Diabetologia* 1993; **36**: 678–681.
- 100 Acerini CL, Harris DA, Matyka KA, Watts AP, Umpleby AM, Russell-Jones DL *et al*. Effects of low-dose recombinant human insulin-like growth factor-I on insulin sensitivity, growth hormone and glucagon levels in young adults with insulin-dependent diabetes mellitus. *Metabolism* 1998; **47**: 1481–1489.
- 101 Carroll PV, Christ ER, Umpleby AM, Gowrie I, Jackson N, Bowes SB *et al*. IGF-I treatment in adults with type 1 diabetes: effects on glucose and protein metabolism in the fasting state and during a hyperinsulinemic-euglycemic amino acid clamp. *Diabetes* 2000; **49**: 789–796.
- 102 Acerini CL, Patton CM, Savage MO, Kernell A, Westphal O, Dunger DB. Randomised placebo-controlled trial of human recombinant insulin-like growth factor I plus intensive insulin therapy in adolescents with insulin-dependent diabetes mellitus. *Lancet* 1997; **350**: 1199–1204.
- 103 Clemmons DR, Moses AC, McKay MJ, Sommer A, Rosen DM, Ruckle J. The combination of insulin-like growth factor I and insulin-like growth factor-binding protein-3 reduces insulin requirements in insulin-dependent type 1 diabetes: evidence for in vivo biological activity. *J Clin Endocrinol Metab* 2000; **85**: 1518–1524.
- 104 Simpson HL, Jackson N, Croos P, Shojaae-Moradie F, Russell-Jones DL, Jones R *et al*. IGF-I has an independent effect on glucose

- metabolism in type 1 diabetes. *Proceedings of the Am Endo Soc Conference* 2001; OR44–1.
- 105 Zenobi PD, Jaeggi-Groisman SE, Riesen WF, Roder ME, Froesch ER. Insulin-like growth factor-I improves glucose and lipid metabolism in type 2 diabetes mellitus. *J Clin Invest* 1992; **90**: 2234–2241.
  - 106 Moses AC, Young SC, Morrow LA, O'Brien M, Clemmons DR. Recombinant human insulin-like growth factor I increases insulin sensitivity and improves glycemic control in type II diabetes. *Diabetes* 1996; **45**: 91–100.
  - 107 Frystyk J, Hussain M, Skjaerbaek C, Porksen N, Froesch ER, Orskov H. The pharmacokinetics of free insulin-like growth factor-I in healthy subjects. *Growth Horm IGF Res* 1999; **9**: 150–156.
  - 108 Clemmons DR, Moses A, Jacobson W, Rogol A, Sommer A, Allan G. IGF-I/IGFBP3 by injection or infusion to type 2 diabetes increases free IGF-I and reduces insulin requirements. *Proceedings of the Am Endo Soc Conference* 2001; OR9–4.
  - 109 Hussain MA, Schmitz O, Mengel A, Glatz Y, Christiansen JS, Zapf J *et al.* Comparison of the effects of growth hormone and insulin-like growth factor I on substrate oxidation and on insulin sensitivity in growth hormone-deficient humans. *J Clin Invest* 1994; **94**: 1126–1133.
  - 110 Mauras N, O'Brien KO, Welch S, Rini A, Helgeson K, Vieira NE *et al.* Insulin-like growth factor I and growth hormone (GH) treatment in GH-deficient humans: differential effects on protein, glucose, lipid, and calcium metabolism. *J Clin Endocrinol Metab* 2000; **85**: 1686–1694.
  - 111 Usala AL, Madigan T, Burguera B, Sinha MK, Caro JF, Cunningham P *et al.* Brief report: treatment of insulin-resistant diabetic ketoacidosis with insulin-like growth factor I in an adolescent with insulin-dependent diabetes. *N Eng J Med* 1992; **327**: 853–857.
  - 112 Quin JD, Fisher BM, Paterson KR, Inoue A, Beastall GH, MacCuish AC. Acute response to recombinant insulin-like growth factor I in a patient with Mendenhall's syndrome. *N Eng J Med* 1990; **323**: 1425–1426.
  - 113 Schoenle EJ, Zenobi PD, Torresani T, Werder EA, Zachmann M, Froesch ER. Recombinant human insulin-like growth factor I (rhIGF I) reduces hyperglycaemia in patients with extreme insulin resistance. *Diabetologia* 1991; **34**: 675–679.
  - 114 Zenobi PD, Glatz Y, Keller A, Graf S, Jaeggi-Groisman SE, Riesen WF *et al.* Beneficial metabolic effects of insulin-like growth factor I in patients with severe insulin-resistant diabetes type A. *Eur J Endocrinol* 1994; **131**: 251–257.
  - 115 Morrow LA, O'Brien MB, Moller DE, Flier JS, Moses AC. Recombinant human insulin-like growth factor-I therapy improves glycemic control and insulin action in the type A syndrome of severe insulin resistance. *J Clin Endocrinol Metab* 1994; **79**: 205–210.
  - 116 Moses AC, Morrow LA, O'Brien M, Moller DE, Flier JS. Insulin-like growth factor I (rhIGF-I) as a therapeutic agent for hyperinsulinemic insulin-resistant diabetes mellitus. *Diabetes Res Clin Pract* 1995; **28**: S185–194.
  - 117 Kuzuya H, Matsuura N, Sakamoto M, Makino H, Sakamoto Y, Kadowaki T *et al.* Trial of insulinlike growth factor I therapy for patients with extreme insulin resistance syndromes. *Diabetes* 1993; **42**: 696–705.
  - 118 Di Cola G, Cool MH, Accili D. Hypoglycemic effect of insulin-like growth factor-1 in mice lacking insulin receptors. *J Clin Invest* 1997; **99**: 2538–2544.
  - 119 Laron Z, Avitzur Y, Klinger B. Carbohydrate metabolism in primary growth hormone resistance (Laron syndrome) before and during insulin-like growth factor-I treatment. *Metabolism* 1995; **44**: 113–118.
  - 120 Laron Z. The essential role of IGF-I: lessons from the long-term study and treatment of children and adults with Laron syndrome. *J Clin Endocrinol Metab* 1999; **84**: 4397–4404.
  - 121 Woods KA, Camacho-Hubner C, Savage MO, Clark AJ. Intrauterine growth retardation and postnatal growth failure associated with deletion of the insulin-like growth factor I gene. *N Eng J Med* 1996; **335**: 1363–1367.
  - 122 Woods KA, Camacho-Hubner C, Bergman RN, Barter D, Clark AJ, Savage MO. Effects of insulin-like growth factor I (IGF-I) therapy on body composition and insulin resistance in IGF-I gene deletion. *J Clin Endocrinol Metab* 2000; **85**: 1407–1411.
  - 123 Jones JL, Clemmons DR. Insulin-like growth factors and their binding proteins: biological actions. *Endocr Rev* 1995; **16**: 3–34.
  - 124 Hwa V, Oh Y, Rosenfeld RG. The insulin-like growth factor-binding protein (IGFBP) superfamily. *Endocr Rev* 1999; **20**: 761–787.
  - 125 Brismar K, Fernqvist-Forbes E, Wahren J, Hall K. Effect of insulin on the hepatic production of insulin-like growth factor-binding protein-1 (IGFBP-1), IGFBP-3, and IGF-I in insulin-dependent diabetes. *J Clin Endocrinol Metab* 1994; **79**: 872–878.
  - 126 Holly JM, Biddlecombe RA, Dunger DB, Edge JA, Amiel SA, Howell R *et al.* Circadian variation of GH-independent IGF-binding protein in diabetes mellitus and its relationship to insulin. A new role for insulin? *Clin Endocrinol (Oxf)* 1988; **29**: 667–675.
  - 127 Batch JA, Baxter RC, Werther G. Abnormal regulation of insulin-like growth factor binding proteins in adolescents with insulin-dependent diabetes. *J Clin Endocrinol Metab* 1991; **73**: 964–968.
  - 128 Lewitt MS, Denyer GS, Cooney GJ, Baxter RC. Insulin-like growth factor-binding protein-1 modulates blood glucose levels. *Endocrinology* 1991; **129**: 2254–2256.
  - 129 Baxter RC, Hizuka N, Takano K, Holman SR, Asakawa K. Responses of insulin-like growth factor binding protein-1 (IGFBP-1) and the IGFBP-3 complex to administration of insulin-like growth factor-I. *Acta Endocrinol (Copenh)* 1993; **128**: 101–108.
  - 130 Frystyk J, Hussain M, Skjaerbaek C, Schmitz O, Christiansen JS, Froesch ER *et al.* Serum free IGF-I during a hyperinsulinemic clamp following 3 days of administration of IGF-I vs. saline. *Am J Physiol* 1997; **273**: E507–513.
  - 131 Holt RIG, Jones JS, Stone NM, Baker AJ, Miell JP. Sequential changes in insulin-like growth factor I (IGF-I) and IGF-binding proteins in children with end-stage liver disease before and after successful orthotopic liver transplantation. *J Clin Endocrinol Metab* 1996; **81**: 160–168.
  - 132 Holt RIG, Miell JP, Jones JS, Mieli-Vergani G, Baker AJ. Nasogastric feeding enhances nutritional status in paediatric liver disease but does not alter circulating levels of IGF-I and IGF binding proteins. *Clin Endocrinol (Oxf)* 2000; **52**: 217–224.
  - 133 Rajkumar K, Krsek M, Dheen ST, Murphy LJ. Impaired glucose homeostasis in insulin-like growth factor binding protein-1 transgenic mice. *J Clin Invest* 1996; **98**: 1818–1825.
  - 134 Crossey PA, Jones JS, Miell JP. Dysregulation of the insulin/IGF binding protein-1 axis in transgenic mice is associated with hyperinsulinemia and glucose intolerance. *Diabetes* 2000; **49**: 457–465.
  - 135 Rajkumar K, Dheen ST, Murphy LJ. Hyperglycemia and impaired glucose tolerance in IGF binding protein-1 transgenic mice. *Am J Physiol* 1996; **270**: E565–571.
  - 136 Rajkumar K, Murphy LJ. Enhanced gluconeogenesis and hepatic insulin resistance in insulin-like growth factor binding protein-1 transgenic mice. *Biochim Biophys Acta* 1999; **1426**: 491–497.
  - 137 Hoefflich A, Wu M, Mohan S, Foll J, Wanke R, Froehlich T *et al.* Overexpression of insulin-like growth factor-binding protein-2 in transgenic mice reduces postnatal body weight gain. *Endocrinology* 1999; **140**: 5488–5496.
  - 138 Zapf J. Role of insulin-like growth factor (IGF) II and IGF binding proteins in extrapancreatic tumour hypoglycaemia. *J Int Med* 1993; **234**: 543–552.
  - 139 Joffe B, Kew M, Beaton G, Kusman B, Seftel H. Serum somatomedin and insulin levels in tumor hypoglycemia. *J Endocrinol Invest* 1978; **1**: 269–271.

- 140 Baxter RC, Daughaday WH. Impaired formation of the ternary insulin-like growth factor-binding protein complex in patients with hypoglycemia due to nonislet cell tumors. *J Clin Endocrinol Metab* 1991; **73**: 696–702.
- 141 Daughaday WH, Trivedi B, Baxter RC. Serum ‘big insulin-like growth factor II’ from patients with tumor hypoglycemia lacks normal E-domain O-linked glycosylation, a possible determinant of normal propeptide processing. *Proc Natl Acad Sci USA* 1993; **90**: 5823–5827.
- 142 Reeve AE, Eccles MR, Wilkins RJ, Bell GI, Millow LJ. Expression of insulin-like growth factor-II transcripts in Wilms’ tumour. *Nature* 1985; **317**: 258–260.
- 143 Ron D, Powers AC, Pandian MR, Godine JE, Axelrod L. Increased insulin-like growth factor II production and consequent suppression of growth hormone secretion: a dual mechanism for tumor-induced hypoglycemia. *J Clin Endocrinol Metab* 1989; **68**: 701–706.
- 144 Zapf J, Futo E, Peter M, Froesch ER. Can ‘big’ insulin-like growth factor II in serum of tumor patients account for the development of extrapancreatic tumor hypoglycemia? *J Clin Invest* 1992; **90**: 2574–2584.
- 145 Katz LE, Liu F, Baker B, Agus MS, Nunn SE, Hintz RL *et al.* The effect of growth hormone treatment on the insulin-like growth factor axis in a child with nonislet cell tumor hypoglycemia. *J Clin Endocrinol Metab* 1996; **81**: 1141–1146.
- 146 Daughaday WH, Emanuele MA, Brooks MH, Barbato AL, Kapadia M, Rotwein P. Synthesis and secretion of insulin-like growth factor II by a leiomyosarcoma with associated hypoglycemia. *N Eng J Med* 1988; **319**: 1434–1440.
- 147 Frystyk J, Skjaerbaek C, Zapf J, Orskov H. Increased levels of circulating free insulin-like growth factors in patients with non-islet cell tumour hypoglycaemia. *Diabetologia* 1998; **41**: 589–594.
- 148 Teale JD, Blum WF, Marks V. Alleviation of non-islet cell tumour hypoglycaemia by growth hormone therapy is associated with changes in IGF binding protein-3. *Ann Clin Biochem* 1992; **29**: 314–323.