### Final exam prep

Tutorial 10

### **Tips**

- Very similar in structure to midterm
- Take your time, but not too long
- Nearly every included is intentional
- Thought process needs to be clear
  - Can use bullet points, may help
- Practice using past exams
- Use short hand for describe questions

### **Tips**

- Include source of data
  - (eg. "It was higher (Table 1)/(\$1:T1)")
- Answer the question
  - (if question says use all data, then use all data!)
- Mechanism questions: Include all relevent info/data before synthesizing data into a mechanism
  - Example in slides to follow as this was an area that many had difficulty with

### **Example answer for mechanism question**

The question (from 2009 midterm):

Propose a mechanism to explain how the different diet and endurance exercise treatments altered the metabolic parameters presented in Table 2. Use only data from Studies 1 and/or 2 to defend your answer.

### **Example answer for mechanism question**

The question (from 2009 midterm):

Propose a mechanism to explain how the different diet and endurance exercise treatments altered the metabolic parameters presented in Table 2. Use only data from Studies 1 and/or 2 to defend your answer.

#### Key:

- only data from \$1 + \$2
- Diet + exercise + metabolism in table 2

### **Example answer for mechanism question**

The question (from 2009 midterm):

Propose a mechanism to explain how the different diet and endurance exercise treatments altered the metabolic parameters presented in Table 2. Use only data from Studies 1 and/or 2 to defend your answer.

#### Key:

- only data from \$1 + \$2
- Diet + exercise + metabolism in table 2

Let's go back to what \$1 + \$2 say (will need background too)

### **Background**

Exercise-induced contractions in skeletal muscle results in increased delivery of carbohydrate (CHO; glucose) into the muscle to maintain metabolism.

This is likely through the regulation of glucose transporters (GLUT), particularly GLUT4, which is the principal isoform in skeletal muscle.

### **Background**

Exercise-induced contractions in skeletal muscle results in increased delivery of carbohydrate (CHO; glucose) into the muscle to maintain metabolism.

This is likely through the regulation of glucose transporters (GLUT), particularly GLUT4, which is the principal isoform in skeletal muscle.

#### Key:

Exercise -> GLUT4 -> CHO into muscles

#### \$1: T1


**Table 1.** Blood glucose concentrations before and during consumption of the meal replacement drink (expressed as glucose area under the curve (gAUC)) during (E1, E3) or after (R1, R2, R3) exercise training.

Treatment Day	Blood glucose (mmol/L)		
	0min	$gAUC_{0\text{-}120min}$	
E1	$8.84 \pm 0.32^a$	541.3 ± 25.9 <sup>a</sup>	
E3	$7.26 \pm 0.32^{b}$	482.1 ± 23.3 <sup>b</sup>	
R1	$7.34 \pm 0.25^{b}$	489.3 ± 24.7 <sup>b</sup>	
R2	8.20 ± 0.19°	589.4 ± 29.3°	
R3	8.41 ± 0.14°	600.1 ± 23.1°	

 $<sup>^{</sup>abc}Data\,(mean\,\pm\,SEM)$  in columns with different superscripts are significantly different from each other at p<0.05.

#### Figure 1:

#### \$1: F1






Figure 1. Data presented are means ± SEM.

- A. The % change in total GLUT4 protein during the exercise regime. Values for E1 were arbitrarily set at 100 and all other days' values were expressed relative to E1.
- B. Total glucose concentrations in quadriceps muscle after consumption of the meal replacement drink.
- C. Total glycogen concentrations in quadriceps muscle after consumption of the meal replacement drink.
- <sup>abc</sup>Bars with different letters are significantly different from each other at p<0.05.</p>

Take the time to think how to answer it simply and concisely. Combine results together to be more efficient and cleaner.

Take the time to think how to answer it simply and concisely. Combine results together to be more efficient and cleaner.

#### **T1**

Fasting glucose + glucose AUC: E1 > (R2 = R3) > (E2 = R1)

Take the time to think how to answer it simply and concisely. Combine results together to be more efficient and cleaner.

#### **T1**

Fasting glucose + glucose AUC: E1 > (R2 = R3) > (E2 = R1)

#### F1

- GLUT4: R3 < (E1 = R2) < (E3 = R1)
- Muscle glucose + glycogen: (E1 = E2 = R1) > (R2 = R3)

### **S2: T2**

Table 2. Subject characteristics before and after 3 months of exercise training.

Measure	Baseline		Post-training	
	LF	HF	LF	HF
Fasting glucose (mmol/L)	9.2 ± 0.2 <sup>a</sup>	9.4 ± 0.1 <sup>a</sup>	8.5 ± 0.2 <sup>b</sup>	9.5 ± 0.1 <sup>a</sup>
HbA1c (%)	$7.3 \pm 0.3^{a}$	$7.5 \pm 0.4^{a}$	$6.9 \pm 0.1^{b}$	$7.4 \pm 0.1^{a}$
Triglycerides(mmol/L)	$2.5 \pm 0.1^{a}$	$2.7 \pm 0.2^{a}$	$2.0 \pm 0.1^{b}$	$2.2 \pm 0.2^{b}$
Total Cholesterol(mmol/L)	$5.2 \pm 0.2^{a}$	$5.3 \pm 0.1^{a}$	$4.8 \pm 0.1^{b}$	$5.0 \pm 0.1^{b}$

Data presented are means  $\pm$  SEM.  $^{abc}$ Data with different letters are significantly different from each other at p<0.05.

Figure 3:

#### S2: F2




Figure 2. Whole-body lipid oxidation at the end of three months of endurance exercise. Data presented are means  $\pm$  SEM. <sup>abc</sup>Data with different letters in the same workload are significantly different from each other at p<0.05.

Again, take the time to think how to answer it simply and concisely.

Again, take the time to think how to answer it simply and concisely.

#### **T2**

- Fasting glucose + HbA1c: (Base LF = Base HF = Post HF)
   Post LF
- TAG + Cholesterol: (Base Lf = Base HF) > (Post LF = Post HF)

Again, take the time to think how to answer it simply and concisely.

#### **T2**

- Fasting glucose + HbA1c: (Base LF = Base HF = Post HF)
   > Post LF
- TAG + Cholesterol: (Base Lf = Base HF) > (Post LF = Post HF)

#### **F2**

Lipid Oxid. at 60, 90, 120, 150 Watts: (Base LF = Base HF)
 < (Post LF = Post HF)</li>

### Let's go back to the question:

Propose a mechanism to explain how the different diet and endurance exercise treatments altered the metabolic parameters presented in Table 2. Use only data from Studies 1 and/or 2 to defend your answer.

Include the background, \$1, and \$2 info first thing in the answer! It may be redundant, but this is good for you + us!

• Background: Exercise -> GLUT4 -> CHO into muscles

- Background: Exercise -> GLUT4 -> CHO into muscles
- S1, T1:

- Background: Exercise -> GLUT4 -> CHO into muscles
- S1, T1:
  - Exercise reduced fasting glucose + glucose AUC acutely, but starting going back to normal after several rests days (E1 > (R2 = R3) > (E2 = R1))

- Background: Exercise -> GLUT4 -> CHO into muscles
- S1, T1:
  - Exercise reduced fasting glucose + glucose AUC acutely, but starting going back to normal after several rests days (E1 > (R2 = R3) > (E2 = R1))
- S1, F1:

- Background: Exercise -> GLUT4 -> CHO into muscles
- S1, T1:
  - Exercise reduced fasting glucose + glucose AUC acutely, but starting going back to normal after several rests days (E1 > (R2 = R3) > (E2 = R1))
- S1, F1:
  - Exercise acutely increased GLUT4, but started dropping right after a single rest day (R3 < (E1 = R2) < (E3 = R1))</li>

- Background: Exercise -> GLUT4 -> CHO into muscles
- S1, T1:
  - Exercise reduced fasting glucose + glucose AUC acutely, but starting going back to normal after several rests days (E1 > (R2 = R3) > (E2 = R1))
- S1, F1:
  - Exercise acutely increased GLUT4, but started dropping right after a single rest day (R3 < (E1 = R2) < (E3 = R1))</li>
  - Exercise acutely increased muscle glucose + glycogen, but dropped after many rest days ((E1 = E2 = R1) > (R2 = R3))

• S2, T2:

- S2, T2:
  - Only LF diet after exercise had lower fasting glucose + HbA1c ((Base LF = Base HF = Post HF) > Post LF)

- S2, T2:
  - Only LF diet after exercise had lower fasting glucose + HbA1c ((Base LF = Base HF = Post HF) > Post LF)
  - Exercise, regardless of diet, reduced TAG + Cholesterol ((Base Lf = Base HF) > (Post LF = Post HF))

- S2, T2:
  - Only LF diet after exercise had lower fasting glucose + HbA1c ((Base LF = Base HF = Post HF) > Post LF)
  - Exercise, regardless of diet, reduced TAG + Cholesterol ((Base Lf = Base HF) > (Post LF = Post HF))
- S2, F2:

- S2, T2:
  - Only LF diet after exercise had lower fasting glucose + HbA1c ((Base LF = Base HF = Post HF) > Post LF)
  - Exercise, regardless of diet, reduced TAG + Cholesterol ((Base Lf = Base HF) > (Post LF = Post HF))
- S2, F2:
  - Exercise, regardless of diet, increased lipid oxidation at 60, 90, 120, 150 Watts ((Base LF = Base HF) < (Post LF = Post HF))

- S2, T2:
  - Only LF diet after exercise had lower fasting glucose + HbA1c ((Base LF = Base HF = Post HF) > Post LF)
  - Exercise, regardless of diet, reduced TAG + Cholesterol ((Base Lf = Base HF) > (Post LF = Post HF))
- S2, F2:
  - Exercise, regardless of diet, increased lipid oxidation at 60, 90, 120, 150 Watts ((Base LF = Base HF) < (Post LF = Post HF))

- S2, T2:
  - Only LF diet after exercise had lower fasting glucose + HbA1c ((Base LF = Base HF = Post HF) > Post LF)
  - Exercise, regardless of diet, reduced TAG + Cholesterol ((Base Lf = Base HF) > (Post LF = Post HF))
- S2, F2:
  - Exercise, regardless of diet, increased lipid oxidation at 60, 90, 120, 150 Watts ((Base LF = Base HF) < (Post LF = Post HF))

Then synthesize the answers...

Explain how the different diet and endurance exercise treatments altered the metabolic parameters presented in Table 2.

Explain how the different diet and endurance exercise treatments altered the metabolic parameters presented in Table 2.

Synthesis:

Explain how the different diet and endurance exercise treatments altered the metabolic parameters presented in Table 2.

#### Synthesis:

Acute exercise reduces blood glucose (\$1:T1), because
of the increase in GLUT4 (\$1:F1; Background), which
also lead to an increase in intramuscular glucose and
glycogen (\$1:F1; Background).

Explain how the different diet and endurance exercise treatments altered the metabolic parameters presented in Table 2.

#### Synthesis:

- Acute exercise reduces blood glucose (\$1:T1), because
  of the increase in GLUT4 (\$1:F1; Background), which
  also lead to an increase in intramuscular glucose and
  glycogen (\$1:F1; Background).
- However, if exercise is not consistent enough, these gains are diminished (S1:T1 + F1)

• Synthesis, con't

- Synthesis, con't
  - The HF diet seems not to influence metabolism much during exercise as both groups decreased TAG + cholesterol equally (S2:T2) and lipid oxidation was equal between diets (S2:F2)

- Synthesis, con't
  - The HF diet seems not to influence metabolism much during exercise as both groups decreased TAG + cholesterol equally (S2:T2) and lipid oxidation was equal between diets (S2:F2)
  - The improvements in glucose + HbA1c in the LF diet in S2:T2 may be due to the greater intake of CHO, letting the muscles use more glucose

- Synthesis, con't
  - The HF diet seems not to influence metabolism much during exercise as both groups decreased TAG + cholesterol equally (\$2:T2) and lipid oxidation was equal between diets (\$2:F2)
  - The improvements in glucose + HbA1c in the LF diet in S2:T2 may be due to the greater intake of CHO, letting the muscles use more glucose
  - But, lipid oxidation is the same between diets, so it could be that the LF diet may increase GLUT4 expression to use the dietary CHO, reducing blood glucose (as seen in \$1:T1) by storing it in the muscles (Background)... However, GLUT4 was not measured in \$2 so this is speculation.