



Modelling competition for nutrients between QFA spots


Daniel Boocock
25/07/2016

QFA

- Microbial cultures are inoculated in a grid on a solid agar surface.
 - Repeatedly photographed.
 - Analysed to get cell density estimates.
 - Growth curves are fit with the logistic model.
-
- Will modelling competition make fitness estimates more reliable?








QFA uses fits of the logistic model


- From this fitness estimates are derived.
- MDP is the number of doublings undergone by the culture population from the inoculum density (g) to carrying capacity (K) throughout the experiment.
- MDR is the inverse of the doubling time at the beginning of the experiment.
- We can obtain the same shape curve using a mechanistic model.



$$\text{rate} = b_i[N][C]$$

- Mass action kinetic logistic equivalent model.
- Assumes all cultures would reach the same final cell density if starting with the same nutrient density.

$$\dot{x} = rx \left(1 - \frac{x}{K}\right) \quad x(t) = \frac{KPe^{rt}}{K + P(e^{rt} - 1)}$$




- Replace with my own plots

$$r = b_i(N_0 + C_0)$$

$$K = (N_0 + C_0)$$

Can we improve the reliability of fitness estimates by modelling competition?



C_i – Cells

N_i – Nutrients

- ← Put correction plot here.

- We use a network diffusion model and mass action kinetics.


$$\frac{dC_i}{dt} = b_i N_i C_i,$$

$$\frac{dN_i}{dt} = -b_i N_i C_i - k_n \sum_{j \in \delta_i} (N_i - N_j)$$


- C_0 , N_0 , and k_n are fit at the plate level.
- b is fit on a culture level.
- If we set $k_n = 0$ we return to the logistic equivalent model.
- In order to fit the logistic equivalent model we have to allow N_0 to vary at the culture level.
- Cell arrest or signalling could also be effects.

We study the dataset p15


- *cdc13-1* is background mutation
- *CDC13* is a gene important for telomere stability
- ~50 deletions chosen for relevance to telomere biology
- Why did we choose this data set?
 - We are interested in detecting competition effects
 - Large variation in fitness of strains (suppress or enhance defect associated with *cdc13-1*).
 - Lots of replicates.
 - Hopefully our results will be interpretable by this audience.



Best Competition Model BC Fit to p15




- Each fit takes ~ 3 hours.
- We cannot yet find global minima. Confounding effect between b and C_0 . (thoughts about fitting approaches/measuring C_0)
- However, ranks of b are correlated for the best fits.




How can we asses the Competition Model?

- For the competition mode, b rank is equal to MDR*MDP rank.
 - Do the rankings on the left match your expectations?
 - Compare variances
 - Are fitness estimates more reliable? (Yet to do this. Will by Monday.)
 - Compare between plates using miniQFA.
(Might be able to do by Monday)



A quick preview of parameter variance.

All rankings are for b.



Further Work

- We still need to work on finding global minima.
- We can decide whether it is worth doing so now.
- Could experiments start to measure C_0 ?
- b as a fitness estimate? Depends on predictions.
- Where can I find references for validation based on existing knowledge?