

Set 3 - Diffusion and OpenMP

Issued: October 10, 2014

Hand in: October 17, 2014, 8:00am

Question 1: Diffusion in 2D

Heat flow in a medium can be described by the diffusion equation

$$\frac{\partial \rho(\mathbf{r}, t)}{\partial t} = D \nabla^2 \rho(\mathbf{r}, t), \quad (1)$$

where $\rho(\mathbf{r}, t)$ is a measure for the amount of heat at position \mathbf{r} and time t and the diffusion coefficient D is constant.

Lets define the domain Ω in two dimensions as $\{x, y\} \in [-1, 1]^2$. Equation 1 then becomes

$$\frac{\partial \rho(x, y, t)}{\partial t} = D \left(\frac{\partial^2 \rho(x, y, t)}{\partial x^2} + \frac{\partial^2 \rho(x, y, t)}{\partial y^2} \right). \quad (2)$$

Equation 2 is discretized with a central finite difference scheme in space and explicit Euler in time:

$$\frac{\rho_{i,j}^n}{\delta t} = D \left(\frac{\rho_{i-1,j}^n - 2\rho_{i,j}^n + \rho_{i+1,j}^n}{\delta x^2} + \frac{\rho_{i,j-1}^n - 2\rho_{i,j}^n + \rho_{i,j+1}^n}{\delta y^2} \right) \quad (3)$$

where $\rho_{i,j}^n = \rho(-1 + i\delta x, -1 + j\delta y, n\delta t)$ and $\delta x = \frac{2}{N}$, $\delta y = \frac{2}{M}$ for a domain discretized with $N \times M$ gridpoints.

We will use Dirichlet boundary conditions

$$\rho(-1, y, t) = \rho(x, -1, t) = \rho(1, y, t) = \rho(x, 1, t) = 0 \quad \forall t \geq 0 \quad (4)$$

and an initial heat distribution

$$\rho(x, y, 0) = \begin{cases} 1 & |x, y| < 1/2 \\ 0 & \text{otherwise} \end{cases}. \quad (5)$$

- Analyze the stability of the scheme given in Equation 3 using the von Neumann stability analysis.
- Parallelize your code from Exercise 1a using OpenMP and make both strong and weak scaling plots up to 24 cores. Define the problem size as the number of grid points in the discretization of Ω . Try different problem sizes for the scaling plots, do they affect the scaling? Try using both the GCC OpenMP and the Intel OpenMP libraries. Can you see a difference?

The total amount of heat in the system $N(t)$ and the second moment of the cloud $\mu^2(t)$ at a time t can be computed as

$$N(t) = \int_{\Omega} dx dy \rho(x, y, t), \quad \mu^2(t) = \int_{\Omega} dx dy \rho(x, y, t)(x^2 + y^2). \quad (6)$$

- c) Make a plot of the time evolution of $N(t)$ and $\mu^2(t)$ for $D = 1, 2, 5$ and $t \in [0, 10]$. What is the effect of the boundary? What is the effect of D ?

Summary

Summarize your answers, results and plots into a PDF document. Furthermore, elucidate the main structure of the code and report possible code details that are relevant in terms of accuracy or performance. Send the PDF document and source code to your assigned teaching assistant.