

MATH 239 Introduction to Combinatorics

University of Waterloo - Fall 2025

2nd December 2025

Instructor: Douglas Stebila

L^AT_EX: Xing Liu

Contents

1	Principles of Enumeration	4
1.1	AND vs. OR - Products vs. Sums	4
1.2	Lists, Permutations, and Subsets	4
1.3	Multisets	7
1.4	Bijections	9
1.5	Summary of Combinatorial and Bijective Proofs	11
2	Generating Series	12
2.1	Binomial Theorem	13
2.2	Generating Series	17
2.3	Sum and Product Lemmas	20
2.4	Compositions	25
3	Binary Strings	30
3.1	Regular Expressions	31
3.2	Unambiguous Expressions	33
3.3	Translation into Generating Series	35
3.4	Block Decomposition	36
3.5	Recursive Decompositions	37
3.6	Excluded Substrings	39

4	Partial Fractions and Recurrence Relations	41
4.1	Rational Function to Explicit Coefficient Formula Using Partial Fractions	41
4.2	Recurrence Relation to Explicit Coefficient Formula	43
4.3	Convenient Conversion Theorems	45
5	Introduction to Graph Theory	47
5.1	Isomorphism	49
5.2	Degree	50
5.3	Bipartite Graphs	52
5.4	How to Specify a Graph	54
5.5	Subgraphs, Paths, and Cycles	56
5.6	Connectedness	60
5.7	Eulerian Circuits	63
5.8	Bridges	64
6	Trees	66
6.1	Trees	66
6.2	Spanning Trees	69
6.3	Characterizing Bipartite Graphs	70
7	Planar Graphs	72
7.1	Planarity	72
7.2	Euler's Formula	75
7.3	Platonic Solids	77
7.4	Nonplanar Graphs	77
7.5	Kuratowski's Theorem	80
7.6	Colouring	81
7.7	Dual Planar Maps	85
8	Matchings	87
8.1	Matching	87
8.2	Covers	88
8.3	König's Theorem	89
8.4	Hall's Theorem	92
8.5	Perfect Matchings in Bipartite Graphs	94

8.6	Edge Colourings	94
-----	---------------------------	----

1 Principles of Enumeration

Lecture 1, 2025/09/03

1.1 AND vs. OR - Products vs. Sums

Example. Let $A = \{\text{apple, orange, banana, grape}\}$ and $B = \{\text{carrot, lettuce, onion}\}$. The # of ways of choosing a fruit from A **AND** a vegetable from B is $|A| \cdot |B| = 4 \cdot 3 = 12$. This is equivalent to choosing one element from the cartesian product of A and B :

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}.$$

For finite sets, the cardinalities satisfy

$$|A \times B| = |A| \cdot |B|.$$

The # of ways of choosing a fruit from A **OR** a vegetable from B is $|A| + |B| = 4 + 3 = 7$. This is equivalent to choosing one element from the union of A and B :

$$A \cup B = \{x : x \in A \text{ or } x \in B\}.$$

For disjoint sets,

$$|A \cup B| = |A| + |B|$$

Note that not all sets are disjoint, some have non-trivial intersection.

The intersection is $A \cap B = \{x : x \in A \text{ and } x \in B\}$. For finite sets,

$$|A \cup B| + |A \cap B| = |A| + |B|.$$

1.2 Lists, Permutations, and Subsets

Definition 1.1 (List).

A **list** of a set S is a listing of the elements of S exactly once, in some order.

Definition 1.2 (Permutation).

A **permutation** is a list of the set $\{1, 2, \dots, n\}$ for some $n \in \mathbb{N}$.

Example. $S = \{a, b, c\}$ has 6 lists:

$$abc \quad acb \quad bac \quad bca \quad cab \quad cba.$$

We can construct a list of S by choosing an element $v \in S$ to be the first element of the list, and then append a list of $S \setminus \{v\}$. If p_n is the # of lists of a set of size n , for $n \in \mathbb{N}$, then

$$p_n = n \cdot p_{n-1}$$

and $p_1 = 1$. So by induction we have the following theorem.

Theorem 1.1. For every $n \geq 1$, the # of lists of an n -element set is $n \cdot (n-1) \cdots 2 \cdot 1 = n!$.

Note. By convention, we take $0! = 1$.

Definition 1.3 (Subset).

A **subset** of S is a set containing some (perhaps none, or perhaps all) of the elements of S , at most once each, and in no particular order.

Note. There are 2^n subsets of an n -element set.

To specify: a subset X of S , for each element $v \in S$, we have to decide whether v is in X or not in X . Equivalently, we have to choose {in, out} n times. So,

$$\underbrace{|\{\text{in, out}\}| \cdot |\{\text{in, out}\}| \cdot \dots \cdot |\{\text{in, out}\}|}_{n \text{ times}} = 2^n.$$

Lecture 2, 2025/09/05

Definition 1.4 (Partial List).

A **partial list** of a set S is a list of a subset of S .

We are interested in k -element partial lists of an n -element set, in other words, k -tuples $(s_1, \dots, s_k)$ of distinct elements of S . There are n choices for s_1 , $n - 1$ choices for s_2 , ..., $(n - k + 1)$ choices for s_k . So we have the following.

Theorem 1.2. For $k, n \geq 0$, the # of partial lists of length k of an n -element set is

$$n(n-1)(n-2) \cdots (n-k+1).$$

Note. For $0 \leq k \leq n$, we can write

$$n(n-1)(n-2) \cdots (n-k+1) = \frac{n(n-1) \cdots 1}{(n-k)(n-k-1) \cdots 1} = \frac{n!}{(n-k)!}.$$

Definition 1.5. For $n, k \geq 0$, let $\binom{n}{k}$ (read “ n choose k ”) denote the # of k -element subsets of an n -element set.

Example.

- $\binom{n}{0} = 1$.
- $\binom{n}{n} = 1$ for all n .
- $\binom{n}{1} = n$ for all n .
- $\binom{n}{k} = 0$ if $k > n$.

Theorem 1.3. For $0 \leq k \leq n$, we have

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

Proof. Let S be a set of size n . From Theorem 1.2, the # of partial lists of S of length k is $\frac{n!}{(n-k)!}$. We will choose a k -element subset X of S AND choose a list of X , which will give a list of a subset of S of length k . Thus, the # of length- k partial lists of S is

$$(\# \text{ of } k\text{-element subsets of } S) \cdot (\# \text{ of lists of a } k\text{-element set}) = \binom{n}{k} \cdot k!.$$

$$\text{So, } \binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

□

Note. This is an example of a combinatorial proof, where we prove an identity by counting the same set in two different ways.

Example. For $n \geq 0$,

$$\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n} = 2^n.$$

Prove this by counting sets, not by algebraic manipulation.

Proof. The RHS, 2^n , is the # of subsets of an n -element set. On the LHS, for $0 \leq k \leq n$, $\binom{n}{k}$ is the # of k -element subsets of an n -element set. And addition corresponds to “OR”, so the LHS is the # of ways of choosing a single subset of an n -element set. That is, choosing one of the $\binom{n}{0}$ 0-element subsets, or one of the $\binom{n}{1}$ 1-element subsets, or ..., or one of the $\binom{n}{n}$ n -element subsets. $\square$

Example. For $k, n \geq 1$,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Proof. The LHS, $\binom{n}{k}$, is the # of k -element subsets of an n -element set S . WLOG, assume $S = \{1, 2, \dots, n\}$. Notice that $\binom{n-1}{k}$ is the # of k -element subsets of $S' = \{1, 2, \dots, n-1\}$, and $\binom{n-1}{k-1}$ is the # of $(k-1)$ -element subsets of S' . Recall our intuition that addition corresponds to “OR”. Any k -element subset of S either includes n or does not include n . If $n \notin X$, then X is a subset of size k of S' . If $n \in X$, then $X \setminus \{n\}$ is a subset of size $k-1$ of S' . Thus, LHS = RHS. $\square$

1.3 Multisets

Suppose I have a bag containing 8 marbles, each of which is either red, green, or blue. How many different bag contents are possible?

First question: how can I represent the contents of a bag? For example, we could have 2 red, 3 green, and 3 blue marbles.

Lecture 3, 2025/09/8

- We might want to write it as a set $\{R, R, G, G, G, B, B, B\}$, but this is not a set since sets do not have duplicates.
- Invent new notation that preserves multiplicity:

$$[[R, R, G, G, G, B, B, B]] \text{ or } \langle\langle R^2, G^3, B^3 \rangle\rangle.$$

- Keep track of multiplicities in a vector. Fix an order, say, R then G then B , and use a tuple or a vector to record the multiplicities: $(2, 3, 3)$.

To count the # of bags containing 8 marbles each of which is either red, green, or blue, we need to count the # of 3-tuples (n_1, n_2, n_3) of integers s.t. $n_1 + n_2 + n_3 = 8$ and $n_1, n_2, n_3 \geq 0$.

Definition 1.6 (Multiset).

Let $n \geq 0, t \geq 1$ be integers. A **multiset** of size n with elements of t types is a sequence of non-negative integers $m_1, m_2, \dots, m_t$ s.t. $m_1 + m_2 + \dots + m_t = n$. The set of all multisets of size n with elements of t -types is denoted $\mathcal{M}(n, t)$.

Theorem 1.4. For $n \geq 0, t \geq 1$, the # of n -element multisets with elements of t types is

$$|\mathcal{M}(n, t)| = \binom{n+t-1}{t-1}.$$

Proof. We know that $\binom{n+t-1}{t-1}$ is the # of $(t-1)$ -element subsets of an $(n+t-1)$ -element set. We will show how to translate between $(t-1)$ -element subsets of an $(n+t-1)$ -element set and multisets of size n with t types. Let us work down a row of $n+t-1$ circles (e.g. $n=8, t=3$):

○ ○ ○ ○ ○ ○ ○ ○ ○ ○

Now, let us choose a $(t-1)$ -element subset of these and cross them out

○ ○ × ○ ○ ○ × ○ ○ ○

Our row now has n circles grouped into t segments each containing 0 or more consecutive circles. Let m_i be the length of the i^{th} segment of consecutive circles, so $m_1 + m_2 + \dots + m_t = n$. So $(m_1, m_2, \dots, m_t)$ is a multiset of size n with elements of t types.

Now, let us go the other way. Let $(m_1, m_2, \dots, m_t)$ be a multiset of size n with t types. Write down m_1 circles, then an $\times$, then m_2 circles, then an $\times$, and so on, upto m_t circles. We will have written down $n+t-1$ symbols total, of which $t-1$ are $\times$'s, these indicate the $(t-1)$ -element subset of an $(n+t-1)$ -element set. □

Note. This is an example of a bijective proof.

1.4 Bijections

Definition 1.7 (Surjective, Injective, Bijective).

Let A and B be sets. Let $f : A \rightarrow B$.

- f is **surjective** (or **onto**) if, for every $b \in B$, $\exists a \in A$ s.t. $f(a) = b$.
- f is **injective** (or **one-to-one**) if, for every $a, a' \in A$, if $f(a) = f(a')$, then $a = a'$.
- f is **bijective** if it is both surjective and injective.

Definition 1.8 (Well-defined).

An expression yields a **well-defined** function $f : A \rightarrow B$ if, for every input $a \in A$, $f(a)$ assigns a single unambiguous value $b \in B$.

Proposition 1.5. A function $f : A \rightarrow B$ is a bijection $\iff f$ has an inverse, namely if $\exists$ a well-defined function $g : B \rightarrow A$ s.t. $g(f(a)) = a$ for all $a \in A$ and $f(g(b)) = b$ for all $b \in B$.

Remark (Notation).

If $\exists$ a bijection between sets A and B , we can write $A \rightleftharpoons B$.

Corollary 1.6. If $A \rightleftharpoons B$ and at least one of A or B is finite, then $|A| = |B|$.

Example. For $k, n \geq 1$,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Proof. Let $S = \{1, 2, \dots, n\}$ and $S' = \{1, 2, \dots, n-1\}$. Let $A = \{k\text{-element subsets of } S\}$ and $B = \{(k-1)\text{-element or } k\text{-element subsets of } S'\}$. By definition, $|A| = \binom{n}{k}$ and we can easily see that $|B| = \binom{n-1}{k-1} + \binom{n-1}{k}$. The goal is to prove $A \rightleftharpoons B$. Define $f : A \rightarrow B$ by

$$f(u) = \begin{cases} u, & \text{if } n \notin u \\ u \setminus \{n\}, & \text{if } n \in u \end{cases}$$

Lecture 4, 2025/09/10

Question: Is f well-defined, i.e. $\forall u \in A$, is $f(u) \in B$?

Answer: Yes, $f(u)$ does not contain n (so it's a subset of S') and has size either $k-1$ or k . So

$f(u) \in B$.

Define $g : B \rightarrow A$ by

$$g(v) = \begin{cases} v, & \text{if } |v| = k \\ v \cup \{n\}, & \text{if } |v| = k - 1 \end{cases}$$

Claim (1). For all $u \in A$, $g(f(u)) = u$.

Proof of Claim (1). First consider $u \in A$ s.t. $n \notin u$. Then, $f(u) = u$ and $|f(u)| = k$, so $g(f(u)) = u$ (first branch of g). Now consider $u \in A$ s.t. $n \in u$. Then, $f(u) = u \setminus \{n\}$ and $|f(u)| = k - 1$, so $g(f(u)) = (u \setminus \{n\}) \cup \{n\} = u$ (second branch of g). $\square$

Claim (2). For all $v \in B$, $f(g(v)) = v$.

Proof of Claim (2). Exercise. $\square$

Thus, f is a bijection, so $|A| = |B|$, i.e. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$. $\square$

Remark (Problem Solving Tips: what steps are needed for a bijective argument?).


To show a function $f : A \rightarrow B$ is a bijection by giving an inverse:

- (1) State the function $f : A \rightarrow B$.
- (2) Show that f is well-defined, i.e. $\forall a \in A, f(a) \in B$.
- (3) State the function $g : B \rightarrow A$.
- (4) Show that g is well-defined.
- (5) Show that $g(f(a)) = a$ for all $a \in A$.
- (6) Show that $f(g(b)) = b$ for all $b \in B$.


Exercise: Show that $\binom{n}{k} = \binom{n}{n-k}$ for $n \geq k \geq 0$ using a combinatorial or bijective argument.

1.5 Summary of Combinatorial and Bijective Proofs

In a combinatorial proof, describe two different ways of counting the size of the same set.



In a bijective proof, describe two different sets and show they're in bijection.



- (1) Give sets A, B and count them.
- (2) Show a bijection between A and B .
- (3) This implies $|A| = |B|$.

2 Generating Series

Definition 2.1 (Formal Power Series).

A **formal power series** is an expression of the form

$$G(x) = \sum_{n=0}^{\infty} g_n x^n$$

in which the coefficients $(g_0, g_1, g_2, \dots)$ are a sequence of real numbers.

Note. We use x as an **indeterminate** for which we do not normally substitute in particular values. This means that we never worry about the convergence of the series, all we require is that the coefficients are finite real numbers.

Example. $G(x) = 0 - 1x + 2x^2 - 3x^3 + \dots = \sum_{n=0}^{\infty} (-1)^n n x^n$.

We can add and multiply formal power series like polynomials. Let

$$F(x) = \sum_{n \geq 0} f_n x^n = f_0 + f_1 x + f_2 x^2 + \dots,$$
$$G(x) = \sum_{n \geq 0} g_n x^n = g_0 + g_1 x + g_2 x^2 + \dots.$$

Then,

$$(F + G)(x) = \sum_{n \geq 0} (f_n + g_n) x^n = (f_0 + g_0) + (f_1 + g_1)x + (f_2 + g_2)x^2 + \dots,$$
$$(F \cdot G)(x) = \sum_{n \geq 0} \left(\sum_{k=0}^n f_k g_{n-k} \right) x^n = (f_0 g_0) + (f_0 g_1 + f_1 g_0)x + (f_0 g_2 + f_1 g_1 + f_2 g_0)x^2 + \dots.$$

Lecture 5, 2025/09/12

Example. Note that

$$(1 + 3x^2 + 5x^4 + \dots) + x(2 + 4x^2 + 6x^4 + \dots) = (1 + 3x^2 + 5x^4 + \dots) + (2x + 4x^3 + 6x^5 + \dots)$$
$$= 1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + \dots$$

Equivalently,

$$\begin{aligned} \sum_{n \geq 0} (2n+1)x^{2n} + x \sum_{n \geq 0} (2n+2)x^{2n} &= \sum_{n \geq 0} (2n+1)x^{2n} + \sum_{n \geq 0} (2n+2)x^{2n+1} \\ &= \sum_{n \geq 0} ((2n+1)x^{2n} + (2n+2)x^{2n+1}) \\ &= \sum_{n \geq 0} (n+1)x^n. \end{aligned}$$

Definition 2.2 (Geometric Series).

The **geometric series** is of the form

$$G(x) = 1 + x + x^2 + x^3 + \cdots = \sum_{n \geq 0} x^n.$$

Consider $xG(x) = x + x^2 + x^3 + \cdots = x \sum_{n \geq 0} x^n$. Then,

$$G(x) - xG(x) = 1 \implies G(x)(1-x) = 1 \implies G(x) = \frac{1}{1-x} = (1-x)^{-1}.$$

Definition 2.3 (Inverse of a Formal Power Series).

If $F(x)$ and $G(x)$ are formal power series such that $F(x)G(x) = 1$, then we write $G(x) = F(x)^{-1}$ or $G(x) = \frac{1}{F(x)}$, and say that $G(x)$ is the **inverse** of $F(x)$.

Corollary 2.1. The inverse of $F(x) = \sum_{n \geq 0} f_n x^n$ exists $\iff f_0 \neq 0$.

2.1 Binomial Theorem

Theorem 2.2 (Binomial Theorem).

For any $n \in \{0, 1, 2, \dots\}$,

$$(1+x)^n = \sum_{k=0}^n \binom{n}{k} x^k = \sum_{k \geq 0} \binom{n}{k} x^k.$$

Proof. We will use a combinatorial proof. Let $\mathcal{P}_n$ be the power set of $\{1, 2, \dots, n\}$, i.e. the set of all subsets of $\{1, 2, \dots, n\}$. Let $\mathcal{P}_{n,k}$ be the set of all k -element subsets of $\{1, 2, \dots, n\}$. We saw before that a

bijection between $\mathcal{P}_n$ and $\{0, 1\}^n$, i.e. a subset S of $\{1, 2, \dots, n\}$ corresponds to an “indicator vector” in $\{0, 1\}^n$. Let $S \in \mathcal{P}_n$ be a subset of $\{1, 2, \dots, n\}$. The corresponding indicator vector will be

$$\vec{a} = (a_1, a_2, \dots, a_n) \quad \text{where} \quad a_i = \begin{cases} 0, & \text{if } i \notin S, \\ 1, & \text{if } i \in S. \end{cases}$$

Note that $|S| = \sum_{i=1}^n a_i$. Let’s introduce an indeterminate x . For every subset S with indicator vector $\vec{a}$, it holds that

$$x^{|S|} = x^{a_1 + a_2 + \dots + a_n}.$$

Because of the bijection between $\mathcal{P}_n$ and $\{0, 1\}^n$, summing over all subsets is equivalent to summing over all indicator vectors. Thus,

$$\sum_{S \in \mathcal{P}_n} x^{|S|} = \sum_{\vec{a} \in \{0, 1\}^n} x^{a_1 + a_2 + \dots + a_n}.$$

Now, let’s simplify the equation.

$$\begin{aligned} \text{LHS} &= \sum_{S \in \mathcal{P}_n} x^{|S|} = \sum_{k=0}^n \sum_{S \in \mathcal{P}_{n,k}} x^{|S|} && \text{since } \mathcal{P}_n = \bigcup_{k=0}^n \mathcal{P}_{n,k} \\ &= \sum_{k=0}^n \sum_{S \in \mathcal{P}_{n,k}} x^k && \text{since } |S| = k \quad \forall S \in \mathcal{P}_{n,k} \\ &= \sum_{k=0}^n x^k |\mathcal{P}_{n,k}| && \text{since } x^k \text{ does not depend on } S \\ &= \sum_{k=0}^n \binom{n}{k} x^k && \text{since } |\mathcal{P}_{n,k}| = \binom{n}{k}. \end{aligned}$$

On the RHS, summing over all $\vec{a} \in \{0, 1\}^n$ can be broken up into summing over all $a_1 \in \{0, 1\}$,

$a_2 \in \{0, 1\}, \dots, a_n \in \{0, 1\}$. Thus,

$$\begin{aligned}
\text{RHS} &= \sum_{\vec{a} \in \{0,1\}^n} x^{a_1+a_2+\dots+a_n} = \sum_{(a_1, a_2, \dots, a_n) \in \{0,1\}^n} x^{a_1} x^{a_2} \dots x^{a_n} && \text{(Cartesian product)} \\
&= \sum_{a_1 \in \{0,1\}} \sum_{a_2 \in \{0,1\}} \dots \sum_{a_n \in \{0,1\}} x^{a_1} x^{a_2} \dots x^{a_n} \\
&= \sum_{a_1=0}^1 x^{a_1} \sum_{a_2=0}^1 x^{a_2} \dots \sum_{a_n=0}^1 x^{a_n} \\
&= \left(\sum_{a_1=0}^1 x^{a_1} \right) \left(\sum_{a_2=0}^1 x^{a_2} \right) \dots \left(\sum_{a_n=0}^1 x^{a_n} \right) \\
&= \underbrace{(1+x)(1+x) \dots (1+x)}_{n \text{ times}} \\
&= (1+x)^n. \quad \square
\end{aligned}$$

Lecture 6, 2025/09/15

Theorem 2.3 (Negative Binomial Theorem).

If $t \geq 1$, then

$$(1-x)^{-t} = \sum_{n \geq 0} \binom{n+t-1}{t-1} x^n.$$

Proof. We will use a combinatorial proof. We recognize that $\binom{n+t-1}{t-1}$ is $|\mathcal{M}(n, t)|$, the # of multisets $(m_1, \dots, m_t)$ of size n with elements of t types. Let $\mathcal{M}(t)$ be the set of all multisets (with any # of elements) of t types. Observe that

$$\mathcal{M}(t) = \bigcup_{n \geq 0} \mathcal{M}(n, t).$$

Note that an element of $\mathcal{M}(t)$ is a tuple $(m_1, \dots, m_t)$ with $m_i \in \mathbb{N}$, so $\mathcal{M}(t) = \mathbb{N}^t$. We will start from

the RHS.

$$\begin{aligned}
\sum_{n \geq 0} \binom{n+t-1}{t-1} x^n &= \sum_{n \geq 0} |\mathcal{M}(n, t)| x^n \\
&= \sum_{n \geq 0} \left(\sum_{\mu \in \mathcal{M}(n, t)} 1 \right) x^n \\
&= \sum_{n \geq 0} \sum_{\mu \in \mathcal{M}(n, t)} x^n \\
&= \sum_{n \geq 0} \sum_{\mu \in \mathcal{M}(t)} x^{|\mu|} && \text{since } n = |\mu| \\
&= \sum_{\mu \in \mathcal{M}(t)} x^{|\mu|} && \text{since } \mathcal{M}(t) = \bigcup_{n \geq 0} \mathcal{M}(n, t) \\
&= \sum_{(m_1, \dots, m_t) \in \mathbb{N}^t} x^{m_1 + m_2 + \dots + m_t} && \text{since } \mathcal{M}(t) = \mathbb{N}^t \\
&= \sum_{m_1 \geq 0} \sum_{m_2 \geq 0} \dots \sum_{m_t \geq 0} x^{m_1 + m_2 + \dots + m_t} \\
&= \sum_{m_1 \geq 0} x^{m_1} \sum_{m_2 \geq 0} x^{m_2} \dots \sum_{m_t \geq 0} x^{m_t} \\
&= \left(\sum_{m_1 \geq 0} x^{m_1} \right) \left(\sum_{m_2 \geq 0} x^{m_2} \right) \dots \left(\sum_{m_t \geq 0} x^{m_t} \right) \\
&= \frac{1}{1-x} \cdot \frac{1}{1-x} \dots \frac{1}{1-x} \\
&= \left(\frac{1}{1-x} \right)^t. \quad \square
\end{aligned}$$

Example. Write $\left(\frac{x^2}{1+3x} \right)^4$ as a formal power series.

Proof. Note that

$$\begin{aligned}
\left(\frac{x^2}{1+3x} \right)^4 &= x^8 \left(\frac{1}{1-(-3x)} \right)^4 \\
&= x^8 \sum_{n \geq 0} \binom{n+4-1}{4-1} (-3x)^n \\
&= \sum_{n \geq 0} \binom{n+3}{3} (-3)^n x^{n+8} \\
&= \sum_{n \geq 8} \binom{n-5}{3} (-3)^{n-8} x^n. \quad \square
\end{aligned}$$

Definition 2.4 (Coefficient Extraction).

Let $G(x) = \sum_{n \geq 0} g_n x^n$ be a formal power series. Let $k \in \mathbb{N}$, define

$$[x^k]G(x) = g_k$$

i.e. $[x^k]$ is an operator that extracts the coefficient of x^k from $G(x)$.

Example (Continued).

We have

$$[x^k] \left(\frac{x^2}{1+3x} \right)^4 = [x^k] \sum_{n \geq 8} \binom{n-5}{3} (-3)^{n-8} x^n = \begin{cases} \binom{k-5}{3} (-3)^{k-8}, & k \geq 8, \\ 0, & 0 \leq k < 8. \end{cases}$$

Remark (Some Rules about Coefficient Extraction).

- (1) $[x^k](aF(x) + bG(x)) = a[x^k]F(x) + b[x^k]G(x)$ for $a, b \in \mathbb{R}$.
- (2) $[x^k](x^\ell F(x)) = [x^{k-\ell}]F(x)$ for $\ell \in \mathbb{Z}$.
- (3) $[x^k](F(x)G(x)) = \sum_{\ell=0}^k ([x^\ell]F(x)) \cdot ([x^{k-\ell}]G(x))$.

2.2 Generating Series

We will use formal power series to record counting information about a set, in which case we call it a **generating series**.

Example. Let $\mathcal{M} = \{\text{Jan, Feb, } \dots, \text{Dec}\}$. Let

$$\mathcal{M}_n = \{\alpha \in \mathcal{M} : \alpha \text{ has exactly } n \text{ days (no leap year)}\}.$$

Note that $\mathcal{M} = \bigcup_{n \geq 0} \mathcal{M}_n$. For example,

$$\mathcal{M}_0 = \emptyset, \quad \mathcal{M}_{28} = \{\text{Feb}\}.$$

Consider

$$\sum_{n \geq 0} |\mathcal{M}_n| x^n = x^{28} + 4x^{30} + 7x^{31}.$$

Lecture 7, 2025/09/17

We could derive the same summation in a different way. Consider the following.

$$\sum_{\alpha \in \mathcal{M}} x^{\# \text{ of days in } \alpha} = x^{31} + x^{28} + x^{31} + x^{30} + \cdots + x^{31} = x^{28} + 4x^{30} + 7x^{31}.$$

Definition 2.5 (Weight Function).

Let S be a set. A **weight function** is a function $\omega : S \rightarrow \mathbb{N}$ s.t. for every $n \in \mathbb{N}$, the # of elements of S of weight n is finite, i.e. $\{\alpha \in S : \omega(\alpha) = n\}$ is finite.

Definition 2.6 (Generating Series).

Let S be a set and ω be a weight function on S . The **generating series** of S with respect to ω is

$$\Phi_S^\omega(x) = \Phi_S(x) = \sum_{\alpha \in S} x^{\omega(\alpha)}.$$

Note. Coefficients in a generating series are always non-negative integers.

In our example above,

$$\Phi_{\mathcal{M}}^{\# \text{ of days}}(x) = \sum_{\alpha \in \mathcal{M}} x^{\# \text{ of days in } \alpha} = x^{28} + 4x^{30} + 7x^{31}.$$

Proposition 2.4. Let ω be a weight function on a set S . Then,

$$\Phi_S^\omega(x) = \sum_{n \geq 0} \underbrace{|\{\alpha \in S : \omega(\alpha) = n\}|}_{\# \text{ of elements of weight } n \text{ in } S} x^n.$$

Equivalently,

$$[x^k] \Phi_S^\omega(x) = |\{\alpha \in S : \omega(\alpha) = k\}|.$$

Proof. Let $S_n = \{\alpha \in S : \omega(\alpha) = n\}$. Note that $S = \bigcup_{n \geq 0} S_n$. Then,

$$\Phi_S^\omega(x) = \sum_{\alpha \in S} x^{\omega(\alpha)} = \sum_{n \geq 0} \sum_{\alpha \in S_n} x^{\omega(\alpha)} = \sum_{n \geq 0} \sum_{\alpha \in S_n} x^n = \sum_{n \geq 0} |S_n| x^n.$$

□

Example. Write the generating series for the set S of all binary strings with respect to the weight function ω which records the length of the string.

Proof. We have

$$\begin{aligned} \Phi_S^\omega(x) &= \sum_{\alpha \in S} x^{\omega(\alpha)} \\ &= \sum_{n \geq 0} |\{\alpha \in S : \omega(\alpha) = n\}| x^n = \sum_{n \geq 0} (\# \text{ of binary strings of length } n) x^n \\ &= \sum_{n \geq 0} 2^n x^n \\ &= \sum_{n \geq 0} (2x)^n = \frac{1}{1 - 2x}. \end{aligned}$$

□

Example. Write the generating series for the set

$$S = \{\text{subsets of } \{1, 2, \dots, t\}\}$$

with the weight function $\omega(\alpha) = |\alpha|$.

Proof. We have

$$\begin{aligned} \Phi_S^\omega(x) &= \sum_{\alpha \subseteq \{1, 2, \dots, t\}} x^{|\alpha|} \\ &= \sum_{n \geq 0} (\# \text{ of subsets of size } n) x^n \\ &= \sum_{n \geq 0} \binom{t}{n} x^n \\ &= (1 + x)^t. \end{aligned}$$

□

2.3 Sum and Product Lemmas

Example. Consider breakfast foods and a weight function that corresponds to their cost. Let

$$S_1 = \{\text{omelette, waffles, pancakes, eggs, cereal}\},$$

and $\omega_1 : S_1 \rightarrow \mathbb{N}$ be defined as follows:

$$\omega_1(\text{omelette}) = 10, \quad \omega_1(\text{waffles}) = 10, \quad \omega_1(\text{pancakes}) = 8, \quad \omega_1(\text{eggs}) = 8, \quad \omega_1(\text{cereal}) = 5.$$

The generating series for breakfast main dishes with respect to cost is

$$\Phi_{S_1}^{\omega_1}(x) = \sum_{\alpha \in S_1} x^{\omega_1(\alpha)} = x^{10} + x^{10} + x^8 + x^8 + x^5 = 2x^{10} + 2x^8 + x^5.$$

Equivalently,

$$\Phi_{S_1}^{\omega_1}(x) = \sum_{n \geq 0} |\{\alpha \in S_1 : \omega_1(\alpha) = n\}| x^n = 2x^{10} + 2x^8 + x^5,$$

where the exponents are the costs and the coefficients are the # of things of that cost.

Let

$$S_2 = \{\text{bacon, hashbrowns, toast}\},$$

and $\omega_2 : S_2 \rightarrow \mathbb{N}$ be defined as follows:

$$\omega_2(\text{bacon}) = 5, \quad \omega_2(\text{hashbrowns}) = 4, \quad \omega_2(\text{toast}) = 3.$$

So the generating series for S_2 with respect to cost is

$$\Phi_{S_2}^{\omega_2}(x) = \sum_{\alpha \in S_2} x^{\omega_2(\alpha)} = x^5 + x^4 + x^3.$$

Now, let's consider

$$S_1 \cup S_2 = \{\text{omelette, waffles, pancakes, eggs, cereal, bacon, hashbrowns, toast}\}.$$

The generating series for any breakfast food (main or side) with respect to cost is

$$\Phi_{S_1 \cup S_2}^\omega(x) = \sum_{\alpha \in S_1 \cup S_2} x^{\omega(\alpha)} = 2x^{10} + 2x^8 + x^5 + x^5 + x^4 + x^3 = 2x^{10} + 2x^8 + 2x^5 + x^4 + x^3,$$

Observe that $\Phi_{S_1 \cup S_2}^\omega(x) = \Phi_{S_1}^{\omega_1}(x) + \Phi_{S_2}^{\omega_2}(x)$.

Note. In this case, S_1 and S_2 are disjoint.

Lecture 8, 2025/09/19

Now, consider

$$\begin{aligned} \Phi_{S_1}^{\omega_1}(x) \cdot \Phi_{S_2}^{\omega_2}(x) &= (2x^{10} + 2x^8 + x^5)(x^5 + x^4 + x^3) \\ &= 2x^{15} + 2x^{14} + 2x^{13} + 2x^{13} + 2x^{12} + 2x^{11} + x^{10} + x^9 + x^8 \\ &= 2x^{15} + 2x^{14} + 4x^{13} + 2x^{12} + 2x^{11} + x^{10} + x^9 + x^8. \end{aligned}$$

For example, $2x^{12}$ indicates that there are two (main, side) combos with total price 12. So this is also the generating series for the cartesian product $S_1 \times S_2$ of breakfast combos with respect to cost. This can be denoted as

$$\Phi_{S_1 \times S_2}^{\omega'}(x)$$

where $\omega'(\alpha_1, \alpha_2) = \omega_1(\alpha_1) + \omega_2(\alpha_2)$.

Lemma 2.5 (Sum Lemma).

Let S_1 and S_2 be disjoint sets and let ω be a weight function on $S_1 \cup S_2$. Then,

$$\Phi_{S_1 \cup S_2}^\omega(x) = \Phi_{S_1}^\omega(x) + \Phi_{S_2}^\omega(x).$$

Proof. The main idea is that sums correspond to disjoint unions.

$$\begin{aligned} \Phi_{S_1}(x) + \Phi_{S_2}(x) &= \sum_{\alpha \in S_1} x^{\omega(\alpha)} + \sum_{\alpha \in S_2} x^{\omega(\alpha)} \\ &= \sum_{\alpha \in S_1 \cup S_2} x^{\omega(\alpha)} && \text{since } S_1 \text{ and } S_2 \text{ are disjoint} \\ &= \Phi_{S_1 \cup S_2}^\omega(x). && \square \end{aligned}$$

Lemma 2.6 (Infinite Sum Lemma).

Let $S_0, S_1, S_2, \dots$ be disjoint sets with union S and let ω be a weight function on S . Then,

$$\Phi_S^\omega(x) = \sum_{n \geq 0} \Phi_{S_n}^\omega(x).$$

Proof. Similar to the proof of the Sum Lemma. □

Lemma 2.7 (Product Lemma).

Let S_1 and S_2 be sets and let ω_1 and ω_2 be weight functions on S_1 and S_2 respectively. Then,

$$\Phi_{S_1 \times S_2}^\omega(x) = \Phi_{S_1}^{\omega_1}(x) \cdot \Phi_{S_2}^{\omega_2}(x)$$

where ω is a weight function on $S_1 \times S_2$ defined by

$$\omega(\alpha, \beta) = \omega_1(\alpha) + \omega_2(\beta).$$

Proof. The main idea is that products correspond to cartesian products.

$$\begin{aligned} \Phi_{S_1}^{\omega_1}(x) \cdot \Phi_{S_2}^{\omega_2}(x) &= \sum_{\alpha_1 \in S_1} x^{\omega_1(\alpha_1)} \cdot \sum_{\alpha_2 \in S_2} x^{\omega_2(\alpha_2)} \\ &= \sum_{\alpha_1 \in S_1} \sum_{\alpha_2 \in S_2} x^{\omega_1(\alpha_1) + \omega_2(\alpha_2)} \\ &= \sum_{(\alpha_1, \alpha_2) \in S_1 \times S_2} x^{\omega(\alpha_1, \alpha_2)} \\ &= \Phi_{S_1 \times S_2}^\omega(x). \end{aligned}$$

□

Example. Let $S = \{2, 4, 6, \dots\}$. How many pairs (a, b) are there with $a, b \in S$, and $a + b = 50$?

Proof. Let $\omega : S \rightarrow \mathbb{N}$ be defined by $\omega(\alpha) = \alpha$ and let $\omega' : S \times S \rightarrow \mathbb{N}$ be defined by

$\omega'(\alpha, \beta) = \alpha + \beta$. Note that we are looking for $[x^{50}]\Phi_{S \times S}^{\omega'}(x)$.

$$\begin{aligned}
[x^{50}]\Phi_{S \times S}^{\omega'}(x) &= [x^{50}](\Phi_S^{\omega}(x))^2 && \text{by Product Lemma} \\
&= [x^{50}]\left(\sum_{\alpha \in S} x^{\omega(\alpha)}\right)^2 \\
&= [x^{50}](x^2 + x^4 + x^6 + \dots)^2 \\
&= [x^{50}](x^2(1 + x^2 + x^4 + \dots))^2 \\
&= [x^{50}]\frac{x^4}{(1 - x^2)^2} \\
&= [x^{46}]\frac{1}{(1 - x^2)^2} && \text{by coefficient extraction rule (2)} \\
&= [x^{46}]\sum_{n \geq 0} \binom{n+2-1}{2-1}(x^2)^n && \text{by Negative Binomial Theorem} \\
&= [x^{46}]\sum_{n \geq 0} (n+1)x^{2n} \\
&= 23 + 1 && \text{taking } n = 23 \\
&= 24. && \square
\end{aligned}$$

Definition 2.7 (Star Operator).

Let A be a set. Define

$$A^* = \bigcup_{k \geq 0} A^k = \{\text{all tuples (of any length) of elements of } A\}.$$

Example. $\{0, 1\}^* = \{\underbrace{()}_{A^0}, \underbrace{(\overline{0}), (\overline{1})}_{A^1}, \underbrace{(\overline{0,0}), (\overline{0,1}), (\overline{1,0}), (\overline{1,1})}_{A^2}, \underbrace{(\overline{0,0,0}), \dots}_{A^3, \dots}\}.$

Example. $\{1, 2, 3, \dots\}^* = \{\underbrace{()}_{A^0}, \underbrace{(1), (2), (3), \dots}_{A^1}, \underbrace{(1,1), (1,2), \dots}_{A^2}, \underbrace{(1,1,1), \dots}_{A^3, \dots}\}.$

Given a weight function ω on A , we can define ω^* on A^* such that

$$\omega^*((\alpha_1, \dots, \alpha_k)) = \sum_{i=1}^k \omega(\alpha_i).$$

This is a well-defined weight function provided no elements of A have weight 0 (otherwise, there would be infinitely many tuples of any given weight).

Lecture 9, 2025/09/22

Example. Let $S = \{0, 1\}$ and let ω count length i.e. $\omega(0) = \omega(1) = 1$. Then,

$$\omega^*((1, 1, 0, 1)) = \omega(1) + \omega(1) + \omega(0) + \omega(1) = 4$$

$$\omega^*((1)) = \omega(1) = 1$$

$$\omega^*() = 0.$$

So ω^* also counts length of tuples. Recall that the definition of ω^* requires no elements to have weight 0 under ω . Why?

Suppose γ had weight 0 under ω and β had weight 1 under ω . So we could have an infinite # of tuples of weight 2: e.g. $(\beta, \beta), (\beta, \beta, \gamma), (\beta, \beta, \gamma, \gamma), \dots$

How does $\Phi_S^\omega(x)$ relate to $\Phi_{S^*}^{\omega^*}(x)$? Note that

$$\begin{aligned} \Phi_{S=\{0,1\}}^\omega(x) &= \sum_{\alpha \in S} x^{\omega(\alpha)} = x^{\omega(0)} + x^{\omega(1)} = 2x \\ \Phi_{S^*}^{\omega^*}(x) &= \Phi_{\{(0),(0),(1),(0,0),(0,1),\dots\}}^{\omega^*}(x) \\ &= \sum_{n \geq 0} (\# \text{ of binary strings of length } n) x^n \\ &= \sum_{n \geq 0} 2^n x^n = \sum_{n \geq 0} (2x)^n \\ &= \frac{1}{1 - 2x} \\ &= \frac{1}{1 - \Phi_S^\omega(x)}. \end{aligned}$$

This is not a coincidence! See the following lemma.

Lemma 2.8 (String Lemma).

Let A be a set with weight function ω such that no elements of A have weight 0. Then,

$$\Phi_{A^*}^{\omega^*}(x) = \frac{1}{1 - \Phi_A^\omega(x)}.$$

Proof. We have

$$\begin{aligned}
\Phi_{A^*}^{\omega^*}(x) &= \Phi_{A^0 \cup A^1 \cup A^2 \cup \dots}^{\omega^*}(x) \\
&= \sum_{n \geq 0} \Phi_{A^n}^{\omega^*}(x) && \text{by Infinite Sum Lemma} \\
&= \sum_{n \geq 0} (\Phi_A^{\omega}(x))^n && \text{by Product Lemma} \\
&= \frac{1}{1 - \Phi_A^{\omega}(x)} && \text{by geometric series.}
\end{aligned}$$

□

2.4 Compositions

Definition 2.8 (Composition).

A **composition** is a finite sequence of positive integers

$$\gamma = (c_1, c_2, \dots, c_k).$$

Each c_i , for $i \in \mathbb{Z}_{>0}$, is called a **part**. The **length** of the composition is the # of parts, i.e. $\text{len}(\gamma) = k$. The **size** of the composition is the sum of the parts, i.e. $|\gamma| = c_1 + c_2 + \dots + c_k$. If γ is a composition of size s , then we say that γ is a composition of s .

Example.

- $(1, 4, 3, 2)$ is a composition of size 10 and with 4 parts.
- $(2, 1, 1)$ is a composition of size 4 and with 3 parts.
- $()$ is a composition of size 0 and with 0 parts (the empty composition).

Theorem 2.9. The generating series for all integer compositions with respect to size is

$$\Phi_{\text{compositions}}^{\text{size}}(x) = \frac{1-x}{1-2x} = 1 + \frac{x}{1-2x}.$$

Example. How many compositions are there of 123?

Proof. By the theorem above, we have

$$\begin{aligned}
 \# \text{ of compositions of } 123 &= [x^{123}] \Phi_{\text{compositions}}^{\text{size}}(x) \\
 &= [x^{123}] \left(\frac{1-x}{1-2x} \right) \\
 &= [x^{123}] (1-x) \sum_{n \geq 0} (2x)^n && \text{by geometric series} \\
 &= [x^{123}] \sum_{n \geq 0} (2x)^n - [x^{123}] \sum_{n \geq 0} x(2x)^n \\
 &= 2^{123} - [x^{122}] \sum_{n \geq 0} (2x)^n \\
 &= 2^{123} - 2^{122} \\
 &= 2^{122}.
 \end{aligned}$$

□

Proof of Theorem 2.9. The generating series for all integer compositions with respect to size is

$$\begin{aligned}
 \Phi_{\text{compositions}}^{\text{size}}(x) &= \Phi_{(\mathbb{Z}_{>0})^*}^{\text{size}}(x) && \text{since compositions are tuples of positive integers} \\
 &= \frac{1}{1 - \Phi_{\mathbb{Z}_{>0}}^{\text{size}}(x)} && \text{by String Lemma} \\
 &= \frac{1}{1 - \left(\sum_{n \geq 1} x^n \right)} \\
 &= \frac{1}{1 - x \sum_{n \geq 0} x^n} \\
 &= \frac{1}{1 - x \cdot \frac{1}{1-x}} = \frac{1-x}{1-2x}.
 \end{aligned}$$

□

Lecture 10, 2025/09/24

Theorem 2.10. For $n \in \mathbb{N}$, the # of compositions of size n is

$$\begin{cases} 1, & \text{if } n = 0, \\ 2^{n-1}, & \text{if } n \geq 1. \end{cases}$$

Example. How many compositions are there of 123 where each part is either 1 or 3?

Proof. Let $R = \{1, 3\}$. Then, $\Phi_R^{\text{size}}(x) = x + x^3$. The set of all compositions with each part being 1 or 3 is R^* . So, we want

$$[x^{123}] \Phi_{R^*}^{\text{size}}(x) = [x^{123}] \frac{1}{1 - \Phi_R^{\text{size}}(x)} = [x^{123}] \frac{1}{1 - x - x^3} \quad \text{by String Lemma.}$$

□

Example. Find the generating series for compositions where each part is odd.

Proof. Let $S = \{1, 3, 5, \dots\}$ be the set of odd positive integers. Note that

$$\Phi_S^{\text{size}}(x) = \sum_{n \geq 0} x^{2n+1} = x \sum_{n \geq 0} (x^2)^n = \frac{x}{1 - x^2}.$$

The set of compositions with odd parts is S^* . So, we want

$$\begin{aligned} \Phi_{S^*}^{\text{size}}(x) &= \frac{1}{1 - \Phi_S^{\text{size}}(x)} && \text{by String Lemma} \\ &= \frac{1}{1 - \frac{x}{1 - x^2}} \\ &= \frac{1 - x^2}{1 - x - x^2} \\ &= 1 + \frac{x}{1 - x - x^2}. \end{aligned}$$

□

Example. Find the generating series for compositions where each part is 2 or greater.

Proof. Let $T = \{2, 3, 4, \dots\}$. Note that

$$\Phi_T^{\text{size}}(x) = \sum_{n \geq 2} x^n = x^2 \sum_{n \geq 0} x^n = \frac{x^2}{1 - x}.$$

The set of compositions where each part is ≥ 2 is T^* . By String Lemma, we have

$$\Phi_{T^*}^{\text{size}}(x) = \frac{1}{1 - \Phi_T^{\text{size}}(x)} = \frac{1}{1 - \frac{x^2}{1-x}} = \frac{1-x}{1-x-x^2} = 1 + \frac{x^2}{1-x-x^2}.$$

□

The generating series in the last two examples look rather similar. In fact, there is a strong relation.

Claim. For $n \geq 2$, the # of compositions of n where each part is ≥ 2 is equal to the # of compositions of $n-1$ where each part is odd.

Proof. We want to show that

$$[x^{n-1}] \Phi_{S^*}^{\text{size}}(x) = [x^n] \Phi_{T^*}^{\text{size}}(x).$$

Let's check both sides.

$$\begin{aligned} \text{LHS} &= [x^{n-1}] \left(1 + \frac{x}{1-x-x^2} \right) = [x^{n-1}] 1 + [x^{n-1}] \frac{x}{1-x-x^2} \\ &= 0 + [x^{n-2}] \frac{1}{1-x-x^2} \end{aligned}$$

And,

$$\begin{aligned} \text{RHS} &= [x^n] \left(1 + \frac{x^2}{1-x-x^2} \right) = [x^n] 1 + [x^n] \frac{x^2}{1-x-x^2} \\ &= 0 + [x^{n-2}] \frac{1}{1-x-x^2} \\ &= \text{LHS}. \end{aligned}$$

□

For fixed n , these sets of compositions are finite and of the same size, so can we find a “natural” bijection between them?

Compositions of $n = 7$ with parts ≥ 2	Compositions of $n - 1 = 6$ with odd parts
(7)	(1,5)
(5,2)	(5,1)
(2,5)	(1,1,1,3)
(4,3)	(1,1,3,1)
(3,4)	(1,3,1,1)
(3,2,2)	(3,1,1,1)
(2,3,2)	(3,3)
(2,2,3)	(1,1,1,1,1,1)

Here is a way to construct a bijection.

- (1) Take a composition of 7 with parts ≥ 2 .
- (2) Subtract 1 from the last part to turn into a composition of $n - 1 = 6$.
- (3) For each even part $2k$, split it into $(1, 2k - 1)$.

Compositions of $n = 7$ with parts ≥ 2		Compositions of $n - 1 = 6$ with odd parts
(7)	$\rightarrow$ (6)	$\rightarrow$ (1, 5)
(5,2)	$\rightarrow$ (5, 1)	$\rightarrow$ (5, 1)
(2,5)	$\rightarrow$ (2, 4)	$\rightarrow$ (1, 1, 1, 3)
(4,3)	$\rightarrow$ (4, 2)	$\rightarrow$ (1, 1, 3, 1)
(3,4)	$\rightarrow$ (3, 3)	$\rightarrow$ (3, 3)
(3,2,2)	$\rightarrow$ (3, 2, 1)	$\rightarrow$ (3, 1, 1, 1)
(2,3,2)	$\rightarrow$ (2, 3, 1)	$\rightarrow$ (1, 1, 3, 1)
(2,2,3)	$\rightarrow$ (2, 2, 2)	$\rightarrow$ (1, 1, 1, 1, 1, 1)

Now we have a composition of 6 with only odd parts. However, we still would have to prove that this is a bijection.

3 Binary Strings

Lecture 11, 2025/09/26

Definition 3.1 (Binary String, Bit).

A **binary string** of length $n \geq 0$ is a finite sequence $\sigma = b_1 b_2 \dots b_n$ where each **bit** $b_i \in \{0, 1\}$.

Example. ϵ (the empty string, which has length 0), 0, 1, 00, 01

Binary strings correspond to elements of the set

$$\{0, 1\}^* = \{(), (0), (1), (0, 0), (0, 1), \dots\}.$$

We already know that the # of binary strings of length n is 2^n , but we can also see this using the String Lemma:

$$\begin{aligned} [x^n] \Phi_{\{0,1\}^*}^{\text{length}}(x) &= [x^n] \frac{1}{1 - \Phi_{\{0,1\}}^{\text{length}}(x)} && \text{(String Lemma)} \\ &= [x^n] \frac{1}{1 - 2x} = 2^n && \text{(Geometric Series)} \end{aligned}$$

Definition 3.2 (Concatenation).

If $\sigma = a_1 a_2 \dots a_m$ and $\tau = b_1 b_2 \dots b_n$ are binary strings, then the **concatenation** $\sigma\tau$ is the binary string $a_1 a_2 \dots a_m b_1 b_2 \dots b_n$.

σ^k denotes the k -fold concatenation of σ with itself, i.e. $\sigma^k = \underbrace{\sigma\sigma\dots\sigma}_{k \text{ times}}$.

Definition 3.3 (Subtring).

We say that σ is a **substring** of τ if there exists (possibly empty) binary strings γ_1, γ_2 such that $\tau = \gamma_1 \sigma \gamma_2$.

Example. Let $\sigma = 011$ and $\tau = 10111$. Note that σ is a substring of τ since

$$\tau = \underbrace{1}_{\gamma_1} \underbrace{011}_{\sigma} \underbrace{1}_{\gamma_2}.$$

Definition 3.4 (Block).

A **block** of a binary string s is a non-empty maximal substring of equal bits.

Example. 00011010000 has 5 blocks: 000, 11, 0, 1, 000.

Example. 000 has 1 block: 000.

Definition 3.5 (Concatenation Product).

If S and T are sets of binary strings and $1 \leq k \in \mathbb{N}$, then

$$ST = \{\sigma\tau : \sigma \in S, \tau \in T\} \quad \text{and} \quad S^k = \underbrace{SS \dots S}_{k \text{ times}}.$$

Example. We have the following concatenation products:

- $\{0, 1\}\{\epsilon, 00, 11\} = \{0, 000, 011, 1, 100, 111\}.$
- $\{00, 01\}^2 = \{0000, 0001, 0100, 0101\}.$
- $\{0, 00\}^2 = \{00, 000, 0000\}.$

Note. From the last example above, the size of the concatenation product ST can sometimes be smaller than $|S| \cdot |T|$.

3.1 Regular Expressions

Definition 3.6 (Regular Expression).

A **regular expression** is defined recursively as any of the following:

- $\epsilon, 0$, or 1 ;
- The expression $R \cup S$ where R and S are regular expressions;
- The expression RS where R and S are regular expressions (with R^k where $k \in \mathbb{N}$);
- The expression R^* where R is a regular expression.

Example. The following are all regular expressions:

- 010 .
- $010 \cup 01$.
- $(010 \cup 01)^*$.
- $(11)(010 \cup 01)^*(\epsilon \cup 0^*)$.
- $(00)^5 \cup (\epsilon \cup 0^*)$.

A regular expression is a formula for **producing** a set of binary strings.

Example.

- $R_1 = \epsilon \cup 0 \cup 1$ produces $\mathcal{R}_1 = \{\epsilon, 0, 1\}$.
- $R_2 = (01)(0 \cup 11)$ produces $\mathcal{R}_2 = \{010, 0111\}$.
- $R_3 = (00 \cup 11)^*$ produces $\mathcal{R}_3 = \{\epsilon, 00, 11, 0000, 0011, 1100, 1111, 000000, \dots\}$.

Definition 3.7 (Production).

We define **production** recursively as follows:

- The regular expressions ϵ , 0 , and 1 produce the sets $\{\epsilon\}$, $\{0\}$, and $\{1\}$ respectively.
- If regular expression R produces the set $\mathcal{R}$ and regular expression S produces the set $\mathcal{S}$, then
 - $R \cup S$ produces $\mathcal{R} \cup \mathcal{S}$.
 - RS produces $\mathcal{RS}$ (concatenation product).
 - R^* produces $\mathcal{R}^* = \bigcup_{k \geq 0} \mathcal{R}^k$ (concatenation powers).

Example.

- 0^* produces $\{\epsilon, 0, 00, 000, \dots\}$.
- $(0 \cup 1)^*$ produces $\{0, 1\}^* = \{\text{all binary strings}\}$.
- $(11)0^*$ produces $\{11, 110, 1100, 11000, \dots\}$.

_____ **Lecture 12, 2025/09/29** _____

- $(00 \cup 000)^*$ produces $\{\epsilon, 00, 000, 0000, 00000, \dots\}$.
- $(0 \cup 00)^*$ produces $\{\epsilon, 0, 00, \dots\}$.

Definition 3.8 (Regular Language).

If $\mathcal{R}$ is a set of binary strings that can be produced by a regular expression R , then we say that $\mathcal{R}$ is a **regular language**. This is also known as a **rational language**.

Example.

- $\{1, 110, 011, \epsilon, 0\}$ is a rational language since it can be produced by $1 \cup 110 \cup 011 \cup \epsilon \cup 0$.
- $\{\underbrace{1010 \dots 10}_{m \text{ times}} \underbrace{00 \dots 0}_{n \text{ times}} : m, n \geq 1\}$ is a rational language and can be produced by $10(10)^*00^*$.
- $\{\epsilon, 01, 0011, 000111, \dots, \underbrace{00 \dots 0}_{n \text{ times}} \underbrace{011 \dots 1}_{n \text{ times}}\}$ is NOT a rational language as no regular expression can produce it.

3.2 Unambiguous Expressions**Example.**

- $(0 \cup 01)(0 \cup 10)$ produces $\{00, 010, 010, 0110\} = \{00, 010, 0110\}$. Note that this regular expression produced 010 in two ways.
- $(0 \cup 1)^*$ produces the set of all binary strings, each string exactly one way.
- $(0 \cup 1 \cup 01)^*$ produces the set of all binary strings but some strings are produced more than once.

Definition 3.9 (Unambiguous Expression).

A regular expression R is **unambiguous** if every string in the language $\mathcal{R}$ produced by R is produced in exactly one way by R . Otherwise, R is called **ambiguous**.

Example. Which of the following are unambiguous? Why?

- 010: unambiguous.
- $(0 \cup 1)^*$: unambiguous.
- $(0 \cup 1)^*(0 \cup 1)^*$: ambiguous. For example, 00 can be produced in two ways as $(00)\epsilon$ or $(0)(0)$.

Example. $0^*(10^*)^*$ is an unambiguous regular expression that produces the set of all binary strings. Note that a string produced by $0^*(10^*)^*$ has the form

$$\underbrace{0 \dots 0}_{m_0} (1 \underbrace{0 \dots 0}_{m_1}) (1 \underbrace{0 \dots 0}_{m_2}) \dots (1 \underbrace{0 \dots 0}_{m_k})$$

for $k \geq 0$ and $m_0, m_1, \dots, m_k \geq 0$.

Every binary string can be written uniquely in this form: k is the number of 1's, and m_i is the number of 0's in the block immediately before the $(i + 1)$ th 1.

Example. $1^*(01^*)^*$ is also an unambiguous regular expression that produces the set of all binary strings.

Example. Give an unambiguous regular expression that produces the set of all binary strings where every 0 is followed by at least three 1's:

$$1^*(01111^*)^*$$

Lecture 13, 2025/10/01

Remark (Problem Solving Tips: how to justify that a regular expression is unambiguous).

- Argue how to uniquely decompose the string, for example by the positions of the 1's in $0^*(10^*)^*$ (from example above).
- Start from a regular expression A that we already know is unambiguous, and create a new regular expression B for the desired set of strings. Use the Restriction Lemma (below) to prove that B is unambiguous by arguing that every part of B produces a subset of what's produced by the corresponding part of A .

Lemma 3.1 (Restriction Lemma).

Let A_1, A_2, B_1, B_2 be unambiguous regular expressions producing sets S_1, S_2, T_1, T_2 respectively such that $T_1 \subseteq S_1$ and $T_2 \subseteq S_2$. Then,

- (1) If $A_1 \smile A_2$ is unambiguous, then $B_1 \smile B_2$ is also unambiguous.
- (2) If $A_1 A_2$ is unambiguous, then $B_1 B_2$ is also unambiguous.
- (3) If A_1^* is unambiguous, then B_1^* is also unambiguous.

Example. Show that $1^*(01111^*)^*$ is an unambiguous regular expression that produces the set of all binary strings where every 0 is followed by at least three 1's.

Proof. Applying the Restriction Lemma to show this is unambiguous. The sub-expression 01111^* is itself unambiguous and generates a subset of the strings generated by 01^* (and all the other parts of the two expressions are the same). So Restriction Lemma applies. $\square$

3.3 Translation into Generating Series

Example. How many strings of length 10 are there where every block of 1's has even length?

Proof. Let $\mathcal{R}$ be the set of such strings. An unambiguous expression producing $\mathcal{R}$ is

$$R = (0 \smile 11)^*.$$

Since R is unambiguous, there is a bijection between $\{0, 11\}^*$ and $\mathcal{R}$. Then,

$$\begin{aligned} \# \text{ of strings in } \mathcal{R} \text{ of length } 10 &= \# \text{ of strings in } \{0, 11\}^* \text{ of length } 10 \\ &= [x^{10}] \Phi_{\{0,11\}^*}^{\text{length}}(x) \\ &= [x^{10}] \frac{1}{1 - \Phi_{\{0,11\}}^{\text{length}}(x)} \quad (\text{String Lemma}) \\ &= [x^{10}] \frac{1}{1 - (x + x^2)}. \end{aligned}$$

We say that the regular expression R **leads to** the rational function $\frac{1}{1-(x+x^2)}$. $\square$

Definition 3.10 (Leads to).

A regular expression **leads to** a rational function, defined recursively as follows:

- Regular expressions $\epsilon, 0, 1$ lead to formal power series $1, x, x$ respectively.
- If R and S lead to $f(x)$ and $g(x)$, then
 - $R \cup S$ leads to $f(x) + g(x)$.
 - RS leads to $f(x) \cdot g(x)$.
 - R^* leads to $\frac{1}{1-f(x)}$.

Theorem 3.2. let R be a regular expression that unambiguously produces the set $\mathcal{R}$. Suppose that R leads to $f(x)$. Then, f is the generating series for $\mathcal{R}$ with respect to length.

Example.

- Regular expression 1100 leads to $x^1 \cdot x^1 \cdot x^1 \cdot x^1 = x^4$.
- Regular expression $11 \cup 000$ leads to $x^2 + x^3$.
- Regular expression $0^*(11)(11 \cup 000)^*$ leads to $\frac{1}{1-x} \cdot x^2 \cdot \frac{1}{1-(x^2+x^3)}$.

3.4 Block Decomposition

Proposition 3.3. The regular expression $0^*(11^*00^*)^*1^*$ is unambiguous and produces the set of all binary strings. Same for $1^*(00^*11^*)^*0^*$.

Example. Show how each of the above expression can generate the string 00001101000 .

Proof.

- From $0^*(11^*00^*)^*1^*$: $\underbrace{(0000)}_{\text{from } 0^*} \underbrace{(110)}_{\text{from } 11^*00^*} \underbrace{(10000)}_{\text{from } 11^*00^*} \underbrace{()}_{\text{from } 1^*}$.
- From $1^*(00^*11^*)^*0^*$: $\underbrace{()}_{\text{from } 1^*} \underbrace{(000011)}_{\text{from } 00^*11^*} \underbrace{(01)}_{\text{from } 00^*11^*} \underbrace{(0000)}_{\text{from } 0^*}$.

Justification of unambiguity of the block decomposition: the “forced” 1 and 0 inside the parenthesis act as block delimiters, starting a new block each time. □

Example. Give an unambiguous regular expression for the set of all binary strings and not containing 0000 as a substring. Find the generating series.

Proof. Start from the block decomposition $0^*(11^*00^*)^*1^*$, and restrict the blocks of 0's to be length at most 3. Then,

$$R = (\epsilon \cup 0 \cup 00 \cup 000)(11^*(0 \cup 00 \cup 000))^*1^*.$$

The generating series is

$$(1 + x + x^2 + x^3) \cdot \frac{1}{1 - \left(x \frac{1}{1-x} (x + x^2 + x^3)\right)} \cdot \frac{1}{1-x}.$$

□

Example. Find the generating series for all binary strings avoiding 011 as a substring.

Proof. Start from the block decomposition $1^*(00^*11^*)^*0^*$, and restrict to at most one 1 after any block of 0's. Then,

$$T = 1^*(00^*1)^*0^*.$$

This leads to the generating series

$$\Phi_T^{\text{length}}(x) = \frac{1}{1-x} \cdot \frac{1}{1 - \left(x \frac{1}{1-x} x\right)} \cdot \frac{1}{1-x} = \frac{1}{1+x^3}.$$

□

3.5 Recursive Decompositions

We will define something more powerful than regular expressions, called **recursive expressions**, which can reference themselves.

Example. A recursive expression that produces the set of all binary strings is

$$S = \epsilon \cup S(0 \cup 1).$$

This is unambiguous since every non-empty binary string can be uniquely expressed as another binary string with a 0 or 1 appended to the end.

If S leads to generating series $f(x)$, then

$$S = \epsilon \cup S(0 \cup 1) \text{ leads to } f(x) = x^0 + f(x)(x + x) \implies f(x) = \frac{1}{1 - 2x}.$$

A **recursive decomposition** of a set S describes S in terms of itself using the syntax of regular expressions + an extra symbol S representing the set itself.

A recursive decomposition for S is **unambiguous** if each side of the equation produces each string exactly once.

Note. We don't need to prove that the recursive decomposition is unambiguous in this course.

Example.

- $S = 1S1 \cup 0$ produces $\mathcal{S} = \{0, 101, 11011, \dots\}$. A string $\sigma \in \mathcal{S}$ is either 0 or $1\sigma'1$ for some $\sigma' \in \mathcal{S}$. It is unambiguous because σ' is uniquely determined by σ .
- $S = \epsilon \cup 0 \cup 1S1$ produces $\mathcal{S} = \{\epsilon, 0, 11, 101, 1111, 11011, 111111, \dots\}$. This is also unambiguous.
- $T = 0 \cup 00 \cup 0T$ produces $\mathcal{T} = \{0, 00, 000, 0000, \dots\}$ ambiguously.

An unambiguous recursive decomposition leads to an equation involving the generating series for the set in question.

Example. Let $S = \epsilon \cup 0 \cup 1S1$. This leads to

$$\begin{aligned} \Phi_S(x) &= \Phi_\epsilon(x) + \Phi_0(x) + \Phi_1(x) \cdot \Phi_S(x) \cdot \Phi_1(x) \\ &= 1 + x + x^2\Phi_S(x) \\ &= \frac{1+x}{1-x^2} = \frac{1}{1-x}. \end{aligned}$$

Recursive decompositions allow us to produce sets of strings that are not rational languages.

Example. Give a recursive expression for the set $S = \{0^n 1^n : n \in \mathbb{N}\}$.

Proof. Note that $S = 0S1 \cup \epsilon$ produces S . □

3.6 Excluded Substrings

Example. Find the generating series for the set of strings without 100 as a substring.

Proof. We could solve this with a clever regular expression:

$$0^*(1 \cup 10)^*$$

which leads to the generating series

$$\frac{1}{1-x} \cdot \frac{1}{1-(x+x^2)} = \frac{1}{1-2x+x^3}.$$

However, let's try two other ways. We can do it directly using recursive expressions or we can apply the following theorem. □

Theorem 3.4 (Excluded Substring Theorem).

Let $\kappa \in \{0, 1\}^*$ be a non-empty string of length n and let A be the set of binary strings that avoid κ as a substring. Let C be the set of all non-empty suffices γ of κ such that $\kappa\gamma = \eta\kappa$ for some non-empty prefix η of κ . Then,

$$\Phi_A(x) = \frac{1 + \Phi_C(x)}{(1-2x)(1 + \Phi_C(x)) + x^n},$$

Example (Technique 1: Apply the Excluded Substring Theorem).

Apply the Excluded Substring Theorem to the previous example where $\kappa = 100$.

Proof. The non-empty suffices of κ are $S = \{0, 00\}$. The non-empty prefixes of κ are $P = \{1, 10\}$. Is there a suffix $\gamma \in S$ and a prefix $\eta \in P$ s.t. $\kappa\gamma = \eta\kappa$?

Note that

$$\kappa\gamma = \{1000, 10000\}, \quad \eta\kappa = \{1100, 10100\}.$$

The overlaps are: $C = \emptyset$. So, $\Phi_C(x) = 0$.

Then, by the Excluded Substring Theorem,

$$\Phi_A(x) = \frac{1+0}{(1-2x)(1+0)+x^3} = \frac{1}{1-2x+x^3},$$

□

Example (Technique 2: Recursive Expressions).

Use recursive expressions directly to find the generating series for the set of strings without 100 as a substring.

Proof. Let A be the set of strings that avoid 100 as a substring. Let B be the set of strings with exactly 1 occurrence of 100 at the very end.

Claim (1). $B = A100$.

Claim (2). $A \cup B = \epsilon \cup A(0 \cup 1)$.

If Claim (1) is true and unambiguous, then it leads to

$$\Phi_B(x) = \Phi_A(x)\Phi_{100}(x) = \Phi_A(x) \cdot x^3.$$

If Claim (2) is true and unambiguous, then it leads to

$$\Phi_A(x) + \Phi_B(x) = 1 + \Phi_A(x)(x + x).$$

Then, we can solve for $\Phi_A(x)$:

$$\Phi_A(x) + \Phi_A(x)x^3 = 1 + 2x\Phi_A(x)$$

$$\Phi_A(x)(1 + x^3 - 2x) = 1$$

$$\Phi_A(x) = \frac{1}{1-2x+x^3}.$$

□

4 Partial Fractions and Recurrence Relations

Goal: We want to be able to convert between several representatives of a formal power series $F(x) = \sum_{n \geq 0} a_n x^n$:

- (1) An explicit formula for the coefficients: $f_n = [x^n]F(x)$.
- (2) A rational formula for $F(x)$: $F(x) = \frac{p(x)}{q(x)}$ for polynomials $p(x)$ and $q(x)$.
- (3) A recurrence relation for the coefficients of $F(x)$, i.e., a recurrence relation with initial conditions for sequences $f_0, f_1, \dots$

4.1 Rational Function to Explicit Coefficient Formula Using Partial Fractions

Let's try to extract coefficients when the generating series has a multi-term denominator using partial fractions.

Example. Let's consider the following.

$$\begin{aligned}
 [x^7] \frac{1+7x}{1-x-6x^2} &= [x^7] \frac{1+7x}{(1-3x)(1+2x)} \\
 &= [x^7] \left(\frac{-1}{(1+2x)} + \frac{2}{(1-3x)} \right) && \text{(partial fractions)} \\
 &= [x^7] \left(-1 \sum_{n \geq 0} (-2x)^n + 2 \sum_{n \geq 0} (3x)^n \right) && \text{(geometric series)} \\
 &= [x^7] \sum_{n \geq 0} ((-1)(-2)^n + 2 \cdot 3^n) x^n \\
 &= 2^7 + 2 \cdot 3^7.
 \end{aligned}$$

Lecture 16, 2025/10/08

Theorem 4.1 (Partial Fractions).

Let

$$G(x) = \frac{p(x)}{q(x)} = \frac{p(x)}{(1-\lambda_1 x)^{d_1} (1-\lambda_2 x)^{d_2} \cdots (1-\lambda_s x)^{d_s}},$$

where $\deg(p) < \deg(q)$, the $\lambda_i \in \mathbb{C}$ are distinct, and $d_i \geq 1$. Then, there exists

$C_1^{(1)}, \dots, C_1^{(d_1)}, C_2^{(1)}, \dots, C_2^{(d_2)}, \dots, C_s^{(1)}, \dots, C_s^{(d_s)} \in \mathbb{C}$ such that

$$G(x) = \sum_{i=1}^s \sum_{j=1}^{d_i} \frac{C_i^{(j)}}{(1 - \lambda_i x)^j}.$$

Example. Let $G(x) = \frac{2+5x}{1-3x^2-2x^3}$. Then,

$$\begin{aligned} G(x) &= \frac{2+5x}{(1+x)^2(1-2x)} \\ &= \frac{A}{(1+x)} + \frac{B}{(1+x)^2} + \frac{C}{(1-2x)} \end{aligned} \quad (\text{partial fractions})$$

$$\begin{aligned} 2+5x &= A(1+x)^2 + B(1-2x)(1+x) + C(1-2x) \\ &= (A+B+C) + (-A-2B+2C)x + (-2A+0B-C)x^2. \end{aligned}$$

This leads to a system of equations:

$$\begin{cases} 2 = A + B + C \\ 5 = -A - 2B + 2C \\ 0 = -2A + 0B - C \end{cases}$$

which implies $(A, B, C) = (1, -1, 2)$. So,

$$G(x) = \frac{1}{(1+x)} + \frac{-1}{(1+x)^2} + \frac{2}{(1-2x)}.$$

Now, we can apply the geometric series and negative binomial series to get a single sum:

$$\begin{aligned} G(x) &= \sum_{n \geq 0} (-x)^n - \sum_{n \geq 0} \binom{n+2-1}{2-1} (-x)^n + 2 \sum_{n \geq 0} (2x)^n \\ &= \sum_{n \geq 0} \left((-1)^n - \binom{n+1}{1} (-1)^n + 2 \cdot 2^n \right) x^n \\ &= \sum_{n \geq 0} \left((-1)^n - (n+1)(-1)^n + 2^{n+1} \right) x^n \\ &= \sum_{n \geq 0} \left(2^{n+1} - n(-1)^n \right) x^n. \end{aligned}$$

Also, $[x^n]G(x) = 2^{n+1} - n(-1)^n$.

Remark (Summary).

To get an explicit formula for coefficients of a formal power series given in rational function form, the steps are:

- (1) Factor the denominator.
- (2) Write it as a sum of negative powers of polynomials using partial fractions.
- (3) Solve for e.g. A, B, C by clearing denominators and equating coefficients.
- (4) Use geometric series and negative binomial series to get a single sum.

4.2 Recurrence Relation to Explicit Coefficient Formula

Example. Recall the Fibonacci sequence defined by

$$f_0 = 1, \quad f_1 = 1, \quad f_n = f_{n-1} + f_{n-2} \text{ for } n \geq 2.$$

Let $F(x) = \sum_{n \geq 0} f_n x^n$ be the generating series for the Fibonacci sequence. The recurrence relation is, for $n \geq 2$,

$$\begin{aligned} f_n &= f_{n-1} + f_{n-2} \\ 0 &= f_n - f_{n-1} - f_{n-2} \\ &= [x^n]F(x) - [x^{n-1}]F(x) - [x^{n-2}]F(x) \\ &= [x^n]F(x) - [x^n](xF(x)) - [x^n](x^2F(x)) \\ &= [x^n](1 - x - x^2)F(x). \end{aligned}$$

So for $n \geq 2$, $[x^n](1 - x - x^2)F(x) = 0$. This means that we must have the form

$$(1 - x - x^2)F(x) = C_0 + C_1x + 0x^2 + 0x^3 + \dots \quad \text{for some } C_0, C_1 \in \mathbb{C}.$$

So,

$$F(x) = \frac{C_0 + C_1x}{1 - x - x^2}.$$

To get an explicit form for coefficients of $F(x)$, we first factor the denominator:

$$F(x) = \frac{C_0 + C_1x}{(1 - \alpha x)(1 - \beta x)},$$

where $\alpha = \frac{1+\sqrt{5}}{2}$ and $\beta = \frac{1-\sqrt{5}}{2}$. Using partial fractions, we have

$$F(x) = \frac{A}{(1 - \alpha x)} + \frac{B}{(1 - \beta x)}.$$

Applying the geometric series,

$$F(x) = \sum_{n \geq 0} (A\alpha^n + B\beta^n)x^n.$$

So, $f_n = [x^n]F(x) = A\alpha^n + B\beta^n$. Substituting the initial conditions $f_0 = 1$ and $f_1 = 1$, we get

$$\begin{aligned} f_0 = 1 &= A\alpha^0 + B\beta^0 = A + B, \\ f_1 = 1 &= A\alpha^1 + B\beta^1 = A\frac{1+\sqrt{5}}{2} + B\frac{1-\sqrt{5}}{2}. \end{aligned}$$

Solve the system of 2 linear equations to get

$$A = \frac{5 + \sqrt{5}}{10}, \quad B = -\frac{5 - \sqrt{5}}{10}.$$

We can conclude that

$$f_n = \frac{5 + \sqrt{5}}{10} \left(\frac{1 + \sqrt{5}}{2} \right)^n - \frac{5 - \sqrt{5}}{10} \left(\frac{1 - \sqrt{5}}{2} \right)^n.$$

Lecture 17, 2025/10/10

Remark (Summary).

To get an explicit formula for coefficients of a formal power series given as a recurrence relation, the steps are:

- (1) Start from the recurrence relation.
- (2) Write a rational expression for the generating series and factor the denominator.

- (3) Apply partial fractions to write expression as a sum.
- (4) Apply geometric and negative binomial theorem to obtain coefficient formula.
- (5) Substitute initial conditions to solve for e.g. A, B .

4.3 Convenient Conversion Theorems

The following theorem shows how to go between a recurrence relation for a generating series and a rational function form for the generating series.

Theorem 4.2 (Recurrence Relation $\leftrightarrow$ Rational Function).

Let $G(x) = \sum_{n \geq 0} g_n x^n$ be a generating series. The following are equivalent:

- (1) The sequence $g_0, g_1, \dots$ satisfies the homogeneous linear recurrence relation

$$g_n + a_1 g_{n-1} + \dots + a_d g_{n-d} = 0$$

for all $n \geq N$ with initial conditions $g_0, g_1, \dots, g_{N-1}$.

- (2) $G(x) = \frac{P(x)}{Q(x)}$ where

$$P(x) = b_0 + b_1 x + b_2 x^2 + \dots + b_{N-1} x^{N-1}$$

$$Q(x) = \underbrace{1 + a_1 x + a_2 x^2 + \dots + a_d x^d}_{\text{auxiliary polynomial}}$$

and

$$b_k = g_k + a_1 g_{k-1} + \dots + a_d g_{k-d}$$

for all $0 \leq k \leq N-1$ with the convention that $g_n = 0$ for all $n < 0$.

The following theorem shows how to go from a recurrence relation to an explicit formula for each coefficient.

Theorem 4.3 (Recurrence Relation $\rightarrow$ Explicit Coefficient Formula).

Suppose $g_0, g_1, \dots$ is a sequence satisfying recurrence relation

$$g_n + a_1 g_{n-1} + \dots + a_d g_{n-d} = 0.$$

Factor the auxiliary polynomial to obtain the “inverse roots” $\lambda_i \in \mathbb{C}$ (distinct) and their multiplicities:

$$1 + a_1x + a_2x^2 + \cdots + a_dx^d = (1 - \lambda_1x)^{d_1}(1 - \lambda_2x)^{d_2} \cdots (1 - \lambda_sx)^{d_s}.$$

Then,

$$g_n = p_1(n)\lambda_1^n + p_2(n)\lambda_2^n + \cdots + p_s(n)\lambda_s^n,$$

where each p_i is a polynomial of degree less than d_i .

Example. Suppose g_n is a sequence given by $(g_0, g_1, g_2) = (0, -5, -1)$ and $g_n = 3g_{n-2} - 2g_{n-3}$ for $n \geq 3$. Give a formula for g_n as a function of n .

Proof. Write the recurrence relation as $g_n - 3g_{n-2} + 2g_{n-3} = 0$. The auxiliary polynomial of the recurrence relation is

$$1 - 3x^2 + 2x^3 = (1 - x)^2(1 + 2x).$$

So by Theorem 4.3, we can write

$$g_n = (An + B)1^n + C(-2)^n.$$

Substituting in our initial conditions $g_0 = 0$, $g_1 = -5$, $g_2 = -1$, and solving for A, B, C yields $(A, B, C) = (-2, -1, 1)$. Thus,

$$g_n = -2n - 1 + (-2)^n.$$

□


5 Introduction to Graph Theory

Lecture 18, 2025/10/20

Definition 5.1 (Graph, Vertices, Edges).

A **graph** G consists of a finite non-empty set $V(G)$ of objects, called **vertices**, and a set $E(G)$ of **edges**, which are unordered pairs of distinct vertices.

Example. Let $V(G) = \{1, 2, 3, 4\}$ and $E(G) = \{\{1, 2\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}\}$. Then, G can be represented as follows.



A graph is the relationship: the set of vertices and edges. But, there are many representations of the same graph.

Definition 5.2 (Adjacent).

Two vertices u and v are **adjacent** if $\{u, v\}$ is an edge.

Example. In the above graph, 1 and 2 are adjacent, but 1 and 3 are not adjacent.

Definition 5.3 (Incident).

If $e = \{u, v\}$ is an edge, then we say that edge e is **incident** with vertices u and v .

Example. In the above graph, $e = \{1, 2\}$ is incident with vertices 1 and 2.

Definition 5.4 (Neighbour).

The vertices adjacent to a vertex v are called **neighbours** of v . The set of neighbours of v is denoted $N(v)$ (sometimes $N_G(v)$).


Example. In the above graph, the neighbours of vertex 2 are 1, 3, and 4. So, $N(2) = \{1, 3, 4\}$.

Sometimes we use the shorthand $e = uv$ to denote the edge $e = \{u, v\}$. Edges in this course are unordered (i.e. undirected), so $e = uv = vu$.

Definition 5.5 (Simple Graph).

A **simple graph** is a graph that has no loops (i.e. edges with the same vertex at both ends) and no multiple edges between the same two vertices.

A loop and multiple edges look like this:




In this course, we only deal with simple graphs.


Definition 5.6 (Planar Graph).

A graph is **planar** if it can be drawn without its edges crossing.


Example.



not planar




planar



not planar


Note that the drawing for the middle graph is not planar, but there is a planar drawing of it:



Lecture 19, 2025/10/22

5.1 Isomorphism

Are these two graphs the same?



Answer: Not exactly the same: different labels on the vertices.


Definition 5.7 (Isomorphism).

Two graphs G_1, G_2 are **isomorphic** if $\exists$ a bijection $f : V(G_1) \rightarrow V(G_2)$ such that u, v are adjacent in $G_1 \iff f(u), f(v)$ are adjacent in G_2 . Such a bijection f is called an **isomorphism**.

To show that two graphs are isomorphic: write down the bijection and argue it preserves all adjacency relationships.

To show that two graphs are not isomorphic: find some property that holds in one graph but not the other, such as number of vertices, number of edges, connected or not, planar or not, etc.

Example.




These are not isomorphic because in G_1 , there is a vertex c that is adjacent to 4 vertices, but in G_2 , no vertex is adjacent to 4 vertices.

5.2 Degree

Definition 5.8 (Degree).

The **degree** of a vertex v in a graph G is the number of edges incident with v , and is denoted $\deg(v)$ or $\deg_G(v)$. Equivalently, degree is the size of the neighbourhood: $\deg(v) = |N(v)|$.

Example.



In this graph, $\deg(a) = 2$, $\deg(b) = 3$, $\deg(c) = 3$, and $\deg(d) = 2$. Note that the sum of all degrees is $2 + 3 + 3 + 2 = 10$, which is twice the number of edges.

Theorem 5.1 (Handshaking Lemma).

For every graph G , we have

$$\sum_{v \in V(G)} \deg(v) = 2|E(G)|.$$

Proof. Every edge $e = uv$ has 2 ends, so when we sum the degrees of the vertices, we are counting each edge twice, once at each end (once in $\deg(u)$ and once in $\deg(v)$). $\square$

Corollary 5.2. In any graph, the number of vertices of odd degree is even.

Proof. Let O and E be the sets of vertices with odd and even degree, respectively. Then,

$$\sum_{v \in V(G)} \deg(v) = \sum_{v \in O} \deg(v) + \sum_{v \in E} \deg(v).$$

By the Handshaking Lemma, the LHS is even. By definition of E , $\deg(v)$ is even for all $v \in E$, so $\sum_{v \in E} \deg(v)$ is even. By definition of O , $\deg(v)$ is odd for all $v \in O$. But they sum to an even number, so there must be an even number of odd terms in the sum. Thus, $|O|$ is even. $\square$


Definition 5.9 (k -regular Graph).

A **k -regular graph** is a graph in which all vertices have degree k .

Definition 5.10 (Complete Graph).

A **complete graph** is a graph where every vertex is adjacent to every other vertex. The complete graph on n vertices is denoted K_n . Equivalently, K_n is an $(n - 1)$ -regular graph.

Example. Here are pictures of K_1, K_2, K_3 , and K_4 .



Question: How many edges does K_n have?

Answer: $\frac{n(n-1)}{2} = \binom{n}{2}$ edges.

Question: How many edges does a k -regular graph with n vertices have?

Answer: By the Handshaking Lemma,


$$\sum_{v \in V(G)} \deg(v) = nk = 2|E(G)| \implies |E(G)| = \frac{nk}{2}.$$

5.3 Bipartite Graphs


Definition 5.11 (Bipartite Graph).

A graph G is **bipartite** if its vertex set can be partitioned into two disjoint sets A, B such that $V(G) = A \cup B$ and every edge in G has one endpoint in A and one endpoint in B .


Example. Here are some graphs:




not bipartite



bipartite



bipartite




not bipartite


Definition 5.12 (Complete Bipartite Graph).

For positive integers m, n , the **complete bipartite graph** $K_{m,n}$ is the graph with bipartition A, B where $|A| = m$, $|B| = n$, containing all possible edges joining vertices in A with vertices in B .


Example. $K_{3,3}$, $K_{1,4}$, and $K_{2,3}$ are shown below.



$K_{3,3}$



$K_{1,4}$



$K_{2,3}$


Question: How many edges are in $K_{m,n}$?

Answer: mn edges.

Definition 5.13 (n -Cube or Hypercube).

For $n \geq 0$, the **n -cube** (or **hypercube**) is the graph whose vertex set is the set of all binary

strings of length n , and two vertices are adjacent if their binary strings differ in exactly one position.



Characteristics of the n -cube:

- Number of vertices: 2^n , i.e. the number of binary strings of length n .
- Regularity: n -regular, since for each string s , we can get a neighbourhood of s by changing one of the n positions of s .
- Number of edges: by the Handshaking Lemma,

$$|E(G)| = \frac{1}{2} \sum_{v \in V(G)} \deg(v) = \frac{1}{2} \cdot 2^n \cdot n = n2^{n-1}.$$


- Bipartiteness: the n -cube is bipartite. To see this, let A be the set of binary strings with an odd number of 1's and let B be the set of binary strings with an even number of 1's. Then, (A, B) partitions the vertex set. Let st be an edge in the graph. By definition, t can be obtained from s by changing one position: either change a 0 to a 1 or a 1 to a 0. In either case, the parity of the number of 1's changes, so the parity of the number of 1's in s and t must be different. Hence, one end of st is in A and the other end is in B . So, the n -cube is bipartite.

We can construct the n -cube recursively:

1. Make 2 copies of the $(n - 1)$ -cube.
2. Pretend 0 in front of all the strings in one copy. Pretend 1 in front of all the strings in the other copy.
3. Joint corresponding copies of the vertices with new edges.

5.4 How to Specify a Graph

Example.



Definition 5.14 (Adjacency Matrix).

The **adjacency matrix** of a graph with vertices $\{v_1, v_2, \dots, v_n\}$ is the $n \times n$ matrix A where

$$A_{ij} = \begin{cases} 1 & \text{if } v_i \text{ and } v_j \text{ are adjacent,} \\ 0 & \text{otherwise.} \end{cases}$$

Example (Continued).

The adjacency matrix for the previous graph is

$$A = \begin{matrix} & \begin{matrix} v_1 & v_2 & v_3 & v_4 & v_5 \end{matrix} \\ \begin{matrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \end{matrix} & \begin{pmatrix} 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \end{pmatrix} \end{matrix}.$$

Note. For simple graphs, A is symmetric and its diagonal entries are all 0.

Lecture 21, 2025/10/27

Definition 5.15 (Incidence Matrix).

The **incidence matrix** of a graph with vertices $\{v_1, \dots, v_n\}$ and edges $\{e_1, \dots, e_m\}$ is the $n \times m$

matrix B where

$$B_{ij} = \begin{cases} 1 & \text{if vertex } v_i \text{ is incident with edge } e_j, \\ 0 & \text{otherwise.} \end{cases}$$

Note. Each column of B contains exactly two 1's and each row sums to the degree of that vertex.

Example. Referring to the graph earlier in the section, its incidence matrix is

$$B = \begin{matrix} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \begin{matrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \end{matrix} & \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix} \end{matrix}.$$

Another way of specifying a graph is to give the **adjacency list**, which is a table listing each vertex and the vertices adjacent to it.

For example,


vertex	adjacent vertices
1	2, 4
2	1, 3
3	2, 4, 5
4	1, 3, 5
5	3, 4

Different specifications may be useful or more efficient in different contexts. For example, in a very sparse graph (many vertices, very few edges per vertex), an adjacency list might be more efficient than an incidence matrix or adjacency matrix.

5.5 Subgraphs, Paths, and Cycles

Definition 5.16 (Subgraph, Spanning Subgraph, Proper Subgraph).

A graph H is a **subgraph** of a graph G if the vertex set of H is a (non-empty) subset of G (i.e. $V(H) \subseteq V(G)$) and the edge set of H is a subset of the edge set of G (i.e. $E(H) \subseteq E(G)$) where both endpoints of any edge in $E(H)$ are in $V(H)$. If $V(H) = V(G)$, then H is called a **spanning subgraph**. If $H \neq G$, then H is called a **proper subgraph**.




Definition 5.17 (Edge Deletion and Addition).

Let $e \in E(H)$. $H - e$ is the subgraph of H with vertex set $V(H)$ and edge set $E(H) \setminus \{e\}$. Let e be an edge such that the endpoints of e are in $V(H)$. $H + e$ is the graph with vertex set $V(H)$ and edge set $E(H) \cup \{e\}$.


Definition 5.18 (Walk, Length, Closed Walk).

A u, v -**walk** is a sequence of alternating vertices and edges $u_0, u_0u_1, u_1, u_1u_2, u_2, \dots, u_{k-1}u_k, u_k$ where $u = u_0$ and $v = u_k$. This walk has **length** k . A walk is called **closed** if $u_0 = u_k$.

Example.



*a 1, 4 walk: 1, 12, 2, 24, 4, 34, 3, 35, 5, 45, 4
of length 5*



*a closed 3, 3-walk: 3, 34, 4, 45, 5, 35, 3
of length 3*


We are allowed to repeat vertices and edges in a walk. For simple graphs, it suffices to list the sequence of vertices; edges are implied.

Definition 5.19 (Path).

A u, v -**path** is a u, v -walk with no repeated vertices.

Since no vertices are repeated, no edges are repeated either.

Example.



*a 1, 3-path: 1, 2, 4, 3
of length 3*

We can have the trivial (empty) path or walk of length 0.

Theorem 5.3. If there is a u, v -walk in a graph G , then there is a u, v -path in G .

Idea: Find vertices that are repeated in the walk. Remove the parts between the repetition. Repeat

until no vertices repeat. Use induction on the number of repeated vertices.

Proof. Prove by strong induction on k on the statement “if there is a u, v -walk with k repeated vertices, then there is a u, v -path”.

Base case: when $k = 0$, a u, v -walk with $k = 0$ repeated vertices is a u, v -path.

Inductive hypothesis: if there is a u, v -walk with less than k repeated vertices, then there is a u, v -path.

Inductive step: suppose there is a u, v -walk $v_0, v_1, \dots, v_t$ with k repeated vertices. Let w be a repeated vertex. Let i and j be the smallest and the largest indices, respectively, where $w = v_i = v_j$. Then, $v_0, v_1, \dots, v_i, v_{j+1}, \dots, v_t$ is a u, v -walk. Since w is no longer repeated, this u, v -walk has fewer than k repeated vertices, so by induction, there is a u, v -path. $\square$

Lecture 22, 2025/10/29

Here is another proof using contradiction:

Proof. Suppose $v_0, \dots, v_t$ is a u, v -walk in G with minimal length among all u, v -walks. If the walk is not a path, then there exists some repeated vertex $v_i = v_j$ for $i < j$. Then, $v_0, \dots, v_i, v_{j+1}, \dots, v_t$ is a shorter u, v -walk, which is a contradiction. So, all vertices in our original minimal length walk must be distinct, so it is a u, v -path. $\square$

Corollary 5.4. If there is a u, v -path and a v, w -path in G , then there is a u, w -path in G .

Proof. Join the u, v -path and the v, w -path to make a u, w -walk. Then apply Theorem 5.3 to get a u, w -path. $\square$

Note. This path does not necessarily go through v .


Aside: for vertices in a graph, the relation “ u and v are in a path together” is an equivalence relation. We will see the equivalence classes soon.

Definition 5.20 (Cycle).


A **cycle** of length $n \geq 1$ is a subgraph with n distinct vertices $v_0, v_1, \dots, v_{n-1}$ and n distinct edges $v_0v_1, v_1v_2, \dots, v_{n-2}v_{n-1}, v_{n-1}v_0$.

Example.

A 6-cycle:




A 3-cycle subgraph:



In a simple graph, the minimum length of a cycle is 3 (the triangle).

We can represent a cycle as a closed walk, starting at any vertex.

If uv is an edge of a cycle C , then $C - uv$ is a u, v -path. If P is a u, v -path and P is not just the edge uv , then $P + uv$ is a cycle.



 $C - uv$ is a u, v -path

Every cycle is a 2-regular graph, in other words, every vertex has degree 2. In fact, we have the following theorem.

Theorem 5.5. If every vertex in G has degree at least 2, then G contains a cycle.


Proof. Let $v_0, v_1, \dots, v_k$ be a longest path in G (such a path exists because G is finite and paths cannot repeat vertices). Since v_0 has degree at least 2, it has two neighbours v_1 and x . If x is not on the path, then $x, v_0, v_1, \dots, v_k$ is a longer path in G which is a contradiction. So $x = v_i$ for some $i \geq 2$, and then $v_0, v_1, \dots, v_i, v_0$ is a cycle. $\square$

Exercise: Is the converse true?

Definition 5.21 (Girth).

The **girth** of a graph is the length of its shortest cycle. If G has no cycle, then its girth is ∞ .

Example.




has girth 3.

Definition 5.22 (Hamiltonian Cycle).

A cycle that is a spanning subgraph (i.e. visits all vertices) is called a **Hamiltonian cycle**.

Example.



— A Hamiltonian cycle


Determining if a graph has a Hamiltonian cycle or not is general difficult - it is NP-complete.

5.6 Connectedness


Definition 5.23 (Connected).

A graph is **connected** if there exists a u, v -path for every pair of vertices u, v .

Example.



is connected.



is not connected.

To check if a graph is connected, we could check for the existence of all $\binom{|V(G)|}{2}$ paths, but we can simplify this only have to check $|V(G)| - 1$ paths. See the theorem below.

Lecture 23, 2025/10/31

Theorem 5.6. Let G be a graph and let $u \in V(G)$. Then, G is connected $\iff$ a u, v -path exists for each $v \in V(G)$.

Proof.

($\Rightarrow$): By definition of connected, there is a u, v -path for each $v \in V(G)$.

($\Leftarrow$): To show that G is connected, we need to show there is an x, y -path for any $x, y \in V(G)$. By assumption, there exists a u, x -path P_1 and a u, y -path P_2 . Reverse P_1 to get a x, u -path P_1' . Then, P_1', P_2 is an x, y -walk. By Theorem 5.3, there is an x, y -path. $\square$



Example. Show the n -cube is connected for every $n \geq 0$.

Proof. Let $u = 00 \dots 0$ and let $x \in V(G)$. Suppose x has k 1's in positions $i_1, i_2, \dots, i_k$. Let v_j be the $\{0, 1\}$ string with 1's in positions $i_1, i_2, \dots, i_j$ and 0's everywhere else. Then, $uv_1v_2 \dots v_k$ is a u, x -path. Applying Theorem 5.6, the n -cube is connected. $\square$

Definition 5.24 (Component).

A **component** C of a graph is a connected subgraph of G such that C is not a proper subgraph of any other connected subgraph of G . Equivalently, C is a maximal connected subgraph of G .

Example.




Some observations:

- A connected graph has exactly 1 component.
- A graph with exactly 1 component is connected.
- A disconnected graph has at least 2 components.
- There are no edges joining a vertex of a component with a vertex outside that component. Otherwise, it is not a maximal connected subgraph.

Definition 5.25 (Cut).


Let $X \subseteq V(G)$. The **cut induced by X** in G is the set of all edges in G with exactly one end in X .

Example.



$X = \{1, 2, 3\}$

Cut induced by X : $\{24, 34, 35\}$



$Y = \{1, 2, 3\}$

Cut induced by $Y = \emptyset$

Note. Even though a cut is generated by a set of vertices, the cut itself is a set of edges.

From the second example above (i.e. $Y = \{1, 2, 3\}$), we have the following observation:

- If a graph is disconnected, then the cut induced by the vertices of a component is empty. The converse is not true. For example $X = \emptyset$ induces an empty cut.
- Suppose a graph has an empty cut induced by some non-empty set of vertices X , is the graph disconnected?

Answer: Not necessarily. For example, $X = V(G)$ also induces an empty cut.

- But these are only degenerate cases, leading to the following theorem.

Theorem 5.7. A graph G is disconnected $\iff$ there exists a non-empty proper subset X of $V(G)$ such that the cut induced by X is empty.

Proof.

($\Rightarrow$): Let G be disconnected and let H_1 be a component of G . Since H_1 is connected and G is not, there exists some vertex not in H_1 , so $V(H_1)$ is a non-empty proper subset of $V(G)$. Since H_1 is a component, if y is adjacent to any vertex in H_1 , y must be in H_1 . Hence, the cut induced by $V(H_1)$ is empty.

($\Leftarrow$): Prove by contrapositive. Suppose G is connected, and let X be a proper non-empty subset of $V(G)$. Choose $u \in X$ and $v \in V(G) \setminus X$. Since G is connected, there exists a u, v -path $u = x_0, x_1, \dots, x_n = v$. Let k be the largest index such that $x_k \in X$. Since $x_n = v \notin X$, it must be that $k < n$ and $x_{k+1} \notin X$. Thus, $x_k x_{k+1}$ is in the cut induced by X , so the cut is non-empty. $\square$

5.7 Eulerian Circuits

Definition 5.26 (Eulerian Circuit).

An **Eulerian circuit** (or **Euler tour**) of a graph G is a closed walk that contains every edge of G exactly once.

Lecture 24, 2025/11/03

Question: Which graphs have Eulerian circuits?

Necessary conditions: If G has an Eulerian circuit, what are its properties?

Answer:

- G is not necessarily connected. For example, we may have an isolated vertex. For convenience, we will focus on connected graphs in the rest of this section.
- Every vertex must have even degree. Each time we visit a vertex, we must go in and out using distinct edges, so each visit to a vertex contributes 2 to the degree.

Sufficient conditions: Suppose G is a connected graph where every vertex has even degree. Is it Eulerian?

Answer: Yes, there is a method for finding an Eulerian circuit.

Theorem 5.8. Let G be a connected graph. Then, G has an Eulerian circuit $\iff$ every vertex of G has even degree.

Proof.

($\Rightarrow$): A closed walk contributes 2 to the degree of a vertex for each visit.

($\Leftarrow$): Prove by strong induction on the number of edges m in G .

Base case: G has 0 edges. Then, G is an isolated vertex which has a trivial Eulerian circuit.

Inductive hypothesis: Assume any connected graph with all even degrees and less than m edges has an Eulerian circuit.


Inductive step: Let G be a connected graph with $m \geq 1$ edges and all even degrees. Each vertex has degree ≥ 2 , so by Theorem 5.5, there exists a cycle C . If C contains all the edges of G , then we are done. Otherwise, remove the edges of C from G to get G' . Since G, C both have even degree vertices, so does G' . Each component of G' has fewer edges than G . By induction, each component of G' has an Eulerian circuit. Since G is connected, each component of G' has at least one vertex in common with C . Obtain an Eulerian circuit of G by attaching the Eulerian circuits of each component to C at their common vertex. $\square$

5.8 Bridges

Definition 5.27 (Bridge).

An edge e of G is a **bridge** (or **cut-edge**) if $G - e$ has more components than G .

Example.



A graph G with 2 components
and 3 bridges.

Note. If G is connected, a bridge is an edge such that $G - e$ is not connected.

Lemma 5.9. If $e = xy$ is a bridge in a connected graph G , then $G - e$ has exactly 2 components, and x and y are in different components of $G - e$.

Proof. If e is a bridge of G , then $G - e$ has at least 2 components. Let H be the component of $G - e$ containing x . Let z be any vertex of $G - e$ not in H . Since G is connected, there is a path P in G from x to z . But there is no path from x to z in $G - e$, so P must have contained e . In particular, the x, z -path P in G must be of the form

$$x, e, y, e_2, v_2, \dots, z.$$

Then, $y, e_2, v_2, \dots, z$ is a y, z -path in $G - e$, so z is in the same component of $G - e$ as y , for every z not in H . Thus, $G - e$ has exactly 2 components, one containing x and the other containing y . $\square$

Theorem 5.10. An edge e is a bridge of $G \iff$ it is not contained in any cycle of G .

Remark. This fully characterizes bridges:

- Edge in cycle $\iff$ edge is not a bridge.
- Edge not in cycle $\iff$ edge is a bridge.

Proof. We will prove the contrapositive: e is in a cycle $\iff e$ is not a bridge.

($\Rightarrow$): Suppose $e = uv$ is in a cycle $C = uvv_1v_2 \dots v_ku$. Then, $C - uv$ contains a v, u -path $v, v_1, v_2, \dots, v_k, u$. This path is also in $G - uv$, so u and v are in the same component of $G - e$. By Lemma 5.9, e is not a bridge.

($\Leftarrow$): Suppose $e = uv$ is not a bridge. Suppose that H is the component of G containing e . Then, $H - e$ is connected, so there exists a u, v -path P in $H - e$. Then, $P + uv$ is a cycle in H and also in G . $\square$

6 Trees


Lecture 25, 2025/11/05

6.1 Trees

Definition 6.1 (Tree).

A **tree** is a connected graph with no cycles.


Example.



Definition 6.2 (Forest).

A **forest** is a graph with no cycles.

Example. Below is a forest with 3 components.



Lemma 6.1. Every edge in a tree or forest is a bridge.

Proof. Since a tree or forest has no cycles, each edge e is not in a cycle. By Theorem 5.10, each edge must be a bridge. $\square$


Lemma 6.2. Let x, y be vertices in a tree T . Then, there exists a unique x, y -path in T .

Proof. Suppose there are two x, y -paths in T . Then, there exists some edge uv in one x, y -path but not in the other. Remove edge uv . We can still find a walk u to v , so u and v are in the same component of $T - uv$, and thus uv is not a bridge. $\square$

A tree is a “minimally-connected graph”: it is connected, but removing any edge disconnects it.

Question: How many edges are there in a tree?

Example.



Theorem 6.3. If T is a tree, then $|E(T)| = |V(T)| - 1$.

Idea: Prove by strong induction on the number of edges. Remove one edge, apply induction to the two trees, then add it up.

Proof. Prove using strong induction on the number of edges m .

Base case: when $m = 0$, there are no edges and only 1 vertex, so the statement holds.

Inductive hypothesis: Assume $|E(T')| = |V(T')| - 1$ for all trees T' with fewer than m edges.

Inductive step: Suppose T is a tree with $m \geq 1$ edges. Let $e \in E(T)$. By Lemma 6.1, e is a bridge. By Lemma 5.9, $T - e$ has two components, T_1, T_2 , each of which are trees and each of which has fewer

than m edges. So,

$$\begin{aligned}
 |E(T)| &= |E(T_1)| + |E(T_2)| + 1 \\
 &= (|V(T_1)| - 1) + (|V(T_2)| - 1) + 1 \quad \text{by the inductive hypothesis} \\
 &= |V(T_1)| + |V(T_2)| - 1 \\
 &= |V(T)| - 1.
 \end{aligned}$$

□


Question: How many edges does a forest with n vertices and k components have?

Answer: $n - k$ edges.

Definition 6.3 (Leaf).

A **leaf** in a tree is a vertex of degree 1.

Example.



Theorem 6.4. If T is a tree with at least 2 vertices, then it has at least 2 leaves.

Proof. Let $P = v_1 v_2 \dots v_k$ be a longest path in T . Since T contains at least one edge, P has at least 2 vertices. Consider v_1 . It has a neighbour v_2 on P . It cannot have any other neighbours: if it had a neighbour u not on P , then we would contradict P being a longest path, and if v_1 has another neighbour $v_i \neq v_2$ on P , then we would have a cycle. So v_1 has degree 1 and hence is a leaf. Similarly for v_k . Thus, T has at least 2 leaves. □

Question: How many leaves are there in a tree?

Answer: Let's consider the degrees of all the vertices. Let T be a tree, and let n_i be the number of vertices in T of degree i for $i \geq 1$. We know that

$$|V(T)| = n_1 + n_2 + n_3 + \dots.$$

By Theorem 6.3,

$$|E(T)| = |V(T)| - 1 = -1 + n_1 + n_2 + n_3 + \dots.$$

By the Handshaking Lemma,

$$2|E(T)| = n_1 + 2n_2 + 3n_3 + \dots.$$

Rearrange and solve for n_1 :

$$n_1 = 2 + n_3 + 2n_4 + 3n_5 + \dots = 2 + \sum_{i \geq 3} (i-2)n_i.$$

Observations:

- We see again that every tree with at least 2 vertices has at least 2 leaves: $n_1 = 2 + \dots \geq 2$.
- Vertices of degree 2 have no effect on the number of leaves.
- Higher degree vertices lead to more leaves.

Lemma 6.5. Every tree is bipartite.

Proof. Exercise. Use induction. □

Lecture 26, 2025/11/07

6.2 Spanning Trees

Definition 6.4 (Spanning Tree).

A **spanning tree** of a graph G is a spanning subgraph of G that is a tree. In other words, $V(T) = V(G)$, $E(T) \subseteq E(G)$, and T is a tree.

Theorem 6.6. A graph G is connected $\iff G$ has a spanning tree.

Proof.

($\Leftarrow$): Let T be a spanning tree of G . Since T is connected, for every $x, y \in V(T)$, there exists an x, y -path in T . Since T is a spanning subgraph of G , $V(T) = V(G)$ and $E(T) \subseteq E(G)$. So this x, y -path in T is also an x, y -path in G . Thus, G is connected.

($\Rightarrow$): **Idea:** Remove edges until we get a spanning tree. Which edges can we remove? Edges that are not bridges. Equivalently, edges in cycles. So use induction on the number of cycles.

Prove by strong induction on the number of cycles k : If G is connected with k cycles, then G has a spanning tree.

Base case: when $k = 0$, G is connected and has no cycles, so G is already a spanning tree.

Inductive hypothesis: Assume any connected graph with fewer than k cycles has a spanning tree.

Inductive step: Let G be a connected graph with $k \geq 1$ cycles. Let e be an edge in cycle C of G . By Theorem 5.10, e is not a bridge, so $G - e$ is connected. Moreover, C is no longer a cycle in $G - e$, so $G - e$ has fewer than k cycles. By the inductive hypothesis, $G - e$ has a spanning tree T . Since $V(T) = V(G - e) = V(G)$ and $E(T) \subseteq E(G - e) \subseteq E(G)$, T is a spanning tree of G . $\square$

Corollary 6.7. If G is a connected graph with n vertices and $n - 1$ edges, then G is a tree.

Proof. Since G is connected, it has a spanning tree T which visits all n vertices. A tree on n vertices has $n - 1$ edges. So $E(T) = E(G)$, and thus G is a tree. $\square$

Corollary 6.8. Let G be a graph with n vertices. If any 2 of the following 3 conditions hold, then G is a tree:

- (1) G is connected.
- (2) G has no cycles.
- (3) G has $n - 1$ edges.

6.3 Characterizing Bipartite Graphs

Theorem 6.9. A graph G is bipartite $\iff$ it has no odd cycles.

Proof.

($\Rightarrow$): Suppose G is bipartite with bipartition (A, B) . Let $C = v_1v_2 \dots v_kv_1$ be a cycle of length k . WLOG, assume $v_1 \in A$. Since v_1v_2 is an edge, $v_2 \in B$. And $v_2v_3 \in E$, so $v_3 \in A$ etc. So $v_i \in A \iff i$ is odd. Since v_kv_1 is an edge and $v_1 \in A$, we must have $v_k \in B$, so k is even. Thus, every cycle in G has even length.

($\Leftarrow$): Prove by contrapositive. Suppose G is not bipartite, show that G has an odd cycle. Since G is not bipartite, it has a non-bipartite component H (note that if all components are bipartite, then G is

bipartite). Since H is connected, there is a spanning tree T of H by Theorem 6.6. Trees are bipartite, so let (A, B) be a bipartition of T . But H is not bipartite, so (A, B) is not a bipartition of H . Thus, there is an edge $uv \in E(H)$ such that both $u, v \in A$ or both $u, v \in B$. WLOG, assume $u, v \in A$.

Since T is connected, there is a u, v -path $P = ux_1x_2 \dots x_kv$ in T . Since T is bipartite, we have that $u \in A, x_1 \in B, x_2 \in A, \dots, x_{2i} \in A, x_{2i+1} \in B, \dots, v \in A$. So x_k must be in B , and thus k is odd. So P of length $k + 1$ is an even-length path in H . And $P + uv$ is an odd cycle in H . Hence, G contains an odd cycle. $\square$

7 Planar Graphs


Lecture 27, 2025/11/10

7.1 Planarity


Definition 7.1 (Planar Embedding, Planar Graph).

A **planar embedding** of a graph G is a drawing of the graph in the plane so that its edges intersect only at their ends (i.e. edges do not cross), and no vertices occupy the same point. A graph that has a planar embedding is called a **planar graph**.


Example.



not a planar embedding of K_4



a planar embedding of K_4



not a planar embedding of K_5

For K_5 , there isn't a planar embedding.

Remark. A graph is planar $\iff$ each of its components is planar.

Definition 7.2 (Face).

A **face** of a planar embedding is an undivided region of the plane.

Definition 7.3 (Boundary).

The **boundary** of a face is the subgraph formed from all vertices and edges that touch the face.

Definition 7.4 (Adjacent Faces).

Two faces are **adjacent** if their boundaries have at least one edge in common.


Definition 7.5 (Boundary Walk).

For a planar embedding of a connected graph, the **boundary walk** of a face is a closed walk once around the perimeter of the face boundary.

Definition 7.6 (Face Degree).

The **degree** of a face f is the length of the boundary walk of the face, denoted $\deg(f)$.

Example.



Face	Boundary	Adjacent faces	Boundary walk	Face deg
f_1		f_4	1, 2, 3, 1	3
f_2		f_3, f_4	5, 6, 7, 8, 9, 11, 10, 9, 8, 5	9
f_3		f_2	9, 10, 11, 9	3
f_4		f_1, f_2	1, 2, 3, 4, 3, 5, 8, 7, 6, 5, 3, 1	11

Remark. To find the boundary walk of a face, imagine that the face is a field, the edges are fences, and the vertices are fence posts. Start at a fence post inside the field, and walk around the entire field (with one hand touching the fence at all times) until you get back to where you started. You cannot jump over any fences.

Note. If the graph is not connected, then the boundary may be disconnected, in which case we sum the lengths of the boundary walks on each component of the boundary to get the degree of the face.

Theorem 7.1 (Handshaking Lemma for Faces).

Let G be a planar graph with a planar embedding, where F is the set of all faces. Then,

$$\sum_{f \in F} \deg(f) = 2|E(G)|.$$

Proof. Each edge has 2 sides, and each side contributes 1 to a boundary walk, so the degree contributes 2 to the sum of degrees. $\square$

Lemma 7.2. In a planar embedding, an edge e is a bridge $\iff$ the two sides of e are in the same face.

Note (Not Testable). Our results on planarity can be formalized properly using topology, but that is beyond the scope of this course. To do so, a central result would be:

Theorem 7.3 (Jordan Curve Theorem).


Every planar embedding of a cycle separates the plane into two regions, one on the inside and one on the outside.

This is true for the plane and the surface of the sphere, but is not necessarily true for other surfaces, such as the torus (donut).

Lecture 28, 2025/11/12

7.2 Euler's Formula

Here are two planar embeddings of the same graph:



– face degree

Question: What characteristics with respect to faces do they share? Which are different?

Answer: They have the same number of faces, but different face degrees.

Theorem 7.4 (Euler's Formula).

Let G be a connected graph with n vertices and m edges. Every planar embedding of G has f faces, where $n - m + f = 2$.

Remark. A consequence of this is that the number of faces f in a planar embedding (if it exists) of a graph with n vertices and m edges is $f = 2 - n + m$.

Idea: Induction. Fix the number of vertices, induct on the number of edges. Remove non-bridge edges to keep it connected. Removing a non-bridge edge causes two faces to merge, so edges and faces each decrease by 1. Eventually we have no non-bridge edges: so we have a tree, the planar embedding of which has a single face.

Proof. Fix the number of vertices n . Use induction on the number of edges m .

Base case: The smallest connected graph on n vertices is a tree with $n - 1$ edges, so the base case is $m = n - 1$. Any planar embedding of a tree has one face, so $f = 1$. Then, $n - m + f = n - (n - 1) + 1 = 2$.

Inductive hypothesis: Assume Euler's formula holds for any connected planar graph with n vertices and $m - 1$ edges.

Inductive step: Consider a connected planar graph G with n vertices, m edges, and f faces. Let e be a non-bridge edge. Then, $G - e$ is connected, is planar, and has $m - 1$ edges. So Euler's formula holds for $G - e$ by induction.

Since e is not a bridge, by Lemma 7.2 the two sides of e are in different faces. When we remove e from an embedding of G to get an embedding of $G - e$, these two faces merge into one since they are no longer separated by e . So $G - e$ has $f - 1$ faces. By Euler's formula for $G - e$, $n - (m - 1) + (f - 1) = 2$, which simplifies to $n - m + f = 2$. Thus, Euler's formula holds for G . $\square$

Exercise: If G has c components, then Euler's formula becomes $n - m + f = 1 + c$.


7.3 Platonic Solids

A polyhedron in which all faces have the same degree and all vertices have the same degree is called a **platonic solid**. There are, surprisingly, only 5 platonic solids: the tetrahedron (4 faces), the cube (6 faces), the octahedron (8 faces), the dodecahedron (12 faces), and the icosahedron (20 faces).

Definition 7.7 (Platonic Graph).

We call a graph **platonic** if it has a planar embedding in which every vertex has degree $d \geq 3$ and every face has degree $d^* \geq 3$.


Theorem 7.5. There are exactly 5 platonic graphs.




7.4 Nonplanar Graphs

Our goal in this section is to prove K_5 and $K_{3,3}$ are nonplanar.

K_5 :



$K_{3,3}$:



- Proving that a graph is planar is easy: draw a planar embedding.
- Proving that a graph is nonplanar is harder: a single drawing is not sufficient.

One way to prove nonplanarity: show that there are “too many” edges.

Lecture 29, 2025/11/14

Lemma 7.6. Let G be a planar graph with n vertices and m edges. If there is a planar embedding of G where every face has degree at least $d \geq 3$, then $m \leq \frac{d(n-2)}{d-2}$.

Proof. First, we deal with the case when G is connected. Let F be the set of all faces in the planar embedding, with $|F| = s$. By the Handshaking Lemma for Faces,

$$2m = \sum_{f \in F} \deg(f) \geq \sum_{f \in F} d = ds.$$

By Euler’s formula, $s = 2 - n + m$. Combining these two,

$$2m \geq ds = d(2 - n + m) \implies m \leq \frac{d(n-2)}{d-2}.$$

If G is not connected, make a new connected planar graph by adding edges in the outer face to join the components. The face degrees have not decreased, so the inequality holds for the new graph. But the original graph had fewer edges, so the inequality holds for it as well. $\square$

Note. This lemma relies on a given planar embedding and gives an inequality relying on face degrees. We would like a similar inequality for planar graphs with an inequality that relies on something dependent of an embedding - in this case, just number of vertices and edges.

Theorem 7.7. Let G be a planar graph with $n \geq 3$ vertices and m edges. Then, $m \leq 3n - 6$.

We will use the following lemma without proof to prove this theorem.

Lemma 7.8. If G contains a cycle, then in any planar embedding of G , every face boundary contains a cycle.

Proof of Theorem 7.7. Suppose G does not have a cycle. Then, G is a forest. So $m \leq n - 1 \leq 3n - 6$ when $n \geq 3$. Now suppose G has a cycle. Then, every face boundary has a cycle by the lemma above. Since each cycle has length at least 3, every face has degree at least 3. So by Lemma 7.6,

$$m \leq \frac{3(n-2)}{3-2} = 3n - 6.$$

□

Corollary 7.9. K_5 is not planar.

Proof. Note that $m = |E(K_5)| = \binom{5}{2} = 10$ and $n = |V(K_5)| = 5$. But $3n - 6 = 3(5) - 6 = 9 < 10 = m$, so by Theorem 7.7, K_5 is not planar. □

This can prove a graph is nonplanar, but cannot be used to prove that a graph is planar. There are examples where $m \leq 3n - 6$ but the graph is still nonplanar, such as $K_{3,3}$.

We can modify Theorem 7.7 for bipartite graphs, in which every cycle has length at least 4.

Theorem 7.10. Let G be a bipartite planar graph with $n \geq 3$ vertices and m edges. Then, $m \leq 2n - 4$.

Proof. Suppose G does not have a cycle. Then, $m \leq n - 1 \leq 2n - 4$ whenever $n \geq 3$. Now suppose G has a cycle. Then, every face boundary has a cycle by Lemma 7.8. Since each cycle in a bipartite graph has length at least 4, every face has degree at least 4. So by Lemma 7.6,

$$m \leq \frac{4(n-2)}{4-2} = 2n - 4.$$

□

Corollary 7.11. $K_{3,3}$ is not planar.

Proof. Note that $m = |E(K_{3,3})| = 9$ and $n = |V(K_{3,3})| = 6$. But $2n - 4 = 2(6) - 4 = 8 < 9 = m$, so by Theorem 7.10, $K_{3,3}$ is not planar. □

7.5 Kuratowski's Theorem

We saw that K_5 and $K_{3,3}$ are nonplanar. Clearly, every graph that contains K_5 or $K_{3,3}$ as a subgraph is also nonplanar. But the following graphs technically do not contain K_5 or $K_{3,3}$ as a subgraph, even though they “look like” or are closely related to K_5 or $K_{3,3}$.




So we need another definition to capture how these graphs are related to K_5 or $K_{3,3}$.

Definition 7.8 (Edge Subdivision).

An **edge subdivision** of G is obtained by replacing each edge of G with a new path of length at least 1.

Note. When introducing vertices or edges in a new path in an edge subdivision, those vertices and edges must be new vertices and edges, and can't be used anywhere else in the graph. This means the following graph H is NOT an edge subdivision of K_5 .



is not a subdivision of K_5

Lecture 30, 2025/11/17

Remark (Fact).

A graph is planar $\iff$ any edge subdivision of the graph is planar.

Idea: drawing an edge on the plane is equivalent (from a planarity perspective) to drawing a path on the plane.

From this, we can conclude the following.

Corollary 7.12. If G contains an edge subdivision of $K_{3,3}$ or K_5 as a subgraph, then G is not planar.

Question: Is the converse true?

Answer: Yes. See the following theorem.

Theorem 7.13 (Kuratowski's Theorem).

A graph is planar $\iff$ it does not contain an edge subdivision of $K_{3,3}$ or K_5 as a subgraph.

Proof. See CO 342. □

Remark (Problem Solving Tip: Demonstrating planarity or nonplanarity).

How to show a graph is planar?

- Draw a planar embedding.

How to show a graph is nonplanar?


- Show it has “too many edges” using Theorem 7.7 or Theorem 7.10.
- Or find an edge subdivision of $K_{3,3}$ or K_5 as a subgraph. But how to find $K_{3,3}$ or K_5 ?
 - Try the same process as above to try drawing a planar embedding; when you get stuck on some vertex, start from that vertex to try to build $K_{3,3}$ or K_5 . Note that if there aren't at least 5 vertices of degree at least 4, then it's not going to contain K_5 .

7.6 Colouring

Definition 7.9 (k -Colouring, k -Colourable).


A k -**colouring** of a graph G is an assignment of a colour to each vertex using one of k colours such that adjacent vertices have different colours. If C is a set of size k , then a k -colouring is a function $f : V(G) \rightarrow C$ such that $f(u) \neq f(v)$ for all $uv \in E(G)$. A graph is k -**colourable** if it has a k -colouring.

Example.




5-colouring of G

$\Rightarrow G$ is 5-colourable



4-colouring of G

$\Rightarrow G$ is 4-colourable



3-colouring of G

$\Rightarrow G$ is 3-colourable

Note that the last graph is not 2-colourable since it is not bipartite.

Note. A k -colouring does not have to use all k colours. So a k -colourable graph is also $(k + 1)$ -colourable.

Question: How many colours does it take to colour a graph?

Answer: Every graph with n vertices can be n -coloured. But what is the minimum number?

Extreme Cases:

- A graph with no edges can be 1-colourable.
- Note that:

Theorem 7.14. A graph is 2-colourable $\iff$ it is bipartite.

- Which graphs are n -colourable?

Theorem 7.15. The complete graph K_n is n -colourable, and not k -colourable for any $k < n$.

In fact, the only graph on n vertices that is also not $(n - 1)$ -colourable is K_n .

Lecture 31, 2025/11/19

Theorem 7.16 (6-Colour Theorem).

Every planar graph is 6-colourable.

Here is a key lemma to prove this theorem.

Lemma 7.17. Every planar graph has at least one vertex of degree at most 5.

Proof. Let G be a planar graph with n vertices. If every vertex has degree at least 6, then by the Handshaking Lemma, the number of edges m is at least $\frac{6n}{2} = 3n$. By Theorem 7.7, every planar graph has at most $3n - 6$ edges, so this is a contradiction. Thus, there exists a vertex of degree at most 5. $\square$

Idea for proof of 6-Colour Theorem: Since there is a vertex of degree at most 5, we can colour it even if all its (≤ 5) neighbours have different colours. Remove that vertex; inductively 6-colour the rest, then add it back in.

Proof of 6-Colour Theorem.

We prove by induction on the statement: “Every planar graph with n vertices is 6-colourable.”

Base case: When $n = 1$, clearly a graph with 1 vertex can be coloured with no more than 6 colours.

Inductive hypothesis: Assume that any planar graph with $n - 1$ vertices is 6-colourable.

Inductive step: Let G be a planar graph with n vertices. Let v be a vertex of degree at most 5, which exists by the lemma above. Let $G' = G - v$, i.e. the graph obtained by removing v and all the edges incident with v . So G' is planar with $n - 1$ vertices, so it is 6-colourable.

Let $f' : V(G') \rightarrow \{1, 2, 3, 4, 5, 6\}$ be a 6-colouring of G' . We obtain a 6-colouring $f : V(G) \rightarrow \{1, 2, 3, 4, 5, 6\}$ as follows. Keep the same colours for all vertices except v , i.e. $f(x) = f'(x)$ for all $x \in V(G) \setminus \{v\}$. Let $U \subseteq \{1, 2, 3, 4, 5, 6\}$ be the set of colours used by the neighbours of v :

$$U = \{f(w) : vw \in E(G)\}.$$

Since v has degree at most 5, $|U| \leq 5$. So there is at least one colour c that is not used, i.e. $\{1, 2, 3, 4, 5, 6\} \setminus U \neq \emptyset$. Set $f(v) = c$. Verify that f is a 6-colouring of G :

- For edges xy where $x, y \neq v$, $f'(x) \neq f'(y)$ since $xy \in E(G')$ and f' is a colouring on G' , and $f(x) = f'(x) \neq f'(y) = f(y)$.
- For edges xv , $f(x) \in U$ but $f(v) \notin U$, so $f(x) \neq f(v)$.

$\square$


Theorem 7.18 (5-Colour Theorem).

Every planar graph is 5-colourable.

Definition 7.10 (Edge Contraction).

Let e be an edge in graph G . The graph G/e formed by **contracting** $e = uv$ removes e and squeezes the two ends of e into one vertex, preserving all edges incident with either end.

Example.



Remark (Facts).

- G/e may not be a simple graph, it may have parallel edges, as in the second example G/e' above.
- If G is planar, then G/e is also planar.

Proof of 5-Colour Theorem.

Proof by strong induction on the statement: "Every planar graph with n vertices is 5-colourable."

Base case: $n = 1$ or $n = 2$ is trivial.

Inductive hypothesis: Assume that any planar graph with fewer than n vertices can be coloured with no more than 5 colours.

Inductive step: Let G be a planar graph with n vertices. Let v be a vertex of degree at most 5, which exists by the lemma above. If $\deg(v) \leq 4$, then we can use a similar argument as in the 6-Colour Theorem to prove the graph is 5-colourable. So suppose $\deg(v) = 5$.

There are two neighbours x, y of v that are not adjacent, otherwise the 5 neighbours of v form a K_5 complete subgraph, which cannot exist inside a planar graph (Kuratowski's Theorem).

Let G' be the graph contained from G by contracting edge vx into vertex w , and then edge wy into vertex z . Then, G' is planar with $n - 2$ vertices, so by induction, G' is 5-colourable with colouring f' .

Create a 5-colouring f of G as follows. For any $a \in V(G) \setminus \{v, x, y\}$, set $f(a) = f'(a)$. Set $f(x) = f(y) = f'(z)$; valid since x and y not adjacent in G , and it was okay for all neighbours of x and y to be adjacent to something coloured $f'(z)$. Since 2 neighbours (x, y) of v have the same colour, at most 4 colours have been used in the 5 neighbours of v . Thus, we have at least one colour left to colour v . $\square$

Theorem 7.19 (4-Colour Theorem).

Every planar graph is 4-colourable.

Proof. Exercise. $\square$


7.7 Dual Planar Maps

Definition 7.11 (Dual of a Planar Embedding).

Let G be a planar graph with a planar embedding. The **dual** G^* of G is defined as follows. G^* has one vertex for each face of G . Two vertices of G^* are joined by an edge $\iff$ the corresponding faces of G have an edge in common. The edge in G^* is typically drawn to cross this common boundary edge in G .

Note. The dual of a simple graph may not be a simple graph.

Example.



Remark (Facts).

- The dual of a planar graph is planar.
- The dual of the dual of a planar graph, $(G^*)^*$, is isomorphic to the original graph G .


8 Matchings

8.1 Matching

Definition 8.1 (Matching).

A **matching** of a graph is a set of edges in which no two edges share a common vertex.

Example.



Example (Co-op Job Matching).

We have a set A of students and a set B of jobs. There is an edge between $a \in A$ and $b \in B$ if student a and job b both rank each other. A matching is an assignment of students to jobs so that no student gets more than 1 job and no job gets more than 1 student.

Question: Is it easier to find a small matching or a large matching?

Answer: Small matchings are easier. The empty edge set is a matching in any graph.. So the harder question is:

Question: Given a graph, what is the largest size of a matching?

Lecture 33, 2025/11/24

Definition 8.2 (Saturated Vertex).


In a matching M , a vertex v is **saturated** by M if v is an endpoint of an edge of M .

Definition 8.3 (Perfect Matching).

A matching is a **perfect matching** if it saturates every vertex.

Question: Does every graph have a perfect matching?

Example.




If the graph has a perfect matching, then clearly that matching is maximum.

8.2 Covers

Definition 8.4 (Cover).

A **cover** of a graph G is a set of vertices C such that every edge of G has at least one endpoint in C .

Example.



Lemma 8.1. If M is a matching of G and C is a cover, then $|M| \leq |C|$.

Proof. For each edge $uv \in M$, at least one of u or v is in C since C is a cover. Additionally, no two edges in M saturate the same vertex in C , since M is a matching. Thus, $|M| \leq |C|$. $\square$

Lemma 8.2. If M is a matching of G and C is a cover of G such that $|M| = |C|$, then M is a maximum matching and C is a minimum cover.

Proof. Let M' be a matching. By Lemma 8.1, $|M'| \leq |C| = |M|$. Thus, M is a maximum matching. Let C' be a cover. By Lemma 8.1, $|M| = |C| \leq |C'|$. Thus, C is a minimum cover. $\square$

Note. We can't always find a matching and a cover with equal sizes. However, König's theorem will tell us that we always can for bipartite graphs.

8.3 König's Theorem

Definition 8.5 (Alternating Path).

An **alternating path** P with respect to a matching M is a path where consecutive edges alternate between being in M and not in M .


Definition 8.6 (Augmenting Path).

An **augmenting path** is an alternating path that starts and ends with distinct unsaturated vertices.

Note. A single edge joining two unsaturated vertices is an augmenting path.

Note. Every augmenting path has odd length.

Example.



An augmenting path can be used to make a larger matching by swapping the edges: remove the matching that were on the path, put in the non-matching edges that were on the path. Now we saturate 2 more vertices than before, so matching size increases by 1.

Given the matching M and an augmenting path P , the new matching is

$$M' = (M \cup (E(P) \setminus M)) \setminus (E(P) \cap M).$$

Lecture 34, 2025/11/26

Lemma 8.3. If a matching M has an augmenting path, then M is not maximum.

Theorem 8.4 (König's Theorem).


In a bipartite graph, the size of a maximum matching is equal to the size of a minimum cover.

Definition 8.7 (Bipartite Matching Algorithm or XY-Construction).

Given a bipartite graph G with bipartition (A, B) , and an initial matching M .

- (1) Let X_0 be the set of unsaturated vertices in A .
- (2) Let $X \leftarrow X_0, Y \leftarrow \emptyset$.
- (3) Let N be the set of all neighbours of X in B not currently in Y .
 - (a) If at least one vertex $v \in N$ is unsaturated, then we have found an augmenting path. Make a larger matching by swapping edges in the augmenting path. Then, start over from step (1).
 - (b) If all vertices in N are saturated, then put all of them in Y . Add their matching neighbours to X , and go to step (3).
 - (c) If no such neighbours exist (if $N = \emptyset$), then stop. The matching is maximum, and a minimum cover is $Y \cup (A \setminus X)$.

Example. Let G be the graph shown below, and suppose we start off with the matching M shown below. See Prof's slides for the rest.



Note. At the end of the algorithm:

- X_0 is the set of unsaturated vertices in A .
- X is the set of all vertices in A reachable via an alternating path starting from a vertex in X_0 .
- Y is the set of all vertices in B reachable via an alternating path starting from a vertex in X_0 .
- $A \setminus X$ is the set of all vertices in A **not** reachable via an alternating path starting from X_0 .
 - Every vertex in $A \setminus X$ is saturated, otherwise it would be in X .
- $B \setminus Y$ is the set of all vertices in B **not** reachable via an alternating path starting from X_0 .
 - Some vertices in B might be unsaturated, but they have to be isolated vertices.

Proof of König's Theorem.

Let M be the matching obtained at the end of the bipartite matching algorithm. Let X, X_0, Y be as above.

Claim (1). There is no edge between X and $B \setminus Y$.

Otherwise, we can use this edge to extend an alternating path to $B \setminus Y$. So every edge has an end in either Y or $A \setminus X$, so $Y \cup (A \setminus X)$ is a cover.

Claim (2). Every vertex in Y is saturated.

Otherwise, there would be an augmenting path from X_0 to Y , so the algorithm would not have terminated.

Claim (3). Every vertex in $A \setminus X$ is saturated.

All unsaturated vertices of A are in X_0 , and $X_0 \subseteq X$, so $A \setminus X$ has no unsaturated vertices.

Claim (4). There are no edges between Y and $A \setminus X$.

Otherwise, we would reach $A \setminus X$ via an alternating path. So, the matching edges saturating Y and $A \setminus X$ are distinct.

So, $|M| = |Y| + |A \setminus X| = |Y \cup (A \setminus X)|$. By Lemma 8.2, M is a maximum matching and $Y \cup (A \setminus X)$ is a minimum cover. $\square$


Corollary 8.5. Let G be a bipartite graph with m edges and bipartition A, B with $|A| = |B| = n$. Then, G has a matching of size at least $\frac{m}{n}$.

Proof. Idea: give a lower bound on the minimum size of a cover, then apply König's theorem.

Let C be a cover of G . Every vertex has degree at most n , so there can be at most $n|C|$ edges incident with all of the vertices in C . But C is a cover, so all m edges must be incident with at least one vertex in C . Thus, $n|C| \geq m \implies |C| \geq \frac{m}{n}$. Since every cover has size $\geq \frac{m}{n}$, the minimum cover has size $\geq \frac{m}{n}$. By König's theorem, the maximum matching also has size $\geq \frac{m}{n}$. $\square$

8.4 Hall's Theorem

Question: In a bipartite graph with bipartition A, B , when is it possible to find a matching that saturates all the vertices in A ?



What could prevent us from finding a matching for every vertex in A ?

- An isolated vertex in A .
- A group of vertices in A that match with fewer vertices in B .

Definition 8.8 (Neighbour Set).

Let $D \subseteq V(G)$. The **neighbour set** of D is the set of all vertices adjacent to at least one vertex in D :

$$N(D) = \{v \in V(G) : uv \in E(G) \wedge u \in D\}.$$

Theorem 8.6 (Hall's Theorem).

A bipartite graph G with bipartition (A, B) has a matching saturating every vertex in $A \iff$ every subset $D \subseteq A$ satisfies $|N(D)| \geq |D|$. This is called the **Hall's condition**.

Proof.


($\Rightarrow$): Suppose that M is a matching saturating A . For any $D \subseteq A$, each matching edge that saturates a vertex in D must have the other end in $N(D)$, and these neighbours are distinct, so $|N(D)| \geq |D|$.

($\Leftarrow$): We prove by contrapositive: if G does not have a matching saturating A , then $\exists D \subseteq A$ s.t. $|N(D)| < |D|$.

Lecture 36, 2025/12/01

Let M be a maximum matching in G . So $|M| < |A|$. By König's theorem, there exists a cover C such that $|C| = |M| < |A|$.

Consider $A \cap C, A \setminus C, B \cap C, B \setminus C$:



Claim. $N(A \setminus C) \subseteq B \cap C$.

Proof of Claim. Since C is a cover, no edge joins $A \setminus C$ with $B \setminus C$. Thus, $|N(A \setminus C)| \leq |B \cap C|$. □

Note that $C = (C \cap A) \cup (C \cap B)$ and $A = (A \cap C) \cup (A \setminus C)$. Since $|C| < |A|$, we have

$$|C \cup A| + |C \cup B| < |A \cap C| + |A \setminus C| \implies |C \cap B| < |A \setminus C|.$$

Thus, $|N(A \setminus C)| < |A \setminus C|$. Let $D = A \setminus C$. Then, $|N(D)| < |D|$. □

8.5 Perfect Matchings in Bipartite Graphs

Corollary 8.7. A bipartite graph with bipartition (A, B) has a perfect matching $\iff |A| = |B|$ and every subset $D \subseteq A$ satisfies $|N(D)| \geq |D|$.

Theorem 8.8. If G is a k -regular bipartite graph with $k \geq 1$, then G has a perfect matching.

Proof. First show $|A| = |B|$. By checking degrees of both sides,

$$\sum_{v \in A} \deg(v) = \sum_{v \in B} \deg(v) \implies k|A| = k|B| \implies |A| = |B| \quad \text{since } k \geq 1.$$

Let $D \subseteq A$. Every edge incident with D has its other end in $N(D) \subseteq B$. So,

$$\sum_{v \in D} \deg(v) \leq \sum_{v \in N(D)} \deg(v) \implies k|D| \leq k|N(D)| \implies |D| \leq |N(D)|.$$


By Corollary 8.7, G has a perfect matching. □

8.6 Edge Colourings

Definition 8.9 (Edge k -Colouring).

An **edge k -colouring** of a graph assigns one of k colours to each edge of the graph such that two edges incident with the same vertex are assigned different colours.

Example. Here is a graph with an edge 3-colouring:



Question: How many colours do we need to edge colour a graph?

Answer: At least $\deg(v)$ colours for each vertex v .

Lemma 8.9. Let G be a bipartite graph with at least one edge. Then, G has a matching saturating every vertex of maximum degree.

Theorem 8.10. Every bipartite graph with maximum degree Δ has an edge Δ -colouring.

Proof. Use induction on Δ .

Base case: $\Delta = 0$. So G has no edges, so is edge 0-colourable.

Inductive hypothesis: Assume every bipartite graph with $\Delta < m$ has an edge Δ -colouring.

Inductive step: Let G be bipartite with maximum degree $\Delta = m$. By the previous lemma, G has a matching M saturating all vertices of degree m . Delete M from G to obtain H , which is still bipartite and has $\Delta = m - 1$, so it has an edge $(m - 1)$ -colouring. Use this colouring, plus one more colour for edges in M , to edge m -colour G . $\square$

END OF MATH 239!