

MATH 239 Introduction to Combinatorics

University of Waterloo - Fall 2025

19th September 2025

Instructor: Douglas Stebila

L^AT_EX: Xing Liu

Contents

1	Principles of Enumeration	2
1.1	AND vs. OR - Products vs. Sums	2
1.2	Lists, Permutations, and Subsets	2
1.3	Multisets	5
1.4	Bijections	7
1.5	Summary of Combinatorial and Bijective Proofs	9
2	Generating Series	10
2.1	Binomial Theorem	11
2.2	Generating Series	15
2.3	Sum and Product Lemmas	18

1 Principles of Enumeration

Lecture 1, 2025/09/03

1.1 AND vs. OR - Products vs. Sums

Example. Let $A = \{\text{apple, orange, banana, grape}\}$ and $B = \{\text{carrot, lettuce, onion}\}$. The # of ways of choosing a fruit from A **AND** a vegetable from B is $|A| \cdot |B| = 4 \cdot 3 = 12$. This is equivalent to choosing one element from the cartesian product of A and B :

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}.$$

For finite sets, the cardinalities satisfy

$$|A \times B| = |A| \cdot |B|.$$

The # of ways of choosing a fruit from A **OR** a vegetable from B is $|A| + |B| = 4 + 3 = 7$. This is equivalent to choosing one element from the union of A and B :

$$A \cup B = \{x : x \in A \text{ or } x \in B\}.$$

For disjoint sets,

$$|A \cup B| = |A| + |B|$$

Note that not all sets are disjoint, some have non-trivial intersection.

The intersection is $A \cap B = \{x : x \in A \text{ and } x \in B\}$. For finite sets,

$$|A \cup B| + |A \cap B| = |A| + |B|.$$

1.2 Lists, Permutations, and Subsets

Definition 1.1 (List).

A **list** of a set S is a listing of the elements of S exactly once, in some order.

Definition 1.2 (Permutation).

A **permutation** is a list of the set $\{1, 2, \dots, n\}$ for some $n \in \mathbb{N}$.

Example. $S = \{a, b, c\}$ has 6 lists:

$$abc \quad acb \quad bac \quad bca \quad cab \quad cba.$$

We can construct a list of S by choosing an element $v \in S$ to be the first element of the list, and then append a list of $S \setminus \{v\}$. If p_n is the # of lists of a set of size n , for $n \in \mathbb{N}$, then

$$p_n = n \cdot p_{n-1}$$

and $p_1 = 1$. So by induction we have the following theorem.

Theorem 1.1. For every $n \geq 1$, the # of lists of an n -element set is $n \cdot (n-1) \cdots 2 \cdot 1 = n!$.

Note. By convention, we take $0! = 1$.

Definition 1.3 (Subset).

A **subset** of S is a set containing some (perhaps none, or perhaps all) of the elements of S , at most once each, and in no particular order.

Note. There are 2^n subsets of an n -element set.

To specify: a subset X of S , for each element $v \in S$, we have to decide whether v is in X or not in X . Equivalently, we have to choose {in, out} n times. So,

$$\underbrace{|\{\text{in, out}\}| \cdot |\{\text{in, out}\}| \cdot \dots \cdot |\{\text{in, out}\}|}_{n \text{ times}} = 2^n.$$

Lecture 2, 2025/09/05

Definition 1.4 (Partial List).

A **partial list** of a set S is a list of a subset of S .

We are interested in k -element partial lists of an n -element set, in other words, k -tuples $(s_1, \dots, s_k)$ of distinct elements of S . There are n choices for s_1 , $n - 1$ choices for s_2 , ..., $(n - k + 1)$ choices for s_k . So we have the following.

Theorem 1.2. For $k, n \geq 0$, the # of partial lists of length k of an n -element set is

$$n(n-1)(n-2) \cdots (n-k+1).$$

Note. For $0 \leq k \leq n$, we can write

$$n(n-1)(n-2) \cdots (n-k+1) = \frac{n(n-1) \cdots 1}{(n-k)(n-k-1) \cdots 1} = \frac{n!}{(n-k)!}.$$

Definition 1.5. For $n, k \geq 0$, let $\binom{n}{k}$ (read “ n choose k ”) denote the # of k -element subsets of an n -element set.

Example.

- $\binom{n}{0} = 1$.
- $\binom{n}{n} = 1$ for all n .
- $\binom{n}{1} = n$ for all n .
- $\binom{n}{k} = 0$ if $k > n$.

Theorem 1.3. For $0 \leq k \leq n$, we have

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

Proof. Let S be a set of size n . From Theorem 1.2, the # of partial lists of S of length k is $\frac{n!}{(n-k)!}$. We will choose a k -element subset X of S AND choose a list of X , which will give a list of a subset of S of length k . Thus, the # of length- k partial lists of S is

$$(\# \text{ of } k\text{-element subsets of } S) \cdot (\# \text{ of lists of a } k\text{-element set}) = \binom{n}{k} \cdot k!.$$

$$\text{So, } \binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

□

Note. This is an example of a combinatorial proof, where we prove an identity by counting the same set in two different ways.

Example. For $n \geq 0$,

$$\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n} = 2^n.$$

Prove this by counting sets, not by algebraic manipulation.

Proof. The RHS, 2^n , is the # of subsets of an n -element set. On the LHS, for $0 \leq k \leq n$, $\binom{n}{k}$ is the # of k -element subsets of an n -element set. And addition corresponds to “OR”, so the LHS is the # of ways of choosing a single subset of an n -element set. That is, choosing one of the $\binom{n}{0}$ 0-element subsets, or one of the $\binom{n}{1}$ 1-element subsets, or ..., or one of the $\binom{n}{n}$ n -element subsets. $\square$

Example. For $k, n \geq 1$,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Proof. The LHS, $\binom{n}{k}$, is the # of k -element subsets of an n -element set S . WLOG, assume $S = \{1, 2, \dots, n\}$. Notice that $\binom{n-1}{k}$ is the # of k -element subsets of $S' = \{1, 2, \dots, n-1\}$, and $\binom{n-1}{k-1}$ is the # of $(k-1)$ -element subsets of S' . Recall our intuition that addition corresponds to “OR”. Any k -element subset of S either includes n or does not include n . If $n \notin X$, then X is a subset of size k of S' . If $n \in X$, then $X \setminus \{n\}$ is a subset of size $k-1$ of S' . Thus, LHS = RHS. $\square$

1.3 Multisets

Suppose I have a bag containing 8 marbles, each of which is either red, green, or blue. How many different bag contents are possible?

First question: how can I represent the contents of a bag? For example, we could have 2 red, 3 green, and 3 blue marbles.

Lecture 3, 2025/09/8

- We might want to write it as a set $\{R, R, G, G, G, B, B, B\}$, but this is not a set since sets do not have duplicates.
- Invent new notation that preserves multiplicity:

$$[[R, R, G, G, G, B, B, B]] \text{ or } \langle\langle R^2, G^3, B^3 \rangle\rangle.$$

- Keep track of multiplicities in a vector. Fix an order, say, R then G then B , and use a tuple or a vector to record the multiplicities: $(2, 3, 3)$.

To count the # of bags containing 8 marbles each of which is either red, green, or blue, we need to count the # of 3-tuples (n_1, n_2, n_3) of integers s.t. $n_1 + n_2 + n_3 = 8$ and $n_1, n_2, n_3 \geq 0$.

Definition 1.6 (Multiset).

Let $n \geq 0, t \geq 1$ be integers. A **multiset** of size n with elements of t types is a sequence of non-negative integers $m_1, m_2, \dots, m_t$ s.t. $m_1 + m_2 + \dots + m_t = n$. The set of all multisets of size n with elements of t -types is denoted $\mathcal{M}(n, t)$.

Theorem 1.4. For $n \geq 0, t \geq 1$, the # of n -element multisets with elements of t types is

$$|\mathcal{M}(n, t)| = \binom{n+t-1}{t-1}.$$

Proof. We know that $\binom{n+t-1}{t-1}$ is the # of $(t-1)$ -element subsets of an $(n+t-1)$ -element set. We will show how to translate between $(t-1)$ -element subsets of an $(n+t-1)$ -element set and multisets of size n with t types. Let us work down a row of $n+t-1$ circles (e.g. $n=8, t=3$):

○ ○ ○ ○ ○ ○ ○ ○ ○ ○

Now, let us choose a $(t-1)$ -element subset of these and cross them out

○ ○ × ○ ○ ○ × ○ ○ ○

Our row now has n circles grouped into t segments each containing 0 or more consecutive circles. Let m_i be the length of the i^{th} segment of consecutive circles, so $m_1 + m_2 + \dots + m_t = n$. So $(m_1, m_2, \dots, m_t)$ is a multiset of size n with elements of t types.

Now, let us go the other way. Let $(m_1, m_2, \dots, m_t)$ be a multiset of size n with t types. Write down m_1 circles, then an $\times$, then m_2 circles, then an $\times$, and so on, upto m_t circles. We will have written down $n+t-1$ symbols total, of which $t-1$ are $\times$'s, these indicate the $(t-1)$ -element subset of an $(n+t-1)$ -element set. □

Note. This is an example of a bijective proof.

1.4 Bijections

Definition 1.7 (Surjective, Injective, Bijective).

Let A and B be sets. Let $f : A \rightarrow B$.

- f is **surjective** (or **onto**) if, for every $b \in B$, $\exists a \in A$ s.t. $f(a) = b$.
- f is **injective** (or **one-to-one**) if, for every $a, a' \in A$, if $f(a) = f(a')$, then $a = a'$.
- f is **bijective** if it is both surjective and injective.

Definition 1.8 (Well-defined).

An expression yields a **well-defined** function $f : A \rightarrow B$ if, for every input $a \in A$, $f(a)$ assigns a single unambiguous value $b \in B$.

Proposition 1.5. A function $f : A \rightarrow B$ is a bijection $\iff f$ has an inverse, namely if $\exists$ a well-defined function $g : B \rightarrow A$ s.t. $g(f(a)) = a$ for all $a \in A$ and $f(g(b)) = b$ for all $b \in B$.

Remark (Notation).

If $\exists$ a bijection between sets A and B , we can write $A \rightleftharpoons B$.

Corollary 1.6. If $A \rightleftharpoons B$ and at least one of A or B is finite, then $|A| = |B|$.

Example. For $k, n \geq 1$,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Proof. Let $S = \{1, 2, \dots, n\}$ and $S' = \{1, 2, \dots, n-1\}$. Let $A = \{k\text{-element subsets of } S\}$ and $B = \{(k-1)\text{-element or } k\text{-element subsets of } S'\}$. By definition, $|A| = \binom{n}{k}$ and we can easily see that $|B| = \binom{n-1}{k-1} + \binom{n-1}{k}$. The goal is to prove $A \rightleftharpoons B$. Define $f : A \rightarrow B$ by

$$f(u) = \begin{cases} u, & \text{if } n \notin u \\ u \setminus \{n\}, & \text{if } n \in u \end{cases}$$

Lecture 4, 2025/09/10

Question: Is f well-defined, i.e. $\forall u \in A$, is $f(u) \in B$?

Answer: Yes, $f(u)$ does not contain n (so it's a subset of S') and has size either $k-1$ or k . So

$f(u) \in B$.

Define $g : B \rightarrow A$ by

$$g(v) = \begin{cases} v, & \text{if } |v| = k \\ v \cup \{n\}, & \text{if } |v| = k - 1 \end{cases}$$

Claim (1). For all $u \in A$, $g(f(u)) = u$.

Proof of Claim (1). First consider $u \in A$ s.t. $n \notin u$. Then, $f(u) = u$ and $|f(u)| = k$, so $g(f(u)) = u$ (first branch of g). Now consider $u \in A$ s.t. $n \in u$. Then, $f(u) = u \setminus \{n\}$ and $|f(u)| = k - 1$, so $g(f(u)) = (u \setminus \{n\}) \cup \{n\} = u$ (second branch of g). $\square$

Claim (2). For all $v \in B$, $f(g(v)) = v$.

Proof of Claim (2). Exercise. $\square$

Thus, f is a bijection, so $|A| = |B|$, i.e. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$. $\square$

Remark (Problem Solving Tips: what steps are needed for a bijective argument?).


To show a function $f : A \rightarrow B$ is a bijection by giving an inverse:

- (1) State the function $f : A \rightarrow B$.
- (2) Show that f is well-defined, i.e. $\forall a \in A, f(a) \in B$.
- (3) State the function $g : B \rightarrow A$.
- (4) Show that g is well-defined.
- (5) Show that $g(f(a)) = a$ for all $a \in A$.
- (6) Show that $f(g(b)) = b$ for all $b \in B$.


Exercise: Show that $\binom{n}{k} = \binom{n}{n-k}$ for $n \geq k \geq 0$ using a combinatorial or bijective argument.

1.5 Summary of Combinatorial and Bijective Proofs

In a combinatorial proof, describe two different ways of counting the size of the same set.



In a bijective proof, describe two different sets and show they're in bijection.



- (1) Give sets A, B and count them.
- (2) Show a bijection between A and B .
- (3) This implies $|A| = |B|$.

2 Generating Series

Definition 2.1 (Formal Power Series).

A **formal power series** is an expression of the form

$$G(x) = \sum_{n=0}^{\infty} g_n x^n$$

in which the coefficients $(g_0, g_1, g_2, \dots)$ are a sequence of real numbers.

Note. We use x as an **indeterminate** for which we do not normally substitute in particular values. This means that we never worry about the convergence of the series, all we require is that the coefficients are finite real numbers.

Example. $G(x) = 0 - 1x + 2x^2 - 3x^3 + \dots = \sum_{n=0}^{\infty} (-1)^n n x^n$.

We can add and multiply formal power series like polynomials. Let

$$F(x) = \sum_{n \geq 0} f_n x^n = f_0 + f_1 x + f_2 x^2 + \dots,$$

$$G(x) = \sum_{n \geq 0} g_n x^n = g_0 + g_1 x + g_2 x^2 + \dots.$$

Then,

$$(F + G)(x) = \sum_{n \geq 0} (f_n + g_n) x^n = (f_0 + g_0) + (f_1 + g_1)x + (f_2 + g_2)x^2 + \dots,$$

$$(F \cdot G)(x) = \sum_{n \geq 0} \left(\sum_{k=0}^n f_k g_{n-k} \right) x^n = (f_0 g_0) + (f_0 g_1 + f_1 g_0)x + (f_0 g_2 + f_1 g_1 + f_2 g_0)x^2 + \dots.$$

Lecture 5, 2025/09/12

Example. Note that

$$\begin{aligned} (1 + 3x^2 + 5x^4 + \dots) + x(2 + 4x^2 + 6x^4 + \dots) &= (1 + 3x^2 + 5x^4 + \dots) + (2x + 4x^3 + 6x^5 + \dots) \\ &= 1 + 2x + 3x^2 + 4x^3 + 5x^4 + 6x^5 + \dots \end{aligned}$$

Equivalently,

$$\begin{aligned} \sum_{n \geq 0} (2n+1)x^{2n} + x \sum_{n \geq 0} (2n+2)x^{2n} &= \sum_{n \geq 0} (2n+1)x^{2n} + \sum_{n \geq 0} (2n+2)x^{2n+1} \\ &= \sum_{n \geq 0} ((2n+1)x^{2n} + (2n+2)x^{2n+1}) \\ &= \sum_{n \geq 0} (n+1)x^n. \end{aligned}$$

Definition 2.2 (Geometric Series).

The **geometric series** is of the form

$$G(x) = 1 + x + x^2 + x^3 + \cdots = \sum_{n \geq 0} x^n.$$

Consider $xG(x) = x + x^2 + x^3 + \cdots = x \sum_{n \geq 0} x^n$. Then,

$$G(x) - xG(x) = 1 \implies G(x)(1-x) = 1 \implies G(x) = \frac{1}{1-x} = (1-x)^{-1}.$$

Definition 2.3 (Inverse of a Formal Power Series).

If $F(x)$ and $G(x)$ are formal power series such that $F(x)G(x) = 1$, then we write $G(x) = F(x)^{-1}$ or $G(x) = \frac{1}{F(x)}$, and say that $G(x)$ is the **inverse** of $F(x)$.

Corollary 2.1. The inverse of $F(x) = \sum_{n \geq 0} f_n x^n$ exists $\iff f_0 \neq 0$.

2.1 Binomial Theorem

Theorem 2.2 (Binomial Theorem).

For any $n \in \{0, 1, 2, \dots\}$,

$$(1+x)^n = \sum_{k=0}^n \binom{n}{k} x^k = \sum_{k \geq 0} \binom{n}{k} x^k.$$

Proof. We will use a combinatorial proof. Let $\mathcal{P}_n$ be the power set of $\{1, 2, \dots, n\}$, i.e. the set of all subsets of $\{1, 2, \dots, n\}$. Let $\mathcal{P}_{n,k}$ be the set of all k -element subsets of $\{1, 2, \dots, n\}$. We saw before that a

bijection between $\mathcal{P}_n$ and $\{0, 1\}^n$, i.e. a subset S of $\{1, 2, \dots, n\}$ corresponds to an “indicator vector” in $\{0, 1\}^n$. Let $S \in \mathcal{P}_n$ be a subset of $\{1, 2, \dots, n\}$. The corresponding indicator vector will be

$$\vec{a} = (a_1, a_2, \dots, a_n) \quad \text{where} \quad a_i = \begin{cases} 0, & \text{if } i \notin S, \\ 1, & \text{if } i \in S. \end{cases}$$

Note that $|S| = \sum_{i=1}^n a_i$. Let's introduce an indeterminate x . For every subset S with indicator vector $\vec{a}$, it holds that

$$x^{|S|} = x^{a_1 + a_2 + \dots + a_n}.$$

Because of the bijection between $\mathcal{P}_n$ and $\{0, 1\}^n$, summing over all subsets is equivalent to summing over all indicator vectors. Thus,

$$\sum_{S \in \mathcal{P}_n} x^{|S|} = \sum_{\vec{a} \in \{0, 1\}^n} x^{a_1 + a_2 + \dots + a_n}.$$

Now, let's simplify the equation.

$$\begin{aligned} \text{LHS} &= \sum_{S \in \mathcal{P}_n} x^{|S|} = \sum_{k=0}^n \sum_{S \in \mathcal{P}_{n,k}} x^{|S|} && \text{since } \mathcal{P}_n = \bigcup_{k=0}^n \mathcal{P}_{n,k} \\ &= \sum_{k=0}^n \sum_{S \in \mathcal{P}_{n,k}} x^k && \text{since } |S| = k \quad \forall S \in \mathcal{P}_{n,k} \\ &= \sum_{k=0}^n x^k |\mathcal{P}_{n,k}| && \text{since } x^k \text{ does not depend on } S \\ &= \sum_{k=0}^n \binom{n}{k} x^k && \text{since } |\mathcal{P}_{n,k}| = \binom{n}{k}. \end{aligned}$$

On the RHS, summing over all $\vec{a} \in \{0, 1\}^n$ can be broken up into summing over all $a_1 \in \{0, 1\}$,

$a_2 \in \{0, 1\}, \dots, a_n \in \{0, 1\}$. Thus,

$$\begin{aligned}
\text{RHS} &= \sum_{\vec{a} \in \{0,1\}^n} x^{a_1+a_2+\dots+a_n} = \sum_{(a_1, a_2, \dots, a_n) \in \{0,1\}^n} x^{a_1} x^{a_2} \dots x^{a_n} && \text{(Cartesian product)} \\
&= \sum_{a_1 \in \{0,1\}} \sum_{a_2 \in \{0,1\}} \dots \sum_{a_n \in \{0,1\}} x^{a_1} x^{a_2} \dots x^{a_n} \\
&= \sum_{a_1=0}^1 x^{a_1} \sum_{a_2=0}^1 x^{a_2} \dots \sum_{a_n=0}^1 x^{a_n} \\
&= \left(\sum_{a_1=0}^1 x^{a_1} \right) \left(\sum_{a_2=0}^1 x^{a_2} \right) \dots \left(\sum_{a_n=0}^1 x^{a_n} \right) \\
&= \underbrace{(1+x)(1+x) \dots (1+x)}_{n \text{ times}} \\
&= (1+x)^n. \quad \square
\end{aligned}$$

Lecture 6, 2025/09/15

Theorem 2.3 (Negative Binomial Theorem).

If $t \geq 1$, then

$$(1-x)^{-t} = \sum_{n \geq 0} \binom{n+t-1}{t-1} x^n.$$

Proof. We will use a combinatorial proof. We recognize that $\binom{n+t-1}{t-1}$ is $|\mathcal{M}(n, t)|$, the # of multisets $(m_1, \dots, m_t)$ of size n with elements of t types. Let $\mathcal{M}(t)$ be the set of all multisets (with any # of elements) of t types. Observe that

$$\mathcal{M}(t) = \bigcup_{n \geq 0} \mathcal{M}(n, t).$$

Note that an element of $\mathcal{M}(t)$ is a tuple $(m_1, \dots, m_t)$ with $m_i \in \mathbb{N}$, so $\mathcal{M}(t) = \mathbb{N}^t$. We will start from

the RHS.

$$\begin{aligned}
\sum_{n \geq 0} \binom{n+t-1}{t-1} x^n &= \sum_{n \geq 0} |\mathcal{M}(n, t)| x^n \\
&= \sum_{n \geq 0} \left(\sum_{\mu \in \mathcal{M}(n, t)} 1 \right) x^n \\
&= \sum_{n \geq 0} \sum_{\mu \in \mathcal{M}(n, t)} x^n \\
&= \sum_{n \geq 0} \sum_{\mu \in \mathcal{M}(t)} x^{|\mu|} && \text{since } n = |\mu| \\
&= \sum_{\mu \in \mathcal{M}(t)} x^{|\mu|} && \text{since } \mathcal{M}(t) = \bigcup_{n \geq 0} \mathcal{M}(n, t) \\
&= \sum_{(m_1, \dots, m_t) \in \mathbb{N}^t} x^{m_1 + m_2 + \dots + m_t} && \text{since } \mathcal{M}(t) = \mathbb{N}^t \\
&= \sum_{m_1 \geq 0} \sum_{m_2 \geq 0} \dots \sum_{m_t \geq 0} x^{m_1 + m_2 + \dots + m_t} \\
&= \sum_{m_1 \geq 0} x^{m_1} \sum_{m_2 \geq 0} x^{m_2} \dots \sum_{m_t \geq 0} x^{m_t} \\
&= \left(\sum_{m_1 \geq 0} x^{m_1} \right) \left(\sum_{m_2 \geq 0} x^{m_2} \right) \dots \left(\sum_{m_t \geq 0} x^{m_t} \right) \\
&= \frac{1}{1-x} \cdot \frac{1}{1-x} \dots \frac{1}{1-x} \\
&= \left(\frac{1}{1-x} \right)^t. \quad \square
\end{aligned}$$

Example. Write $\left(\frac{x^2}{1+3x} \right)^4$ as a formal power series.

Proof. Note that

$$\begin{aligned}
\left(\frac{x^2}{1+3x} \right)^4 &= x^8 \left(\frac{1}{1-(-3x)} \right)^4 \\
&= x^8 \sum_{n \geq 0} \binom{n+4-1}{4-1} (-3x)^n \\
&= \sum_{n \geq 0} \binom{n+3}{3} (-3)^n x^{n+8} \\
&= \sum_{n \geq 8} \binom{n-5}{3} (-3)^{n-8} x^n. \quad \square
\end{aligned}$$

Definition 2.4 (Coefficient Extraction).

Let $G(x) = \sum_{n \geq 0} g_n x^n$ be a formal power series. Let $k \in \mathbb{N}$, define

$$[x^k]G(x) = g_k$$

i.e. $[x^k]$ is an operator that extracts the coefficient of x^k from $G(x)$.

Example (Continued).

We have

$$[x^k] \left(\frac{x^2}{1+3x} \right)^4 = [x^k] \sum_{n \geq 8} \binom{n-5}{3} (-3)^{n-8} x^n = \begin{cases} \binom{k-5}{3} (-3)^{k-8}, & k \geq 8, \\ 0, & 0 \leq k < 8. \end{cases}$$

Remark (Some Rules about Coefficient Extraction).

- (1) $[x^k](aF(x) + bG(x)) = a[x^k]F(x) + b[x^k]G(x)$ for $a, b \in \mathbb{R}$.
- (2) $[x^k](x^\ell F(x)) = [x^{k-\ell}]F(x)$ for $\ell \in \mathbb{Z}$.
- (3) $[x^k](F(x)G(x)) = \sum_{\ell=0}^k ([x^\ell]F(x)) \cdot ([x^{k-\ell}]G(x))$.

2.2 Generating Series

We will use formal power series to record counting information about a set, in which case we call it a **generating series**.

Example. Let $\mathcal{M} = \{\text{Jan, Feb, } \dots, \text{Dec}\}$. Let

$$\mathcal{M}_n = \{\alpha \in \mathcal{M} : \alpha \text{ has exactly } n \text{ days (no leap year)}\}.$$

Note that $\mathcal{M} = \bigcup_{n \geq 0} \mathcal{M}_n$. For example,

$$\mathcal{M}_0 = \emptyset, \quad \mathcal{M}_{28} = \{\text{Feb}\}.$$

Consider

$$\sum_{n \geq 0} |\mathcal{M}_n| x^n = x^{28} + 4x^{30} + 7x^{31}.$$

Lecture 7, 2025/09/17

We could derive the same summation in a different way. Consider the following.

$$\sum_{\alpha \in \mathcal{M}} x^{\# \text{ of days in } \alpha} = x^{31} + x^{28} + x^{31} + x^{30} + \cdots + x^{31} = x^{28} + 4x^{30} + 7x^{31}.$$

Definition 2.5 (Weight Function).

Let S be a set. A **weight function** is a function $\omega : S \rightarrow \mathbb{N}$ s.t. for every $n \in \mathbb{N}$, the # of elements of S of weight n is finite, i.e. $\{\alpha \in S : \omega(\alpha) = n\}$ is finite.

Definition 2.6 (Generating Series).

Let S be a set and ω be a weight function on S . The **generating series** of S with respect to ω is

$$\Phi_S^\omega(x) = \Phi_S(x) = \sum_{\alpha \in S} x^{\omega(\alpha)}.$$

Note. Coefficients in a generating series are always non-negative integers.

In our example above,

$$\Phi_{\mathcal{M}}^{\# \text{ of days}}(x) = \sum_{\alpha \in \mathcal{M}} x^{\# \text{ of days in } \alpha} = x^{28} + 4x^{30} + 7x^{31}.$$

Proposition 2.4. Let ω be a weight function on a set S . Then,

$$\Phi_S^\omega(x) = \sum_{n \geq 0} \underbrace{|\{\alpha \in S : \omega(\alpha) = n\}|}_{\# \text{ of elements of weight } n \text{ in } S} x^n.$$

Equivalently,

$$[x^k] \Phi_S^\omega(x) = |\{\alpha \in S : \omega(\alpha) = k\}|.$$

Proof. Let $S_n = \{\alpha \in S : \omega(\alpha) = n\}$. Note that $S = \bigcup_{n \geq 0} S_n$. Then,

$$\Phi_S^\omega(x) = \sum_{\alpha \in S} x^{\omega(\alpha)} = \sum_{n \geq 0} \sum_{\alpha \in S_n} x^{\omega(\alpha)} = \sum_{n \geq 0} \sum_{\alpha \in S_n} x^n = \sum_{n \geq 0} |S_n| x^n.$$

□

Example. Write the generating series for the set S of all binary strings with respect to the weight function ω which records the length of the string.

Proof. We have

$$\begin{aligned} \Phi_S^\omega(x) &= \sum_{\alpha \in S} x^{\omega(\alpha)} \\ &= \sum_{n \geq 0} |\{\alpha \in S : \omega(\alpha) = n\}| x^n = \sum_{n \geq 0} (\# \text{ of binary strings of length } n) x^n \\ &= \sum_{n \geq 0} 2^n x^n \\ &= \sum_{n \geq 0} (2x)^n = \frac{1}{1 - 2x}. \end{aligned}$$

□

Example. Write the generating series for the set

$$S = \{\text{subsets of } \{1, 2, \dots, t\}\}$$

with the weight function $\omega(\alpha) = |\alpha|$.

Proof. We have

$$\begin{aligned} \Phi_S^\omega(x) &= \sum_{\alpha \subseteq \{1, 2, \dots, t\}} x^{|\alpha|} \\ &= \sum_{n \geq 0} (\# \text{ of subsets of size } n) x^n \\ &= \sum_{n \geq 0} \binom{t}{n} x^n \\ &= (1 + x)^t. \end{aligned}$$

□

2.3 Sum and Product Lemmas

Example. Consider breakfast foods and a weight function that corresponds to their cost. Let

$$S_1 = \{\text{omelette, waffles, pancakes, eggs, cereal}\},$$

and $\omega_1 : S_1 \rightarrow \mathbb{N}$ be defined as follows:

$$\omega_1(\text{omelette}) = 10, \quad \omega_1(\text{waffles}) = 10, \quad \omega_1(\text{pancakes}) = 8, \quad \omega_1(\text{eggs}) = 8, \quad \omega_1(\text{cereal}) = 5.$$

The generating series for breakfast main dishes with respect to cost is

$$\Phi_{S_1}^{\omega_1}(x) = \sum_{\alpha \in S_1} x^{\omega_1(\alpha)} = x^{10} + x^{10} + x^8 + x^8 + x^5 = 2x^{10} + 2x^8 + x^5.$$

Equivalently,

$$\Phi_{S_1}^{\omega_1}(x) = \sum_{n \geq 0} |\{\alpha \in S_1 : \omega_1(\alpha) = n\}| x^n = 2x^{10} + 2x^8 + x^5,$$

where the exponents are the costs and the coefficients are the # of things of that cost.

Let

$$S_2 = \{\text{bacon, hashbrowns, toast}\},$$

and $\omega_2 : S_2 \rightarrow \mathbb{N}$ be defined as follows:

$$\omega_2(\text{bacon}) = 5, \quad \omega_2(\text{hashbrowns}) = 4, \quad \omega_2(\text{toast}) = 3.$$

So the generating series for S_2 with respect to cost is

$$\Phi_{S_2}^{\omega_2}(x) = \sum_{\alpha \in S_2} x^{\omega_2(\alpha)} = x^5 + x^4 + x^3.$$

Now, let's consider

$$S_1 \cup S_2 = \{\text{omelette, waffles, pancakes, eggs, cereal, bacon, hashbrowns, toast}\}.$$

The generating series for any breakfast food (main or side) with respect to cost is

$$\Phi_{S_1 \cup S_2}^\omega(x) = \sum_{\alpha \in S_1 \cup S_2} x^{\omega(\alpha)} = 2x^{10} + 2x^8 + x^5 + x^5 + x^4 + x^3 = 2x^{10} + 2x^8 + 2x^5 + x^4 + x^3,$$

Observe that $\Phi_{S_1 \cup S_2}^\omega(x) = \Phi_{S_1}^{\omega_1}(x) + \Phi_{S_2}^{\omega_2}(x)$.

Note. In this case, S_1 and S_2 are disjoint.

Lecture 8, 2025/09/19

Now, consider

$$\begin{aligned} \Phi_{S_1}^{\omega_1}(x) \cdot \Phi_{S_2}^{\omega_2}(x) &= (2x^{10} + 2x^8 + x^5)(x^5 + x^4 + x^3) \\ &= 2x^{15} + 2x^{14} + 2x^{13} + 2x^{13} + 2x^{12} + 2x^{11} + x^{10} + x^9 + x^8 \\ &= 2x^{15} + 2x^{14} + 4x^{13} + 2x^{12} + 2x^{11} + x^{10} + x^9 + x^8. \end{aligned}$$

For example, $2x^{12}$ indicates that there are two (main, side) combos with total price 12. So this is also the generating series for the cartesian product $S_1 \times S_2$ of breakfast combos with respect to cost. This can be denoted as

$$\Phi_{S_1 \times S_2}^{\omega'}(x)$$

where $\omega'(\alpha_1, \alpha_2) = \omega_1(\alpha_1) + \omega_2(\alpha_2)$.

Lemma 2.5 (Sum Lemma).

Let S_1 and S_2 be disjoint sets and let ω be a weight function on $S_1 \cup S_2$. Then,

$$\Phi_{S_1 \cup S_2}^\omega(x) = \Phi_{S_1}^\omega(x) + \Phi_{S_2}^\omega(x).$$

Proof. The main idea is that sums correspond to disjoint unions.

$$\begin{aligned} \Phi_{S_1}(x) + \Phi_{S_2}(x) &= \sum_{\alpha \in S_1} x^{\omega(\alpha)} + \sum_{\alpha \in S_2} x^{\omega(\alpha)} \\ &= \sum_{\alpha \in S_1 \cup S_2} x^{\omega(\alpha)} && \text{since } S_1 \text{ and } S_2 \text{ are disjoint} \\ &= \Phi_{S_1 \cup S_2}^\omega(x). && \square \end{aligned}$$

Lemma 2.6 (Infinite Sum Lemma).

Let $S_0, S_1, S_2, \dots$ be disjoint sets with union S and let ω be a weight function on S . Then,

$$\Phi_S^\omega(x) = \sum_{n \geq 0} \Phi_{S_n}^\omega(x).$$

Proof. Similar to the proof of the Sum Lemma. □

Lemma 2.7 (Product Lemma).

Let S_1 and S_2 be sets and let ω_1 and ω_2 be weight functions on S_1 and S_2 respectively. Then,

$$\Phi_{S_1 \times S_2}^\omega(x) = \Phi_{S_1}^{\omega_1}(x) \cdot \Phi_{S_2}^{\omega_2}(x)$$

where ω is a weight function on $S_1 \times S_2$ defined by

$$\omega(\alpha, \beta) = \omega_1(\alpha) + \omega_2(\beta).$$

Proof. The main idea is that products correspond to cartesian products.

$$\begin{aligned} \Phi_{S_1}^{\omega_1}(x) \cdot \Phi_{S_2}^{\omega_2}(x) &= \sum_{\alpha_1 \in S_1} x^{\omega_1(\alpha_1)} \cdot \sum_{\alpha_2 \in S_2} x^{\omega_2(\alpha_2)} \\ &= \sum_{\alpha_1 \in S_1} \sum_{\alpha_2 \in S_2} x^{\omega_1(\alpha_1) + \omega_2(\alpha_2)} \\ &= \sum_{(\alpha_1, \alpha_2) \in S_1 \times S_2} x^{\omega(\alpha_1, \alpha_2)} \\ &= \Phi_{S_1 \times S_2}^\omega(x). \end{aligned}$$

□

Example. Let $S = \{2, 4, 6, \dots\}$. How many pairs (a, b) are there with $a, b \in S$, and $a + b = 50$?

Proof. Let $\omega : S \rightarrow \mathbb{N}$ be defined by $\omega(\alpha) = \alpha$ and let $\omega' : S \times S \rightarrow \mathbb{N}$ be defined by

$\omega'(\alpha, \beta) = \alpha + \beta$. Note that we are looking for $[x^{50}]\Phi_{S \times S}^{\omega'}(x)$.

$$\begin{aligned}
[x^{50}]\Phi_{S \times S}^{\omega'}(x) &= [x^{50}](\Phi_S^{\omega}(x))^2 && \text{by Product Lemma} \\
&= [x^{50}]\left(\sum_{\alpha \in S} x^{\omega(\alpha)}\right)^2 \\
&= [x^{50}](x^2 + x^4 + x^6 + \dots)^2 \\
&= [x^{50}](x^2(1 + x^2 + x^4 + \dots))^2 \\
&= [x^{50}]\frac{x^4}{(1 - x^2)^2} \\
&= [x^{46}]\frac{1}{(1 - x^2)^2} && \text{by coefficient extraction rule (2)} \\
&= [x^{46}]\sum_{n \geq 0} \binom{n+2-1}{2-1}(x^2)^n && \text{by Negative Binomial Theorem} \\
&= [x^{46}]\sum_{n \geq 0} (n+1)x^{2n} \\
&= 23 + 1 && \text{taking } n = 23 \\
&= 24. && \square
\end{aligned}$$

Definition 2.7 (Star Operator).

Let A be a set. Define

$$A^* = \bigcup_{k \geq 0} A^k = \{\text{all tuples (of any length) of elements of } A\}.$$

Example. $\{0, 1\}^* = \{\underbrace{()}_{A^0}, \underbrace{(0), (1)}_{A^1}, \underbrace{(0, 0), (0, 1), (1, 0), (1, 1)}_{A^2}, \underbrace{(0, 0, 0), \dots}_{A^3, \dots}\}.$

Example. $\{1, 2, 3, \dots\}^* = \{\underbrace{()}_{A^0}, \underbrace{(1), (2), (3), \dots}_{A^1}, \underbrace{(1, 1), (1, 2), \dots}_{A^2}, \underbrace{(1, 1, 1), \dots}_{A^3, \dots}\}.$

Given a weight function ω on A , we can define ω^* on A^* such that

$$\omega^*((\alpha_1, \dots, \alpha_k)) = \sum_{i=1}^k \omega(\alpha_i).$$

This is a well-defined weight function provided no elements of A have weight 0 (otherwise, there would be infinitely many tuples of any given weight).