Logistic Regression

• Hypothesis Representation

Want

$$0 \le h_{\theta}(x) \le 1$$

Sigmoid function:

$$g(z) = \frac{1}{1 + e^{-z}}$$

Logistic function:

$$h_{\theta}(x) = g(\theta^T x) = \frac{1}{1 + e^{-\theta^T x}}$$

Interpretation:

 $h_{\theta}(x) = estimated probability that y = 1 on input x$

Decision Boundary

$$h_{\theta}(x) = g(\theta^T x)$$

$$g(z) = \frac{1}{1 + e^{-z}}$$

Suppose

$$predict y = 1, if h_{\theta}(x) \ge 0.5$$

predict
$$y = 0$$
, if $h_{\theta}(x) < 0.5$

Decision boundary

$$\theta^T x = 0$$

Non-linear decision boundaries

$$h_{\theta}(x) = g(-1 + x_1^2 + x_2^2)$$

$$Predict \ y = 1, if - 1 + x_1^2 + x_2^2 \ge 0$$

$$Predict \ y = 0, if - 1 + x_1^2 + x_2^2 < 0$$

Decision boundary

$$x_1^2 + x_2^2 = 1$$

Cost function

$$J(\theta) = \frac{1}{m} \sum_{i=1}^{m} Cost(h_{\theta}(x^{(i)}), y^{(i)})$$

$$Cost(h_{\theta}(x), y) = \begin{cases} -\log(h_{\theta}(x)), & \text{if } y = 1\\ -\log(1 - h_{\theta}(x)), & \text{if } y = 0 \end{cases}$$

$$\to J(\theta) = -\frac{1}{m} \sum_{i=1}^{m} (y^{(i)} \log(h_{\theta}(x^{(i)})) + (1 - y^{(i)}) \log(1 - h_{\theta}(x^{(i)}))$$

To filter parameters θ :

$$minJ(\theta)$$

To make a prediction given new x:

Output
$$h_{\theta}(x) = \frac{1}{1 + e^{-\theta^T x}}$$

Gradient descent

$$J(\theta) = -\frac{1}{m} \sum_{i=1}^{m} (y^{(i)} \log \left(h_{\theta}(x^{(i)}) \right) + \left(1 - y^{(i)} \right) \log \left(1 - h_{\theta}(x^{(i)}) \right))$$

Want $min_{\theta}J(\theta)$:

Repeat {

$$\theta_j \coloneqq \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta)$$

$$= \theta_j - \alpha \frac{1}{m} \sum_{i=1}^{m} (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

}

• One-vs-all

Train a logistic regression classifier $h_{\theta}^{(i)}(x)$ for each class i to predict the probability that y=i.

On a new input x, to make a prediction, pick the class i that maximizes

$$maxh_{\theta}^{(i)}(x)$$