abbvie

Immunology

Competitive Intelligence Update

Brought to you by:
APEX & IMABI CI

Takeda Announced Positive Results from the Ph3 KEPLER Trial of Entyvio in Pediatric Patients with Moderately to Severely Active UC

Company: Takeda
Drug: Entyvio

Date: February 19, 2026
Source(s): press release

What happened:

- Takeda announced positive results from the pivotal Ph3 KEPLER trial of Entyvio (vedolizumab, α4β7 integrin inhibitor, **IV/IV**) in **pediatric patients aged 2 years and older with moderately to severely active ulcerative colitis**

- Results presented at the ECCO 2026 (abstract link)

Impact (CI PoV): MEDIUM

- The positive data strengthen Takeda's position in the IBD market and bolster its strategy to expand Entyvio's indications. With this approval, Takeda addresses a segment where therapeutic options remain limited.

- Takeda is moving toward regulatory filings in the US, EU, and other markets for Entyvio IV in pediatric patients aged 2–17 years with moderately to severely UC

Why it matters

- The KEPLER trial results represent a significant clinical milestone for Takeda, demonstrating that Entyvio can effectively induce and maintain clinical remission in pediatric patients as young as two years old. These results address a critical treatment gap for children and adolescents with moderate to severe UC
- Entyvio is currently positioned as the #1 prescribed brand in IBD in the US; due to its tolerable safety profile, Entyvio could be preferred over anti-TNFs currently approved in the pediatric population

Tell me more:

- The KEPLER study enrolled 120 pediatric patients aged 2 to 17 years with moderately to severely active UC, who had shown inadequate response to conventional treatments, including steroids and immunomodulators, and/or TNF antagonists
- Patients initially received open-label IV vedolizumab during a 14-week induction phase. At week 14, 93/ 120 patients achieved a clinical response, who were randomized into maintenance therapy groups
    - 47 patients were assigned to a low-dose vedolizumab regimen, and 46 patients were assigned to a high-dose vedolizumab regimen
- Among the 93 patients randomized to maintenance therapy,
    - 47.3% achieved the primary endpoint of clinical remission at week 54
    - 34.7% of patients reached clinical remission at week 14; 29% of participants achieved sustained clinical remission at both week 14 and week 54
- No new safety signals identified. The most common treatment-emergent adverse events (occurring in ≥10% of patients) were upper respiratory infection (30%), worsening of UC (17.5%), and pyrexia (12.5%)

Image from the data presented at ECCO 2026