

MACMAHON'S SUM-OF-DIVISORS FUNCTIONS, CHEBYSHEV POLYNOMIALS, AND QUASI-MODULAR FORMS

GEORGE E. ANDREWS AND SIMON C. F. ROSE

ABSTRACT. We investigate a relationship between MacMahon's generalized sum-of-divisors functions and Chebyshev polynomials of the first kind. This determines a recurrence relation to compute these functions, as well as proving a conjecture of MacMahon about their general form by relating them to quasi-modular forms. These functions arise as solutions to a curve-counting problem on Abelian surfaces.

1. INTRODUCTION

The sum-of-divisors function $\sigma_k(n)$ is defined to be

$$\sigma_k(n) = \sum_{d|n} d^k.$$

For $k = 1$, this has as a generating function

$$A_1(q) = \sum_{n=1}^{\infty} \sigma_1(n) q^n = \sum_{k=1}^{\infty} \frac{q^k}{(1-q^k)^2}.$$

As a generalization of this notion, MacMahon introduces in the paper [5, pp. 303, 309] the generating functions

$$A_k = \sum_{0 < m_1 < \dots < m_k} \frac{q^{m_1 + \dots + m_k}}{(1-q^{m_1})^2 \dots (1-q^{m_k})^2}$$

$$C_k = \sum_{0 < m_1 < \dots < m_k} \frac{q^{2m_1 + \dots + 2m_k - k}}{(1-q^{2m_1-1})^2 \dots (1-q^{2m_k-1})^2}.$$

These provide generalizations in the following sense.

Fix a positive integer k . We define $a_{n,k}$ to be the sum

$$a_{n,k} = \sum s_1 \dots s_k$$

where the sum is taken over all possible ways of writing $n = s_1 m_1 + \dots + s_k m_k$ with $0 < m_1 < \dots < m_k$. Note that for $k = 1$ this is nothing but $\sigma_1(n)$, the usual sum-of-divisors function. It can then be shown that we have

$$A_k(q) = \sum_{n=1}^{\infty} a_{n,k} q^n.$$

Similarly, we define $c_{n,k}$ to be

$$c_{n,k} = \sum_1 s_1 \dots s_k$$

where the sum is over all partitions of n into

$$n = s_1(2m_1 - 1) + \cdots + s_k(2m_k - 1)$$

with, as before $0 < m_1 < \cdots < m_k$. For $k = 1$ this is the sum over all divisors whose conjugate is an odd number. As for the case of $a_{n,k}$, we have

$$C_k(q) = \sum_{n=1}^{\infty} c_{n,k} q^n.$$

We recall also that Chebyshev polynomials [1, p. 101] are defined via the relation

$$T_n(\cos \theta) = \cos(n\theta).$$

With these we form the following generating functions.

$$F(x, q) := 2 \sum_{n=0}^{\infty} T_{2n+1}\left(\frac{1}{2}x\right) q^{n^2+n}$$

$$G(x, q) := 1 + 2 \sum_{n=1}^{\infty} T_{2n}\left(\frac{1}{2}x\right) q^{n^2}.$$

The results of this paper are the following.

Theorem 1. *We have the following equalities:*

$$F(x, q) = (q^2; q^2)_{\infty}^3 \sum_{k=0}^{\infty} A_k(q^2) x^{2k+1}$$

$$G(x, q) = \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} \sum_{k=0}^{\infty} C_k(q) x^{2k}$$

where $(a; q)_{\infty} = \prod_{k=0}^{\infty} (1 - aq^k)$.

Corollary 2. *The functions $A_k(q)$ and $C_k(q)$ can be written as*

$$A_k(q) = \frac{(-1)^k}{(2k+1)!(q; q)_{\infty}^3} \sum_{n=0}^{\infty} (-1)^n (2n+1) \frac{(n+k)!}{(n-k)!} q^{\frac{1}{2}n(n+1)}$$

$$C_k(q) = \frac{(-1)^k (-q; q)_{\infty}}{(2k)!(q; q)_{\infty}} \sum_{n=1}^{\infty} (-1)^n 2n \frac{(n+k-1)!}{(n-k)!} q^{n^2}.$$

Corollary 3. *The functions A_k and C_k satisfy the recurrence relations*

$$A_k(q) = \frac{1}{(2k+1)2k} \left((6A_1(q) + k(k-1))A_{k-1}(q) - 2q \frac{d}{dq} A_{k-1}(q) \right)$$

$$C_k(q) = \frac{1}{2k(2k-1)} \left((2C_1(q) + (k-1)^2)C_{k-1}(q) - q \frac{d}{dq} C_{k-1}(q) \right).$$

Our final result settles a long-standing conjecture of MacMahon. In MacMahon's paper [5, p. 328] he makes the claim

The function $A_k = \sum a_{n,k} q^n$ has apparently the property that the coefficient $a_{n,k}$ is expressible as a linear function of the sum of the uneven powers of the divisors of n . I have not succeeded in reaching the general theory...

What we prove is the following.

Corollary 4. *The functions $A_k(q)$ are in the ring of quasi-modular forms.*

We will also discuss in section 3 some applications of this result to an enumerative problem involving counting curves on abelian surfaces.

2. PROOFS

Proof of theorem 1. Beginning with the series $F(x, q)$, and letting $x = 2 \cos \theta$ we find

$$\begin{aligned} F(x, q) &= 2 \sum_{n=0}^{\infty} T_{2n+1}(\cos \theta) q^{n^2+n} \\ &= 2 \sum_{n=0}^{\infty} \cos((2n+1)\theta) q^{n^2+n} \\ &= \sum_{n=0}^{\infty} \left(e^{i(2n+1)\theta} + e^{-i(2n+1)\theta} \right) q^{n^2+n} \\ &= \sum_{n=0}^{\infty} e^{i(2n+1)\theta} q^{n^2+n} + \sum_{n=0}^{\infty} e^{-i(2n+1)\theta} q^{n^2+n} \end{aligned}$$

where in the latter sum, letting $n \mapsto -n-1$ we obtain

$$F(x, q) = e^{i\theta} \sum_{n=-\infty}^{\infty} e^{2ni\theta} q^{n^2+n}.$$

Using the Jacobi triple product [1, p. 497, Thm 10.4.1] we see that this is equal to

$$\begin{aligned} e^{i\theta} \sum_{n=-\infty}^{\infty} e^{2ni\theta} q^{n^2+n} &= e^{i\theta} (-e^{-2i\theta}; q^2)_{\infty} (-q^2 e^{2i\theta}; q^2)_{\infty} (q^2; q^2)_{\infty} \\ &= \underbrace{(e^{i\theta} + e^{-i\theta})}_x (q^2; q^2)_{\infty} \prod_{m=1}^{\infty} \underbrace{(q + 2 \cos(2\theta) q^{2m} + q^{4m})}_{x^2-2} \\ &= x (q^2; q^2)_{\infty} \prod_{m=1}^{\infty} ((1 - q^{2m})^2 + x^2 q^{2m}) \\ &= x (q^2; q^2)_{\infty}^3 \prod_{m=1}^{\infty} \left(1 + x^2 \frac{q^{2m}}{(1 - q^{2m})^2} \right) \\ &= (q^2; q^2)_{\infty}^3 \sum_{k=0}^{\infty} A_k(q^2) x^{2k+1} \end{aligned}$$

and thus comparing coefficients of x^{2k+1} yields the result.

We ply a similar trick for $G(x, k)$. In that case we have

$$\begin{aligned} G(x, q) &= 1 + 2 \sum_{n>0} T_{2n}(\cos \theta) q^{n^2} \\ &= 1 + 2 \sum_{n>0} \cos(2n\theta) q^{n^2} \\ &= \sum_{n=-\infty}^{\infty} e^{2ni\theta} q^{n^2} \end{aligned}$$

which, again, by the Jacobi triple product yields

$$\begin{aligned}
\sum_{n=-\infty}^{\infty} e^{2ni\theta} q^{n^2} &= (q^2; q^2)_{\infty} (-qe^{2i\theta}; q^2)_{\infty} (-qe^{-2i\theta}; q^2)_{\infty} \\
&= (q^2; q^2)_{\infty} \prod_{m=1}^{\infty} \underbrace{(1 + 2\cos(2\theta)q^{2m-1} + q^{4m-2})}_{x^2-2} \\
&= (q^2; q^2)_{\infty} \prod_{m=1}^{\infty} ((1 - q^{2m-1})^2 + x^2 q^{2m-1}) \\
&= \underbrace{(q^2; q^2)_{\infty} (q; q^2)_{\infty}^2}_{\frac{(q; q)_{\infty}}{(-q; q)_{\infty}}} \prod_{m=1}^{\infty} \left(1 + x^2 \frac{q^{2m-1}}{(1 - q^{2m-1})^2}\right) \\
&= \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} \sum_{k=0}^{\infty} C_k(q) x^{2k}
\end{aligned}$$

which completes the theorem. \square

To deduce Corollary 2, we begin by expanding the series $F(x, q)$ (and similarly, $G(x, q)$) in powers of x , i.e.

$$\begin{aligned}
F(x, q) &= x f_0(q) + x^3 f_1(q) + x^5 f_2(q) + \cdots + x^{2k+1} f_k(q) + \cdots \\
G(x, q) &= g_0(q) + x^2 g_1(q) + x^4 g_2(q) + \cdots + x^{2k} g_k(q) + \cdots
\end{aligned}$$

Now, it can be shown that the coefficients of x^{2k} in $2T_{2n}(\frac{1}{2}x)$ and of x^{2k+1} in $2T_{2n+1}(\frac{1}{2}x)$ are respectively given by

$$2n(-1)^{n-k} \frac{(n+k-1)!}{(n-k)!(2k)!} \quad (-1)^{n-k} (2n+1) \frac{(n+k)!}{(n-k)!(2k+1)!}$$

and thus we have

$$\begin{aligned}
f_k(q) &= \frac{(-1)^k}{(2k+1)!} \sum_{n=0}^{\infty} (-1)^n (2n+1) \frac{(n+k)!}{(n-k)!} q^{n^2+n} \\
g_k(q) &= \frac{(-1)^k}{(2k)!} 2 \sum_{n=1}^{\infty} (-1)^n n \frac{(n+k-1)!}{(n-k)!} q^{n^2}.
\end{aligned}$$

As theorem 1 implies that $f_k(q) = (q^2; q^2)_{\infty}^3 A_k(q^2)$ and $g_k(q) = \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} C_k(q)$, we see that Corollary 2 follows.

Next, letting

$$\begin{aligned}
f_0(q) &= \sum_{n=0}^{\infty} (-1)^n (2n+1) q^{n^2+n} = (q^2; q^2)_{\infty}^3 \\
g_0(q) &= 1 + 2 \sum_{n=1}^{\infty} (-1)^n q^{n^2} = \frac{(q; q)_{\infty}}{(-q; q)_{\infty}}
\end{aligned}$$

and defining the operators $D_\ell = q \frac{d}{dq} - \ell(\ell - 1)$ and $D'_\ell = q \frac{d}{dq} - (\ell - 1)^2$, we then have that

$$\begin{aligned} f_k(q) &= \frac{(-1)^k}{(2k+1)!} D_k \cdots D_1 f_0(q) \\ g_k(q) &= \frac{(-1)^k}{(2k)!} D'_k \cdots D'_1 g_0(q). \end{aligned}$$

From these formulae we note that the functions f_k , g_k satisfy the recursion relations

$$\begin{aligned} f_k(q) &= \frac{-1}{(2k+1)2k} \left(q \frac{d}{dq} - k(k-1) \right) f_{k-1}(q) \\ g_k(q) &= \frac{-1}{2k(2k-1)} \left(q \frac{d}{dq} - (k-1)^2 \right) g_{k-1}(q). \end{aligned}$$

Noting again that $f_k(q) = (q^2; q^2)_\infty^3 A_k(q^2)$ and $g_k(q) = \frac{(q; q)_\infty}{(-q; q)_\infty} C_k(q)$, we now obtain the recurrence relation of Corollary 3 between the functions $A_k(q)$ and $C_k(q)$.

Our final result requires a bit of explanation. It is well known that the ring of modular forms for the full modular group $\Gamma = PSL_2(\mathbb{Z})$ is the polynomial ring in the generators E_4 , E_6 , where

$$E_{2k}(q) = 1 + \frac{2}{\zeta(1-2k)} \sum_{n=1}^{\infty} \sigma_{2k-1}(n) q^n$$

are the Eisenstein series of weights $2k$. There are no modular forms of weight 2 for Γ , but $E_2 = 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^n$ is a *quasi*-modular form (See [4]).

The relevant fact for this paper is that the ring of all such objects (which contains the ring of modular forms as a subring) is the ring generated either by E_2 , E_4 , and E_6 , or by $q \frac{d}{dq}$ and by E_2 . Noting then that $A_1(q) = \frac{1-E_2(q)}{24}$, the recurrence relation from Corollary 3 implies that each $A_k(q)$ lies in this ring, and hence the conclusion follows.

3. APPLICATIONS

The functions $A_k(q)$ and $C_k(q)$ arise naturally in the following enumerative problem.

Let $A \subset \mathbb{P}^N$ be a generic polarized abelian surface. There are then a finite number of hyperplane sections which are hyperelliptic curves of geometric genus g and having $\delta = N - g + 2$ nodes. The number of such curves, $N(g, \delta)$ is independent of the choice of A and these numbers can be assembled into a generating function

$$F(x, u) = \sum_{g, \delta} N(g, \delta) x^{2g+2} u^{g+\delta-1}.$$

The coefficient of x^{2g+2} in F is given by a certain homogeneous polynomial of degree $g-1$ in the functions $A_k(u^4)$ and $C_k(u^2)$.

This formula is derived by relating hyperelliptic curves on A to genus zero curves on the Kummer surface $A/\pm 1$. The latter is computed using orbifold Gromov-Witten theory, the Crepant resolution conjecture [2] and the Yau-Zaslow formula [6] [3]. This will be described further in the second author's thesis.

REFERENCES

1. G. E. Andrews, R. Askey, and R. Roy, *Special functions*, vol. 71, pp. 101, 497, Cambridge University Press, 1999.
2. Jim Bryan and Tom Graber, *The crepant resolution conjecture*, Algebraic geometry—Seattle 2005. Part 1, Proc. Sympos. Pure Math., vol. 80, Amer. Math. Soc., Providence, RI, 2009, pp. 23–42. MR 2483931 (2009m:14083)
3. Jim Bryan and Naichung Conan Leung, *The enumerative geometry of K3 surfaces and modular forms*, J. Amer. Math. Soc. **13** (2000), no. 2, 371–410 (electronic). MR 1750955 (2001i:14071)
4. Masanobu Kaneko and Don Zagier, *A generalized Jacobi theta function and quasimodular forms*, The moduli space of curves (Texel Island, 1994), Progr. Math., vol. 129, Birkhäuser Boston, Boston, MA, 1995, pp. 165–172. MR 1363056 (96m:11030)
5. P. A. MacMahon, *Divisors of numbers and their continuations in the theory of partitions*, Reprinted: Percy A. MacMahon Collected Papers (G. Andrews, ed.), MIT Press, Cambridge, 1986, pp. 305–341.
6. Shing-Tung Yau and Eric Zaslow, *BPS states, string duality, and nodal curves on K3*, Nuclear Phys. B **471** (1996), no. 3, 503–512. MR 1398633 (97e:14066)

PENN STATE UNIVERSITY, USA

E-mail address: `andrews@math.psu.edu`

UNIVERSITY OF BRITISH COLUMBIA, CANADA

E-mail address: `scfr@math.ubc.ca`