

# CHARACTERIZING THE NUMBER OF $m$ -ARY PARTITIONS MODULO $m$

GEORGE E. ANDREWS, AVIEZRI S. FRAENKEL, AND JAMES A. SELLERS

ABSTRACT. Motivated by a recent conjecture of the second author related to the ternary partition function, we provide an elegant characterization of the values  $b_m(mn)$  modulo  $m$  where  $b_m(n)$  is the number of  $m$ -ary partitions of the integer  $n$  and  $m \geq 2$  is a fixed integer.

2010 Mathematics Subject Classification: 05A17, 11P83

Keywords: partition, congruence, generating function

## 1. INTRODUCTION

Congruences for partition functions have been studied extensively for the last century or so, beginning with the discoveries of Ramanujan [7]. In this note, we will focus our attention on congruence properties for the partition functions which enumerate restricted integer partitions known as  $m$ -ary partitions. These are partitions of an integer  $n$  wherein each part is a power of a fixed integer  $m \geq 2$ . Throughout this note, we will let  $b_m(n)$  denote the number of  $m$ -ary partitions of  $n$ .

As an example, note that there are five 3-ary partitions of  $n = 9$ :

$$\begin{aligned} 9, \quad & 3 + 3 + 3, \quad 3 + 3 + 1 + 1 + 1, \\ & 3 + 1 + 1 + 1 + 1 + 1 + 1, \quad 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 \end{aligned}$$

Thus,  $b_3(9) = 5$ .

In the late 1960s, Churchhouse [3, 4] initiated the study of congruence properties of binary partitions ( $m$ -ary partitions with  $m = 2$ ). By his own admission, he did so serendipitously. To quote Churchhouse [4], “It is however salutary to realise that the most interesting results were discovered because I made a mistake in a hand calculation!”

Within months, other mathematicians proved Churchhouse’s conjectures and proved natural extensions of his results. These included Rødseth [8] who extended Churchhouse’s results to include the functions  $b_p(n)$  where  $p$  is any prime as well as Andrews [2] and Gupta [5, 6] who proved that corresponding results also held for  $b_m(n)$  where  $m$  could be any integer greater than 1. As part of an infinite family of results, these authors proved that, for any  $m \geq 2$  and any nonnegative integer  $n$ ,  $b_m(m(mn - 1)) \equiv 0 \pmod{m}$ .

We now fast forward forty years. In 2012, the second author conjectured the following absolutely remarkable result related to the ternary partition function  $b_3(n)$ :

- For all  $n \geq 0$ ,  $b_3(3n)$  is divisible by 3 if and only if at least one 2 appears as a coefficient in the base 3 representation of  $n$ .

- Moreover,  $b_3(3n) \equiv (-1)^j \pmod{3}$  whenever no 2 appears in the base 3 representation of  $n$  and  $j$  is the number of 1s in the base 3 representation of  $n$ .

This conjecture is remarkable for at least two reasons. First, it provides a complete characterization of  $b_3(3n)$  modulo 3. Such **characterizations** in the world of integer partitions are rare. Secondly, the result depends on the base 3 representation of  $n$  and nothing else.

Just to “see” what the second author saw, let’s quickly look at some data related to this conjecture.

| $n$ | Base 3 Representation of $n$                 | $b_3(3n)$ | $b_3(3n) \pmod{3}$ |
|-----|--|-----------|--------------------|
| 1   | $1 \times 3^0$                               | 2         | 2                  |
| 2   | $2 \times 3^0$                               | 3         | 0                  |
| 3   | $0 \times 3^0 + 1 \times 3^1$                | 5         | 2                  |
| 4   | $1 \times 3^0 + 1 \times 3^1$                | 7         | 1                  |
| 5   | $2 \times 3^0 + 1 \times 3^1$                | 9         | 0                  |
| 6   | $0 \times 3^0 + 2 \times 3^1$                | 12        | 0                  |
| 7   | $1 \times 3^0 + 2 \times 3^1$                | 15        | 0                  |
| 8   | $2 \times 3^0 + 2 \times 3^1$                | 18        | 0                  |
| 9   | $0 \times 3^0 + 0 \times 3^1 + 1 \times 3^2$ | 23        | 2                  |
| 10  | $1 \times 3^0 + 0 \times 3^1 + 1 \times 3^2$ | 28        | 1                  |
| 11  | $2 \times 3^0 + 0 \times 3^1 + 1 \times 3^2$ | 33        | 0                  |
| 12  | $0 \times 3^0 + 1 \times 3^1 + 1 \times 3^2$ | 40        | 1                  |
| 13  | $1 \times 3^0 + 1 \times 3^1 + 1 \times 3^2$ | 47        | 2                  |
| 14  | $2 \times 3^0 + 1 \times 3^1 + 1 \times 3^2$ | 54        | 0                  |
| 15  | $0 \times 3^0 + 2 \times 3^1 + 1 \times 3^2$ | 63        | 0                  |

In recent days, the authors succeeded in proving this conjecture. Thankfully, the proof was both elementary and elegant. After just a bit of additional consideration, we were able to alter the proof to provide a completely unexpected generalization. We describe this generalized result, and provide its proof, in the next section.

## 2. THE FULL RESULT

Our main theorem, which includes the above conjecture in a very natural way, provides a complete characterization of  $b_m(mn)$  modulo  $m$ :

**Theorem 2.1.** *Let  $m \geq 2$  be a fixed integer and let*

$$n = a_0 + a_1m + \cdots + a_jm^j$$

*be the base  $m$  representation of  $n$  (so that  $0 \leq a_i \leq m-1$  for each  $i$ ). Then*

$$b_m(mn) \equiv \prod_{i=0}^j (a_i + 1) \pmod{m}.$$

Notice that the conjecture mentioned above is exactly the  $m = 3$  case of Theorem 2.1.

In order to prove Theorem 2.1, we need a few elementary tools. We describe these tools here.

First, it is important to note that the generating function for  $b_m(n)$  is given by

$$(1) \quad B_m(q) := \prod_{j=0}^{\infty} \frac{1}{1 - q^{mj}}.$$

Note that  $B_m(q)$  satisfies the functional equation

$$(1 - q)B_m(q) = B_m(q^m).$$

From here it is straightforward to prove that

$$b_m(mn) = b_m(mn + i)$$

for all  $1 \leq i \leq m - 1$ . Thus, we see that Theorem 2.1 actually provides a characterization of  $b_m(N) \pmod{m}$  for **all**  $N$ , not just for those  $N$  which are multiples of  $m$ .

With this information in hand, we now prove a small number of lemmas which we will use in our proof of Theorem 2.1.

**Lemma 2.2.** *For  $|x| < 1$ ,*

$$\frac{1 - x^m}{(1 - x)^2} \equiv \sum_{k=1}^m kx^{k-1} \pmod{m}.$$

*Proof.* This elementary congruence can be proven rather quickly using well-known mathematical tools. We begin with the geometric series identity

$$\frac{1}{1 - x} = \sum_{k=0}^{\infty} x^k.$$

Differentiating both sides yields

$$\frac{1}{(1 - x)^2} = \sum_{k=1}^{\infty} kx^{k-1}.$$

We then multiply both sides by  $1 - x^m$  and simplify as follows:

$$\begin{aligned} \frac{1 - x^m}{(1 - x)^2} &= \sum_{k=1}^{\infty} kx^{k-1} - x^m \sum_{k=1}^{\infty} kx^{k-1} \\ &= \sum_{k=1}^{\infty} kx^{k-1} - \sum_{k=m+1}^{\infty} (k - m)x^{k-1} \\ &= \sum_{k=1}^m kx^{k-1} + \sum_{k=m+1}^{\infty} mx^{k-1} \\ &\equiv \sum_{k=1}^m kx^{k-1} \pmod{m} \end{aligned}$$

■

**Lemma 2.3.** *Let  $\zeta$  be the  $m^{\text{th}}$  root of unity given by  $\zeta = e^{2\pi i/m}$ . Then*

$$\sum_{k=0}^{m-1} \frac{1}{1 - \zeta^k q} = m \left( \frac{1}{1 - q^m} \right).$$

*Proof.* Using geometric series and elementary series manipulations, we have

$$\begin{aligned}
\sum_{k=0}^{m-1} \frac{1}{1 - \zeta^k q} &= \sum_{k=0}^{m-1} \sum_{r=0}^{\infty} \zeta^{kr} q^r \\
&= \sum_{k=0}^{m-1} \left( \sum_{r \mid m} \zeta^{kr} q^r + \sum_{r \nmid m} \zeta^{kr} q^r \right) \\
&= \sum_{k=0}^{m-1} \sum_{j=0}^{\infty} \zeta^{k(jm)} q^{jm} + \sum_{k=0}^{m-1} \sum_{r \nmid m} \zeta^{kr} q^r \\
&= \sum_{k=0}^{m-1} \frac{1}{1 - q^m} \quad \text{using facts about roots of unity} \\
&= m \left( \frac{1}{1 - q^m} \right).
\end{aligned}$$

■

**Lemma 2.4.** *Let  $T_m(q) := \sum_{n \geq 0} b_m(mn)q^n$ . Then*

$$T_m(q) = \frac{1}{1 - q} B_m(q).$$

*Proof.* As in Lemma 2.3, let  $\zeta = e^{2\pi i/m}$ . Note that

$$\begin{aligned}
T_m(q^m) &= \sum_{n \geq 0} b_m(mn)q^{mn} \\
&= \frac{1}{m} (B_m(q) + B_m(\zeta q) + \cdots + B_m(\zeta^{m-1} q)) \\
&= \left( \prod_{j=1}^{\infty} \frac{1}{1 - q^{mj}} \right) \times \frac{1}{m} \sum_{k=0}^{m-1} \frac{1}{1 - \zeta^k q} \\
&= \frac{1}{1 - q^m} \prod_{j=1}^{\infty} \frac{1}{1 - q^{mj}}
\end{aligned}$$

thanks to Lemma 2.3. Lemma 2.4 then follows by replacing  $q^m$  by  $q$ . ■

We now combine these elementary facts from the lemmas above to prove one last lemma. This lemma will, in essence, allow us to “move” from considering  $T_m(q)$  modulo  $m$  to a new function modulo  $m$  which makes the result of Theorem 2.1 transparent.

**Lemma 2.5.** *Let  $U_m(q) = \prod_{j=0}^{\infty} (1 + 2q^{m^j} + 3q^{2m^j} + \cdots + mq^{(m-1)m^j})$ . Then*

$$T_m(q) \equiv U_m(q) \pmod{m}.$$

*Proof.* Lemma 2.5 will follow if we can prove that  $\frac{1}{T_m(q)} \cdot U_m(q) \equiv 1 \pmod{m}$ , and this will be our means of attack. Thankfully, this follows from a novel generating function manipulation which we demonstrate here. Using (1) and Lemma 2.4, we

know that

$$\begin{aligned}
& \frac{1}{T_m(q)} \cdot U_m(q) \\
&= (1-q)^2 \prod_{j=1}^{\infty} (1-q^{m^j}) \prod_{j=0}^{\infty} \left(1 + 2q^{m^j} + 3q^{2m^j} + \cdots + mq^{(m-1)m^j}\right) \\
&\equiv (1-q)^2 \prod_{j=1}^{\infty} (1-q^{m^j}) \prod_{j=0}^{\infty} \frac{1-q^{m^{j+1}}}{(1-q^{m^j})^2} \pmod{m} \quad \text{thanks to Lemma 2.2} \\
&= \frac{\prod_{j=0}^{\infty} 1-q^{m^{j+1}}}{\prod_{j=1}^{\infty} 1-q^{m^j}} \\
&= 1.
\end{aligned}$$

■

We can now utilize all of the above results to prove Theorem 2.1.

*Proof.* First, we remember that

$$\sum_{n \geq 0} b_m(mn)q^n = T_m(q) \equiv U_m(q) \pmod{m}.$$

So we simply need to consider  $U_m(q)$  modulo  $m$  to obtain our proof. Note that

$$U_m(q) = \prod_{j=0}^{\infty} \left(1 + 2q^{m^j} + 3q^{2m^j} + \cdots + mq^{(m-1)m^j}\right).$$

If we expand this product as a power series in  $q$ , then each term of the form  $q^n$  can occur at most once (because the terms  $q^{i \cdot m^j}$  are serving as the building blocks for the **unique** base  $m$  representation of  $n$ ). Thus, if

$$n = a_0 + a_1m + \cdots + a_jm^j,$$

then the coefficient of  $q^n$  in this expansion is

$$\prod_{i=0}^j (a_i + 1) \pmod{m}.$$

■

## REFERENCES

1. G. E. Andrews, *The Theory of Partitions*, Addison-Wesley, Reading 1976; reprinted, Cambridge University Press, Cambridge, 1984, 1998
2. G. E. Andrews, Congruence properties of the  $m$ -ary partition function, *J. Number Theory* **3** (1971), 104-110
3. R. F. Churchhouse, Congruence properties of the binary partition function, *Proc. Cambridge Philos. Soc.* **66** (1969), 371-376
4. R. F. Churchhouse, Binary partitions, in *Computers in Number Theory*, edited A. O. L. Atkin and B. J. Birch, Academic Press, London, 1971
5. H. Gupta, A simple proof of the Churchhouse conjecture concerning binary partitions, *Indian J. Pure Appl. Math.* **3** (1972), no. 5, 791-794
6. H. Gupta, On  $m$ -ary partitions, *Proc. Cambridge Philos. Soc.* **71** (1972), 343-345
7. S. Ramanujan, Some properties of  $p(n)$ , the number of partitions of  $n$ , *Proc. Cambridge Philos. Soc.* **19** (1919), 207-210
8. Ø. Rødseth, Some arithmetical properties of  $m$ -ary partitions, *Proc. Cambridge Philos. Soc.* **68** (1970), 447-453

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA,  
ANDREWS@MATH.PSU.EDU

DEPARTMENT OF COMPUTER SCIENCE AND APPLIED MATHEMATICS, WEIZMANN INSTITUTE OF  
SCIENCE, 76100 REHOVOT, ISRAEL, AVIEZRI.FRAENKEL@WEIZMANN.AC.IL

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA,  
SELLERSJ@PSU.EDU