

THE LANCET

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Moskowitz CH, Nademanee A, Masszi T, et al, for the AETHERA Study Group. Brentuximab vedotin as consolidation therapy after autologous stem-cell transplantation in patients with Hodgkin's lymphoma at risk of relapse or progression (AETHERA): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2015; published online March 19. [http://dx.doi.org/10.1016/S0140-6736\(15\)60165-9](http://dx.doi.org/10.1016/S0140-6736(15)60165-9).

Web Extra Material

Table of Contents

AETHERA Study Group2

Table S1: Dose Modification Schedule3

Table S2: PFS Analysis Components4

Table S3: Subsequent Anti-Tumor Therapies5

Table S4: Serious Adverse Events Occurring in ≥ 2 Patients on the BV Arm6

Table S5: Summary of Deaths7

Figure S1: Progression-Free Survival by Response to Frontline Therapy8

AETHERA Study Group

Bulgaria

Dobrin Konstantinov

Georgi Mihaylov

Czech Republic

Jan Koren

Jana Markova

Pavla Stepankova

France

Pauline Brice

Noel Milpied

Gilles Salles

Aspasia Stamatoullas

Germany

Bastian von Tresckow

Hungary

Zita Borbenyi

Arpad Illes

Tamas Masszi

Arpad Szomor

Italy

Angelo Carella

Alessandro M. Gianni

Pier Luigi Zinzani

Poland

Andrzej Hellmann

Jerzy Holowiecki

Iwona Hus

Wieslaw Jedrzejczak

Slawomira Kyrz-Krzemien

Tadeusz Robak

Aleksander Skotnicki

Jan Walewski

Krzysztof Warzocha

Romania

Istvan Benedek

Andreea Delia Moicean

Alina Daniela Tanase

Russia

Boris Afanasjev

Julia Alexeeva

Anait Davtyan

Tatiana Konstantinova

Sergey Kravchenko

Irina Kryuchkova

Aleksandr Myasnikov

Dzhelil Osmanov

Elleonora Podoltseva

Oleg Rukavitsyn

Tatiana Shneider

Serbia

Slobodan Marjanovic

Aleksandar Savic

Milena Todorovic

Spain

Javier Briones

Monica Estevez

Ramon Garcia-Sanz

Carmen Martinez

Mercedes Rodriguez-Calvillo

United Kingdom

Catherine Burton

Roderick Johnson

John Radford

United States

Muneer H. Abidi

Edward Agura

Weiyun Z. Ai

Paolo Anderlini

Sally Arai

Ernesto Ayala

Veronika Bachanova

Asad Bashey

Amanda Cashen

Andy I. Chen

Steven M. Devine

Suzanne Fanning

Tatyana Feldman

Jonathan W. Friedberg

Brian T. Hill

Leona Holmberg

Eric Jacobsen

Mark S. Kaminski

Yvette Kasamon

Mary Ellen Martin

Jeffrey Matous

John M. McCarty

Craig H. Moskowitz

Auayporn Nademanee

Thomas Shea

Gary Spitzer

Patrick Stiff

Jasmine Zain

Table S1: Dose Modification Schedule

Toxicity	Grade 1	Grade 2	Grade 3	Grade 4
Non-hematologic	Continue at same dose level.	Continue at same dose level, except in the event of Grade 2 neuropathy. For Grade 2 neuropathy, withhold dose until toxicity is \leq Grade 1 or has returned to baseline, then reduce the dose to 1.2 mg/kg and resume treatment.	Withhold dose until toxicity is \leq Grade 1 or has returned to baseline, then resume treatment at the same dose level ^a . For Grade 3 or higher neuropathy, discontinue treatment.	Withhold dose until toxicity is \leq Grade 1 or has returned to baseline, then reduce dose to 1.2 mg/kg and resume treatment ^a .
Hematologic	Continue at same dose level.	Continue at same dose level.	Withhold dose until toxicity is \leq Grade 2, or has returned to baseline, then resume treatment at the same dose level ^b . Consider growth factor support (G-CSF or GM-CSF) for treatment of neutropenia and prophylaxis in subsequent cycles.	Withhold dose until toxicity is \leq Grade 2, then resume treatment at the same dose level. Consider growth factor support (G-CSF or GM-CSF) for treatment of neutropenia and prophylaxis in subsequent cycles. For the second occurrence of Grade 4 toxicity (if neutropenia, while receiving growth factor support), withhold dose until toxicity is \leq Grade 2, then reduce the dose to 1.2 mg/kg and resume treatment ^b .

^a Patients who develop Grade 3 or 4 electrolyte laboratory abnormalities may continue study treatment without interruption.

^b Patients who develop Grade 3 or 4 lymphopenia may continue study treatment without interruption. G-CSF: granulocyte-colony stimulating factor; GM-CSF: granulocyte macrophage-colony stimulating factor

Table S2: PFS Analysis Components

Analysis	CT scans (per IRF)	CT Scans (per investigator)	Biopsy Reports	Lymphoma Assessments	Death
IRF	X		X		X
Investigator		X	X	X	X

Table S3: Subsequent Anti-Tumor Therapies

	Treatment Group	
	BV (n=165)	Placebo (n=164)
Subsequent treatment ever received, n (%)	51 (31)	85 (52)
Stem cell transplant*	13 (8)	24 (15)
Single-agent BV	8 (5)	72 (44)
Multi-agent therapy including BV	1 (1)	1 (1)
Multi-agent therapy	35 (21)	34 (21)
Single-agent therapy	22 (13)	22 (13)
Radiation	22 (13)	23 (14)
Donor lymphocyte infusion	2 (1)	1 (1)
Other treatment	1 (1)	2 (1)
First subsequent treatment, n (%)		
Allogeneic stem cell transplant	3 (2)	0 (0)
Single-agent BV	3 (2)	63 (38)
Multi-agent therapy including BV	1 (1)	0 (0)
Multi-agent therapy	27 (16)	12 (7)
Single-agent therapy	7 (4)	5 (3)
Radiation	10 (6)	4 (2)
Other treatment	0 (0)	1 (1)

* Transplant type: BV: 12/13 allogeneic; placebo: 23/24 allogeneic

BV: brentuximab vedotin

Table S4: Serious Adverse Events Occurring in ≥ 2 Patients on the BV Arm

Preferred Term	Treatment Group	
	BV (n=167)	Placebo (n=160)
Any Event, n (%)	41 (25)	20 (13)
Pneumonia	7 (4)	4 (3)
Pyrexia	6 (4)	2 (1)
Vomiting	5 (3)	1 (1)
Nausea	4 (2)	1 (1)
Hepatotoxicity	3 (2)	1 (1)
Peripheral sensory neuropathy	3 (2)	0 (0)
Acute respiratory distress syndrome	2 (1)	1 (1)
Constipation	2 (1)	0 (0)
Headache	2 (1)	0 (0)
Herpes zoster	2 (1)	1 (1)
Pneumonitis	2 (1)	0 (0)

BV: brentuximab vedotin


Table S5: Summary of Deaths

	BV (n=167)	Placebo (n=160)
All Deaths, n (%)	28 (17)	25 (16)
Disease Related	18 (11)	17 (11)
Acute respiratory distress syndrome	0	1 (1)
Disease progression	5 (3)	9 (6)
Hodgkin lymphoma	13 (8)	7 (4)
Not Disease Related	9 (5)	7 (4)
Acute respiratory distress syndrome*	2 (1)	0
Aplastic anemia	0	1 (1)
Bladder cancer	1 (1)	0
Cardiac arrest	1 (1)	0
Graft versus host disease	0	3 (2)
Influenza	0	1 (1)
Lung infection	1 (1)	0
Myelodysplastic syndrome	1 (1)	1 (1)
Myocardial infarction	1 (1)	0
Pancreatic carcinoma	1 (1)	0
Pneumonia	0	1 (1)
Sepsis	1 (1)	0
Disease Relationship Unknown	1 (1)	1 (1)
Fungal pneumonia	0	1 (1)
Other	1 (1)	0
Death prior to progression per IRF, n (%)	4 (2)	3 (2)
Death prior to progression by INV, n (%)	5 (3)	3 (2)

*1 death occurred within 30 days of last dose of study treatment

BV: brentuximab vedotin; IRF: independent review facility; INV: investigator

Figure S1: Progression-Free Survival by Response to Frontline Therapy



Progression-free survival (PFS) by response to frontline therapy. Kaplan-Meier plots showing PFS per investigator between the brentuximab vedotin (BV) arm (dark blue) and the placebo arm (light blue) by response to frontline therapy: refractory (top), relapse in less 12 months (middle), and relapse at least 12 months after frontline therapy and with extranodal involvement (bottom). Symbols indicate censored patients.