

# ASTRO C207 Radiative Processes in Astrophysics

## Fall 2021

## Problem Set 2

### 1. Blackbody Flux

$$\begin{aligned}
 B_\nu &= \frac{2h\nu^3}{c^2} \frac{1}{e^{\frac{h\nu}{k_B T}} - 1} \\
 \int_0^\infty \int_0^{2\pi} \int_0^{\pi/2} B_\nu \cos \theta \sin \theta d\theta d\phi d\nu &= 2\pi \int_0^{\pi/2} \cos \theta \sin \theta d\theta \int_0^\infty B_\nu d\nu \\
 &= 2\pi \cdot \frac{1}{2} \cdot \int_0^\infty \frac{2h\nu^3}{c^2} \frac{1}{e^{\frac{h\nu}{k_B T}} - 1} d\nu \\
 &= \pi \int_0^\infty \frac{2h\frac{c^3}{\lambda^3}}{c^2} \frac{1}{e^{\frac{hc}{\lambda k_B T}} - 1} d\nu \\
 &= \pi \int_0^\infty \frac{2hc^2}{\lambda^5} \frac{1}{e^{\frac{hc}{\lambda k_B T}} - 1} d\lambda \leftarrow \begin{cases} \nu = \frac{c}{\lambda} \\ d\nu = -\frac{c}{\lambda^2} d\lambda \end{cases} \\
 &= \pi \cdot 2hc^2 \cdot \frac{hc}{k_B T} \cdot \left( \frac{k_B T}{hc} \right)^5 \int_0^\infty \frac{1}{x(e^{1/x} - 1)} dx \leftarrow \begin{cases} x = \lambda \frac{k_B T}{hc} \\ dx = d\lambda \frac{k_B T}{hc} \end{cases} \\
 &= \pi \cdot \frac{2k_B^4 T^4}{h^3 c^2} \cdot \frac{\pi^4}{15} \\
 &= \frac{2\pi^5 k_B^4}{15h^3 c^2} T^4 \\
 &= \sigma T^4
 \end{aligned}$$

## 2. Flat Disks

For an annulus in the ring at distance  $r$ , power in must equal power out

$$\begin{aligned} P_{\text{in}} &= P_{\text{out}} \\ F_{\text{in}} &= F_{\text{out}} \\ &= \sigma T^4 \end{aligned}$$

where  $T$  is the temperature in question. For the flux coming from the star where  $\tan \theta_c = \frac{R_*}{r}$ ,

$$\begin{aligned} F_{\text{in}} &\approx F_* \cdot 4\pi R_*^2 \cdot \frac{1}{4\pi r^2} \cdot C \sin \theta_c \\ &\approx \sigma T_*^4 \frac{R_*^2}{r^2} \frac{R_*}{r} \end{aligned}$$


where  $C$  (roughly 1) accounts for error from  $r - R_* \approx r$  and the fact that there should be an integral over  $\theta$  going from 0 to  $+\theta_c$ .

$$\begin{aligned} \sigma T^4 &\approx \sigma T_*^4 \frac{R_*^2}{r^2} \frac{R_*}{r} \\ T &\approx T_* \left( \frac{R_*}{r} \right)^{\frac{3}{4}} \end{aligned}$$

This feels weird, but I guess as you get further away a slice of the ring of the same area occupies a smaller solid angle from the view of some  $z \neq 0$  point on the star?

### 3. A Simplified Greenhouse Effect

I suppose we'll assume the atmosphere doesn't emit as a blackbody and Earth doesn't reflect anything coming off of the sun?



$$F_{\text{Earth}} = F_{\text{scat}} + F_{\text{sun}}$$

$$F_{\text{out}} = F_{\text{sun}}$$

For scattering, we consider the volume of scatterers with some arbitrary column area and depth  $z$ .

optically thin atmosphere

$$z = z_{\text{atm}}$$

optically thick atmosphere

$$z = \lambda_{\text{mfp}} = \frac{1}{n\sigma_v} = \frac{z_{\text{atm}}}{N\sigma_v}$$

Considering  $j_{v,\text{scat}} = n\sigma_{\text{scat}}J_v = \frac{N}{z_{\text{atm}}}\sigma_{\text{scat}}J_v$ ,

$$j_{v,\text{scat}}V_{\text{scat}} \propto \begin{cases} NJ_v & \text{optically thin} \\ J_v & \text{optically thick} \end{cases}$$

where  $J_v = KF_{\text{Earth}}$  for some scalar  $K$ .

$$F_{\text{Earth}} = F_{\text{scat}} + F_{\text{sun}}$$

$$\sigma T_{\text{gnd}}^4 = F_{\text{scat}} + F_{\text{sun}}$$

$$\sigma T_{\text{gnd}}^4 = \begin{cases} \sigma T_{\text{gnd}}^4 C_{\text{thin}} N + F_{\text{sun}} & \text{optically thin} \\ \sigma T_{\text{gnd}}^4 C_{\text{thick}} + F_{\text{sun}} & \text{optically thick} \end{cases}$$

where  $C_{\text{thick/thin}}$  are constants.

$$T_{\text{gnd}} \propto \begin{cases} \sqrt[4]{\frac{1}{1-C_{\text{thin}}N}} & \text{optically thin} \\ \text{constant} & \text{optically thick} \end{cases}$$