

SI 211: Numerical Analysis

Solution of Homework 1

Provided by 陈宏博

1. (a) $x = \pi$

Exact solution is: 0 and Matlab solution is: -1.54597494809480e-13.

So the numerical approximation error is: -1.54597494809480e-13.

The reason for this error is that π is stored with finite precision in the computers.

- (b) $f(x)$ is factorable, we can rewrite it as

$$\begin{aligned}a_0 &= x, \\a_1 &= 10^4 a_0 \\a_2 &= \sin(a_1) \\a_3 &= \frac{a_2}{a_0}\end{aligned}$$

In the worst case, the numerical errors associated with evaluating $f(x)$

$$\begin{aligned}\Delta a_0 &\approx \text{eps} \\ \Delta a_1 &\approx |10^4| * \Delta a_0 + \text{eps} \\ \Delta a_2 &\approx |\cos(a_1)| * \Delta a_1 + \text{eps} \\ \Delta a_3 &\approx \left| \frac{a_2}{a_0^2} \right| * \Delta a_0 + \left| \frac{1}{a_0} \right| * \Delta a_2 + \text{eps}\end{aligned}$$

We have error equals $\Delta a_3 \approx 0.04$ or $1\text{e-}2$ (calculated by MATLAB with respect to above steps).

2. (a) Function *diff* code in Matlab:

```
function r = newdiff(f,x,h)
    b = 2*h;
    r = (f(x+h)-f(x-h))/b;
end
```

- (b) Evaluate the derivative of the function $f(x)=\exp(x)$ at $x=0$, and $h \in [10^{-15}, 10^{-1}]$


```
for i = 1 : 15
    h(i) = 10^-i;
end
x = 0.0;
exact_value = 1.0;
f = @(x) exp(x);
for i = 1 : 15
    e(i)=abs(newdiff(f,x,h(i))-exact_value);
end
```

```

x = log10(h);
y = log10(e);
plot(x,y);hold on
plot(x,y,'r.','MarkerSize',10);


```

The result is shown in the figure 1



Interpret the result:

The numerical differentiation error is produced by numerical error and mathematical error together, the error dominated by h^2 when h is greater than h^* , in the other hand, the error dominated by $\frac{eps}{h}$ when h is less than h^* .



$$\begin{aligned}
 \text{error}(h) &\approx h^2 + \frac{eps}{h} \\
 \frac{d}{dh} \text{error}(h^*) &= 2h^* - \frac{eps}{h^{*2}} = 0 \\
 h^* &= \sqrt[3]{eps} \approx 10^{-5}
 \end{aligned}$$

3. Algorithmic differentiation

$$\begin{aligned}b_0 &= 1 \\b_1 &= -\sin(a_0) * b_0 \\b_2 &= a_1 * b_1 + b_1 * a_1 \\b_3 &= \cos(a_1) * b_1 \\b_4 &= a_2 * b_3 + b_2 * a_3 \\f'(x) &= b_4\end{aligned}$$

The derivative of $f(x)$ at $x = 0$ is: 0 using AD code, the actual derivative is 0, so the numerical error is 0, and the order of magnitude of the numerical error also is 0.

Julia Code

```
import Base.*
import Base.sin
import Base.cos
mutable struct ADV
    a
    b
end

function *(A::ADV,B::ADV)
    return ADV(A.a*B.a,A.a*B.b+A.b*B.a);
end

function sin(A::ADV)
    return ADV(sin(A.a),cos(A.a)*A.b)
end

function cos(A::ADV)
    return ADV(cos(A.a),-sin(A.a)*A.b)
end

function f(x)
    return sin(cos(x))*cos(x*x)
end
```

Get the Result

```
x = ADV(0,1)
f(x)
ADV(0.8414709848078965, -0.0)
```