




12.4 PARTIAL DERIVATIVES

NOTE: KNOW YOUR DERIVATIVES
(i.e. e^x , $\ln(x)$...
and product, chain
rules!

- First Degree Partial Derivative: for $f(x, y)$:
 - Calc f_x = Derivative of $f(x, y)$ treating all non- x var's as const.
 - Calc f_y = Derivative $f(x, y)$ where all non- y var's are const.
 - NOTE: $f_x = \frac{\partial f}{\partial x}$, $f_y = \frac{\partial f}{\partial y}$
- First Deg. P. D. Using Limits: for $f(x, y)$
 - $f_x = f_x(a, b) = \lim_{h \rightarrow 0} \frac{f(a+h, b) - f(a, b)}{h}$
 - $f_y = f_y(c, d) = \lim_{h \rightarrow 0} \frac{f(c, d+h) - f(c, d)}{h}$
- Second Degree P. D.'s: for function $f(x, y)$:
 - Perform FDPD for the first var, then again on result for second: f_{xy} would be f_x derived with y ...

12.5 DA CHAIN RULE

- f' for $f \in x \dots y$ where x is a function of another var, and so is y
 - $f' = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial t}$ 
- f' for $f(x, y)$ where $x(s, t)$, $y(s, t)$ 
 - $f'_s = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial s}$, $f'_t = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial t}$
- If $f(x, y, z)$: add another term, $\frac{\partial f}{\partial z}$, and chain rule
- solution process: evaluate $f'(\dots)$; then substitute
 $x =$, $y =$, $z = \dots$ and simplify
- Implicit Differentiation $\frac{dy}{dx} = -\frac{F_x}{F_y}$



12.6 GRADIENTS AND DIRECTIONAL DERIVATIVES

- Gradients: $\nabla f(x,y)$ = "gradient" = $\langle f_x, f_y \rangle$. If evaluated at a point, substitute x, y from point, solve as a vector.

- Directional Derivative:

- 1) Find if \vec{v} provided is unit vector, else make it one
- 2) Find $\nabla f(x,y)$ (see above for help there :))
- 3) Find $D_v f(x,y) = \vec{v} \cdot \nabla f(x,y) = v_x \cdot f_x + v_y \cdot f_y$
- 4) Substitute Point "P".

NOTE: STEPS 3 and 4 CAN BE DONE INTERCHANGEABLY

- Directions of steepest change, no change

→ Steepest ascent = $(\nabla f(x,y) / \|\nabla f(x,y)\|)$ AKA unit in dir of $\nabla f(x,y)$

→ Steepest descent = $-[\text{steepest ascent}]$

→ NO change = \oplus [swapped variables and one made neg, i.e. $\langle 8,1 \rangle \rightarrow \langle -1,8 \rangle$]
from pos steepest or neg steepest.

- Tangent Line to Level Curves: $f(x,y)$ at point $P = (a,b)$

→ $y'(x) = -\frac{f_x(a,b)}{f_y(a,b)}$: eval, if UNO = VERT TAN, ~~FE~~

→ CHECK: should be \perp to $\nabla f(x,y) \mid P \dots$