CH 13.1 - 11/6/2018 Lyen ( Red [13.1] Double Integrals over Rectangular Areas • To compute Ssof(x,4) dx, dy first evaluate Sof(x,4) dx, then substitute that into Se Tay) a dy where Fair) = SFLAIN) · Notation: R f(x,y) on {(x,y): a < x < b, c < y < d} just means: -> I fex, y) dy dx which is Equal to for fex, y) dx dy Average Function Value: Area of Rf(x,4) on {rectangle} Average Square distance: area of Rf(x,y) on (rectangle) where

f(x,y) = d² = (x,-x2)² + (y,-y2)² (Distance Function, Squared) 13.2 DOUBLE INTEGRALS OVER NON-RECTANGULAR AREAS · Writing a double integral based on a graph, or functions. > If you are doing dx dy the first integral is the y range -> the inside integral, the dx one, is from the "closer" function to the y axis to the "hither". In a Sodx · Evaluating an integral with equations as limits of Integration: just do as you would a normal integral · Example: Ry 3A bounded by X=1, -x-1, 2x+2 y dy dx

· Volume Calwatton using double Integrals: Integrate over the range of the base, the value of the top. to find the range of the base, with a plane, set z=0, solve for y = ....

· Changing the order of integration; Drawout The xy shape, and find the equations, then integrals, for the opposite

Variable. The passage as a complete of the moderation of

\* france marker value: then to " ( fray) or freehow u)

· Amage Space distance: mant. I then on freshandled about

a winter or course intering based on a stage, or functions.

e Evallation an integral with equations as limits of

\* Example: If y 3A bounded by xell -x-11, 2x+3