

Arithmetic of Fourier coefficients of Gan-Gurevich lifts on G_2

Naomi Sweeting

(joint work with Petar Bakic, Alex Horawa, and Siyan Daniel Li-Huerta)

June 10, 2025

International Seminar on Automorphic Forms

Introduction: modular forms on G_2

G = reductive group, $\pi = \otimes' \pi_v$ AR of $G(\mathbb{A})$.

Some π_∞ are more arithmetic than others.

Introduction: modular forms on G_2

G = reductive group, $\pi = \otimes' \pi_v$ AR of $G(\mathbb{A})$.

Some π_∞ are more arithmetic than others.

Most arithmetic: π_∞ is a **holomorphic discrete series**, e.g. for SL_2

Introduction: modular forms on G_2

G = reductive group, $\pi = \otimes' \pi_v$ AR of $G(\mathbb{A})$.

Some π_∞ are more arithmetic than others.

Most arithmetic: π_∞ is a **holomorphic discrete series**, e.g. for SL_2 , Sp_{2g} , ...

Introduction: modular forms on G_2

G = reductive group, $\pi = \otimes' \pi_v$ AR of $G(\mathbb{A})$.

Some π_∞ are more arithmetic than others.

Most arithmetic: π_∞ is a **holomorphic discrete series**, e.g. for SL_2 , Sp_{2g} , ...

You can use the relation to Shimura varieties to construct Galois representations, study Fourier expansions, and much more

Introduction: modular forms on G_2

G = reductive group, $\pi = \otimes' \pi_v$ AR of $G(\mathbb{A})$.

Some π_∞ are more arithmetic than others.

Most arithmetic: π_∞ is a **holomorphic discrete series**, e.g. for SL_2 , Sp_{2g} , ...

You can use the relation to Shimura varieties to construct Galois representations, study Fourier expansions, and much more

But not all groups admit holomorphic discrete series!

Introduction: modular forms on G_2

On G_2 , the **quaternionic discrete series** are the most arithmetic.

π_k : QDS representation of $G_2(\mathbb{R})$ of weight $k \geq 2$.

Introduction: modular forms on G_2

On G_2 , the **quaternionic discrete series** are the most arithmetic.

π_k : QDS representation of $G_2(\mathbb{R})$ of weight $k \geq 2$.

Definition

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ is a **quaternionic modular form** of weight $k \geq 2$ if φ generates the lowest K -type in π_k under $G_2(\mathbb{R})$.

Introduction: modular forms on G_2

On G_2 , the **quaternionic discrete series** are the most arithmetic.

π_k : QDS representation of $G_2(\mathbb{R})$ of weight $k \geq 2$.

Definition

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ is a **quaternionic modular form** of weight $k \geq 2$ if φ generates the lowest K -type in π_k under $G_2(\mathbb{R})$.

Fact

The Fourier coefficients $c_A(\varphi)$ of a cuspidal QMF φ of weight k are indexed by totally real cubic rings A (and these determine φ)

Introduction: modular forms on G_2

On G_2 , the **quaternionic discrete series** are the most arithmetic.

π_k : QDS representation of $G_2(\mathbb{R})$ of weight $k \geq 2$.

Definition

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ is a **quaternionic modular form** of weight $k \geq 2$ if φ generates the lowest K -type in π_k under $G_2(\mathbb{R})$.

Fact

The Fourier coefficients $c_A(\varphi)$ of a cuspidal QMF φ of weight k are indexed by totally real cubic rings A (and these determine φ)

- A free of rank 3 over \mathbb{Z} , $A \otimes \mathbb{R} = \mathbb{R}^3$ as a ring
- e.g. $A = \mathbb{Z}^3$, $A = \mathcal{O}_E$ with E/\mathbb{Q} totally real cubic field

Introduction: modular forms on G_2

On G_2 , the **quaternionic discrete series** are the most arithmetic.

π_k : QDS representation of $G_2(\mathbb{R})$ of weight $k \geq 2$.

Definition

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ is a **quaternionic modular form** of weight $k \geq 2$ if φ generates the lowest K -type in π_k under $G_2(\mathbb{R})$.

Fact*

The Fourier coefficients $c_A(\varphi)$ of a cuspidal QMF φ of weight k are indexed by totally real cubic rings A (and these determine φ)

- A free of rank 3 over \mathbb{Z} , $A \otimes \mathbb{R} = \mathbb{R}^3$ as a ring
- e.g. $A = \mathbb{Z}^3$, $A = \mathcal{O}_E$ with E/\mathbb{Q} totally real cubic field
- *: actually, only if φ has a nice level (like $\Gamma_0(N)$)

Fourier coefficients and arithmetic

The Fourier coefficients $c_A(\varphi)$ are arithmetic in nature.

Fourier coefficients and arithmetic

The Fourier coefficients $c_A(\varphi)$ are arithmetic in nature.

Some examples:

- There exists a natural family of Eisenstein series E_{2k} of weight $2k \geq 4$, such that

$$c_A(E_{2k}) = \zeta_A(1 - 2k)$$

when A is maximal. (Jiang-Rallis 97, Gan-Gross-Savin 02, Xiong 17)

Fourier coefficients and arithmetic

The Fourier coefficients $c_A(\varphi)$ are arithmetic in nature.

Some examples:

- There exists a natural family of Eisenstein series E_{2k} of weight $2k \geq 4$, such that

$$c_A(E_{2k}) = \zeta_A(1 - 2k)$$

when A is maximal. (Jiang-Rallis 97, Gan-Gross-Savin 02, Xiong 17)

- For $k \geq 6$, there is a basis of level one forms with all coefficients in \mathbb{Q}^{cyc} (Pollack 22)

Gross's Conjecture

Let f be a cusp form for $\mathrm{SL}_2(\mathbb{Z})$ of weight k .

Assuming $L(1/2, f) \neq 0$, there exists a cuspidal **Gan-Gurevich lift** φ of f to G_2

Conjecture (Gross)

For all maximal totally real cubic rings A ,

$$c_A(\varphi)^2 = L(1/2, f \otimes \rho_A) \mathrm{disc}(A)^{\frac{k-1}{2}}$$

where $\rho_A =$ two-dimensional Artin representation associated to A

Gross's Conjecture

Let f be a cusp form for $\mathrm{SL}_2(\mathbb{Z})$ of weight k .

Assuming $L(1/2, f) \neq 0$, there exists a cuspidal **Gan-Gurevich lift** φ of f to G_2

Conjecture (Gross)

For all maximal totally real cubic rings A ,

$$c_A(\varphi)^2 = L(1/2, f \otimes \rho_A) \mathrm{disc}(A)^{\frac{k-1}{2}}$$

where $\rho_A =$ two-dimensional Artin representation associated to A

- Kim-Yamauchi 24: true when $A = \mathbb{Z} \times \mathcal{O}_F$ for F/\mathbb{Q} quadratic

Gross's Conjecture

Let f be a cusp form for $\mathrm{SL}_2(\mathbb{Z})$ of weight k .

Assuming $L(1/2, f) \neq 0$, there exists a cuspidal **Gan-Gurevich lift** φ of f to G_2

Conjecture (Gross)

For all maximal totally real cubic rings A ,

$$c_A(\varphi)^2 = L(1/2, f \otimes \rho_A) \mathrm{disc}(A)^{\frac{k-1}{2}}$$

where $\rho_A =$ two-dimensional Artin representation associated to A

- Kim-Yamauchi 24: true when $A = \mathbb{Z} \times \mathcal{O}_F$ for F/\mathbb{Q} quadratic
- Today: Gross's conjecture when f is a CM form (always has level!)

Main result

- $f = f_\chi$ CM form of weight k , trivial character, and any level N , associated to K/\mathbb{Q} and $\chi : K^\times \backslash \mathbb{A}_K^\times \rightarrow \mathbb{C}^\times$.
- Assume: $L(1/2, \chi) \neq 0$.
- $\mathcal{A}_{GG}(f_\chi)$ space of “Gan-Gurevich lifts” $G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$.

Theorem (Bakic–Horawa–Li–Huerta–S., in progress)

For all $\ell|N$, fix a cubic ring A_ℓ/\mathbb{Z}_ℓ , such that

$$\prod_{\ell|N} \epsilon_\ell(A_\ell, \chi_\ell) = -\epsilon(1/2, \chi^3)$$

Then \exists a QMF $\varphi \in \mathcal{A}_{GG}(f_\chi)$ s.t. for A maximal outside N

$$|c_A(\varphi)|^2 = \begin{cases} L(1/2, f_\chi \otimes \rho_A) \operatorname{disc}(A)^{\frac{k-1}{2}} & A \otimes \mathbb{Z}_\ell = A_\ell \ \forall \ell|N \\ 0 & \text{otherwise} \end{cases}$$

Plan of talk

1. Theory of Gan-Gurevich lifts, and role of epsilon factors
2. A construction of $\mathcal{A}_{GG}(f_\chi)$
3. Sketch of proof

1. Theory of Gan-Gurevich lifts

Arthur parameters

Langlands philosophy, G/F reductive group:

$$\{\text{aut. reps of } G\} \longleftrightarrow \left\{ \text{“Galois representations” } L_{\mathbb{Q}} \rightarrow {}^L G \right\}$$

Arthur parameters

Langlands philosophy, G/F reductive group:

$$\{\text{aut. reps of } G\} \longleftrightarrow \left\{ \text{“Galois representations” } L_{\mathbb{Q}} \rightarrow {}^L G \right\}$$

Arthur's conjecture:

$$\mathcal{A}_{\text{disc}}(G) = \bigoplus_{\psi} \mathcal{A}_{\psi}(G)$$

where

$$\psi : L_{\mathbb{Q}} \times \text{SL}_2(\mathbb{C}) \rightarrow {}^L G,$$

$\psi|_{L_{\mathbb{Q}}}$ is tempered, and $\psi|_{\text{SL}_2(\mathbb{C})}$ is algebraic

Arthur parameters

Langlands philosophy, G/F reductive group:

$$\{\text{aut. reps of } G\} \longleftrightarrow \left\{ \text{“Galois representations” } L_{\mathbb{Q}} \rightarrow {}^L G \right\}$$

Arthur's conjecture:

$$\mathcal{A}_{\text{disc}}(G) = \bigoplus_{\psi} \mathcal{A}_{\psi}(G)$$

where

$$\psi : L_{\mathbb{Q}} \times \text{SL}_2(\mathbb{C}) \rightarrow {}^L G,$$

$\psi|_{L_{\mathbb{Q}}}$ is tempered, and $\psi|_{\text{SL}_2(\mathbb{C})}$ is algebraic

The more nontrivial the $\text{SL}_2(\mathbb{C})$, the more nontempered (and degenerate) the representation.

Arthur parameters

Langlands philosophy, G/F reductive group:

$$\{\text{aut. reps of } G\} \longleftrightarrow \left\{ \text{“Galois representations” } L_{\mathbb{Q}} \rightarrow {}^L G \right\}$$

Arthur's conjecture:

$$\mathcal{A}_{\text{disc}}(G) = \bigoplus_{\psi} \mathcal{A}_{\psi}(G)$$

where

$$\psi : L_{\mathbb{Q}} \times \text{SL}_2(\mathbb{C}) \rightarrow {}^L G,$$


$\psi|_{L_{\mathbb{Q}}}$ is tempered, and $\psi|_{\text{SL}_2(\mathbb{C})}$ is algebraic

The more nontrivial the $\text{SL}_2(\mathbb{C})$, the more nontempered (and degenerate) the representation. e.g. for $G = \text{GL}_2$, ${}^L G = \text{GL}_2(\mathbb{C})$: $\psi|_{\text{SL}_2(\mathbb{C})}$ nontrivial corresponds to the characters

Theory of Gan-Gurevich lifts: Arthur parameters

$$G = G_2, {}^L G = G_2(\mathbb{C})$$


$$\psi : L_{\mathbb{Q}} \times SL_2(\mathbb{C}) \rightarrow SL_{2,\text{short}}(\mathbb{C}) \times SL_{2,\text{long}}(\mathbb{C}) \rightarrow G_2(\mathbb{C})$$



Theory of Gan-Gurevich lifts: Arthur parameters

$$G = G_2, {}^L G = G_2(\mathbb{C})$$

$$\psi : L_{\mathbb{Q}} \times SL_2(\mathbb{C}) \rightarrow SL_{2,\text{short}}(\mathbb{C}) \times SL_{2,\text{long}}(\mathbb{C}) \rightarrow G_2(\mathbb{C})$$




For any $\tau =$ cuspidal AR of PGL_2 , consider ψ_τ where $L_{\mathbb{Q}} \rightarrow SL_{2,\text{short}}(\mathbb{C})$ corresponds to τ .

Theory of Gan-Gurevich lifts: Arthur parameters

$$G = G_2, {}^L G = G_2(\mathbb{C})$$

$$\psi : L_{\mathbb{Q}} \times SL_2(\mathbb{C}) \rightarrow SL_{2,\text{short}}(\mathbb{C}) \times SL_{2,\text{long}}(\mathbb{C}) \rightarrow G_2(\mathbb{C})$$



For any $\tau =$ cuspidal AR of PGL_2 , consider ψ_τ where

$L_{\mathbb{Q}} \rightarrow SL_{2,\text{short}}(\mathbb{C})$ corresponds to τ .

Specialize to $\tau = \tau_\chi$ CM, $\mathcal{A}_{GG}(f_\chi) := \mathcal{A}_{\psi_{\tau_\chi}}(G_2)$

Epsilons

Arthur has a precise conjectural description for $\mathcal{A}_\psi(G)$ in terms of **local packets** and **global multiplicities**.

Epsilons

Arthur has a precise conjectural description for $\mathcal{A}_\psi(G)$ in terms of **local packets** and **global multiplicities**.

Arthur's prediction for structure of global GG packet (partially known, Alonso-He-Ray-Roset 23 and BHL-HS24):

$$\mathcal{A}_{GG}(f_\chi) = \bigoplus_{(\epsilon_v)_v} m((\epsilon_v)_v) \bigotimes_v \pi_v^{\epsilon_v}$$

where $\{\pi_v^+, \pi_v^-\}$ is a local packet depending only on χ_v , with $\pi_v^- = 0$ almost everywhere, and

$$m((\epsilon_v)_v) = \begin{cases} 1 & \prod \epsilon_v = \epsilon(1/2, \chi^3) \\ 0 & \text{else} \end{cases}$$

Epsilons

Arthur has a precise conjectural description for $\mathcal{A}_\psi(G)$ in terms of **local packets** and **global multiplicities**.

Arthur's prediction for structure of global GG packet (partially known, Alonso-He-Ray-Roset 23 and BHL-HS24):

$$\mathcal{A}_{GG}(f_\chi) = \bigoplus_{(\epsilon_v)_v} m((\epsilon_v)_v) \bigotimes_v \pi_v^{\epsilon_v}$$

where $\{\pi_v^+, \pi_v^-\}$ is a local packet depending only on χ_v , with $\pi_v^- = 0$ almost everywhere, and

$$m((\epsilon_v)_v) = \begin{cases} 1 & \prod \epsilon_v = \epsilon(1/2, \chi^3) \\ 0 & \text{else} \end{cases}$$

It is always interesting to see how arithmetic conspires to enforce the multiplicity formula.

Epsilons

Arthur has a precise conjectural description for $\mathcal{A}_\psi(G)$ in terms of **local packets** and **global multiplicities**.

Arthur's prediction for structure of global GG packet (partially known, Alonso-He-Ray-Roset 23 and BHL-HS24):

$$\mathcal{A}_{GG}(f_\chi) = \bigoplus_{(\epsilon_v)_v} m((\epsilon_v)_v) \bigotimes_v \pi_v^{\epsilon_v}$$

where $\{\pi_v^+, \pi_v^-\}$ is a local packet depending only on χ_v , with $\pi_v^- = 0$ almost everywhere, and

$$m((\epsilon_v)_v) = \begin{cases} 1 & \prod \epsilon_v = \epsilon(1/2, \chi^3) \\ 0 & \text{else} \end{cases}$$

It is always interesting to see how arithmetic conspires to enforce the multiplicity formula. Here, it comes in via local-global obstruction for existence of Hermitian spaces.

Epsilons

Arthur's prediction for structure of GG packet:

$$\mathcal{A}_{GG}(f_\chi) = \bigoplus_{(\epsilon_v)_v} m((\epsilon_v)_v) \bigotimes'_v \pi_v^{\epsilon_v}$$

$$m((\epsilon_v)_v) = \begin{cases} 1 & \prod \epsilon_v = \epsilon(1/2, \chi^3) \\ 0 & \text{else} \end{cases}$$

Given $\pi = \bigotimes'_v \pi_v^{\epsilon_v}$:

- $\varphi \in \pi$ can be a QMF only if $\epsilon_\infty = -1$.

Epsilons

Arthur's prediction for structure of GG packet:

$$\mathcal{A}_{GG}(f_\chi) = \bigoplus_{(\epsilon_v)_v} m((\epsilon_v)_v) \bigotimes'_v \pi_v^{\epsilon_v}$$

$$m((\epsilon_v)_v) = \begin{cases} 1 & \prod \epsilon_v = \epsilon(1/2, \chi^3) \\ 0 & \text{else} \end{cases}$$

Given $\pi = \bigotimes'_v \pi_v^{\epsilon_v}$:

- $\varphi \in \pi$ can be a QMF only if $\epsilon_\infty = -1$.
- $\varphi \in \pi$ can have $c_A(\varphi) \neq 0$ only if $\epsilon_\ell = \epsilon_\ell(A \otimes \mathbb{Z}_\ell, \chi_\ell)$ for all $\ell | N$.

Epsilons

Arthur's prediction for structure of GG packet:

$$\mathcal{A}_{GG}(f_\chi) = \bigoplus_{(\epsilon_v)_v} m((\epsilon_v)_v) \bigotimes'_v \pi_v^{\epsilon_v}$$

$$m((\epsilon_v)_v) = \begin{cases} 1 & \prod \epsilon_v = \epsilon(1/2, \chi^3) \\ 0 & \text{else} \end{cases}$$

Given $\pi = \bigotimes'_v \pi_v^{\epsilon_v}$:

- $\varphi \in \pi$ can be a QMF only if $\epsilon_\infty = -1$.
- $\varphi \in \pi$ can have $c_A(\varphi) \neq 0$ only if $\epsilon_\ell = \epsilon_\ell(A \otimes \mathbb{Z}_\ell, \chi_\ell)$ for all $\ell | N$.

\implies to see $c_A(\varphi)$, need $\prod \epsilon_\ell(A \otimes \mathbb{Z}_\ell, \chi_\ell) = -\epsilon(1/2, \chi^3)$

Main result

- $f = f_\chi$ CM form of weight k , trivial character, and any level N , associated to K/\mathbb{Q} and $\chi : K^\times \backslash \mathbb{A}_K^\times \rightarrow \mathbb{C}^\times$.
- Assume: $L(1/2, \chi) \neq 0$.
- $\mathcal{A}_{GG}(f_\chi)$ space of Gan-Gurevich lifts $G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$.

Theorem (Bakic–Horawa–Li–Huerta–S., in progress)

For all $\ell|N$, fix a cubic ring A_ℓ/\mathbb{Z}_ℓ , such that

$$\prod_{\ell|N} \epsilon_\ell(A_\ell, \chi_\ell) = -\epsilon(1/2, \chi^3)$$

Then \exists a QMF $\varphi \in \mathcal{A}_{GG}(f_\chi)$ s.t. for A maximal outside N :

$$c_A(\varphi) = \begin{cases} L(1/2, f_\chi \otimes \rho_A) \operatorname{disc}(A)^{\frac{k-1}{2}} & A \otimes \mathbb{Z}_\ell = A_\ell \ \forall \ell|N \\ 0 & \text{otherwise} \end{cases}$$

2. construction of $\mathcal{A}_{GG}(f_\chi)$

Exceptional theta correspondences

- A theta correspondence is a construction

$$\Theta : \{\text{ARs of } G\} \rightsquigarrow \{\text{ARs of } H\}$$

for “dual reductive pairs” H, G .

- Means can embed as mutual centralizers $H \times G \hookrightarrow \tilde{G}$, where \tilde{G} is a group with theta functions (used as a kernel).

Exceptional theta correspondences

- A theta correspondence is a construction

$$\Theta : \{\text{ARs of } G\} \rightsquigarrow \{\text{ARs of } H\}$$

for “dual reductive pairs” H, G .

- Means can embed as mutual centralizers $H \times G \hookrightarrow \tilde{G}$, where \tilde{G} is a group with theta functions (used as a kernel).

$H \times G$	\tilde{G}
$SO_n \times Sp_{2m}$	Sp_{2mn}
$U(n) \times U(m)$	Sp_{2mn}

Exceptional theta correspondences

- A theta correspondence is a construction

$$\Theta : \{\text{ARs of } G\} \rightsquigarrow \{\text{ARs of } H\}$$

for “dual reductive pairs” H, G .

- Means can embed as mutual centralizers $H \times G \hookrightarrow \tilde{G}$, where \tilde{G} is a group with theta functions (used as a kernel).

$H \times G$	\tilde{G}	$H \times G$	\tilde{G}
$SO_n \times Sp_{2m}$	Sp_{2mn}	$G_2 \times PGL_3$	E_6
$U(n) \times U(m)$	Sp_{2mn}	$G_2 \times PGSp_6$	E_7
		$G_2 \times F_4$	E_8

Exceptional theta correspondences

- A theta correspondence is a construction

$$\Theta : \{\text{ARs of } G\} \rightsquigarrow \{\text{ARs of } H\}$$

for “dual reductive pairs” H, G .

- Means can embed as mutual centralizers $H \times G \hookrightarrow \tilde{G}$, where \tilde{G} is a group with theta functions (used as a kernel).

$H \times G$	\tilde{G}	$H \times G$	\tilde{G}
$SO_n \times Sp_{2m}$	Sp_{2mn}	$G_2 \times PU_3$	E'_6
$U(n) \times U(m)$	Sp_{2mn}	$G_2 \times PGSp_6$	E_7
		$G_2 \times F_4$	E_8

Exceptional theta correspondences and Gan-Gurevich lifts

Gan-Gurevich:

$$\begin{array}{c} \sigma \\ \text{PGL}_2 \rightsquigarrow \text{PGSp}_6 \xrightarrow{\Theta} G_2 \end{array}$$

Exceptional theta correspondences and Gan-Gurevich lifts

Gan-Gurevich:

$$\begin{array}{ccc} & \sigma & \\ & \text{PGL}_2 \rightsquigarrow \text{PGSp}_6 & \\ & \searrow \Theta & \\ & & G_2 \\ & \nearrow \Theta & \\ \text{PU}_3 & & \end{array}$$

Alternative approach when $\sigma = \text{CM form associated to } \chi$, cf.
[BHL-HS24]

Exceptional theta correspondences and Gan-Gurevich lifts

Gan-Gurevich:

$$\begin{array}{ccccc} & \sigma & & & \\ & \text{PGL}_2 & \rightsquigarrow & \text{PGSp}_6 & \xrightarrow{\Theta} G_2 \\ & U(1) & \xrightarrow{\Theta_\chi} & \text{PU}_3 & \nearrow \Theta \\ & \mathbb{1} & & & \end{array}$$

Alternative approach when $\sigma = \text{CM form associated to } \chi$, cf.
[BHL-HS24]

Exceptional theta correspondences and Gan-Gurevich lifts

Gan-Gurevich:


$$\begin{array}{ccccc} & \sigma & & & \\ & & & & \\ \mathrm{PGL}_2 & \rightsquigarrow & \mathrm{PGSp}_6 & \xrightarrow{\Theta} & G_2 \\ & & & \nearrow \Theta & \\ U(1) & \xrightarrow{\Theta_\chi} & \mathrm{PU}_3 & & \\ \mathbb{1} & & & & \end{array}$$

Alternative approach when $\sigma = \text{CM form associated to } \chi$, cf. [BHL-HS24]

Easier to understand packet structure (comes from Howe–Piatetskii-Shapiro CAP forms on PU_3)

3. sketch of proof

What are Fourier coefficients?



$$\{\text{characters of } N(\mathbb{Q}) \backslash N(\mathbb{A})\} \longleftrightarrow \{\lambda = (a, b, c, d) \in \mathbb{Q}^4\}$$
$$\psi_\lambda \longleftrightarrow \lambda$$

$$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$$

$$c_\lambda(\varphi) := \int_{[N]} \psi_\lambda^{-1}(n) \varphi(n) dn$$

What are Fourier coefficients?

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ modular form of weight $k \geq 2$

$$c_\lambda(\varphi) := \int_{[N]} \psi_\lambda^{-1}(n) \varphi(n) dn$$

What are Fourier coefficients?

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ modular form of weight $k \geq 2$

$$c_\lambda(\varphi) := \int_{[N]} \psi_\lambda^{-1}(n) \varphi(n) dn$$

- φ invariant by $P(\widehat{\mathbb{Z}}) \implies c_\lambda(\varphi)$ is supported on $\lambda \in \mathbb{Z}^4$

What are Fourier coefficients?

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ modular form of weight $k \geq 2$

$$c_\lambda(\varphi) := \int_{[N]} \psi_\lambda^{-1}(n) \varphi(n) dn$$

- φ invariant by $P(\widehat{\mathbb{Z}}) \implies c_\lambda(\varphi)$ is supported on $\lambda \in \mathbb{Z}^4$ and depends only on the **cubic ring** A_λ
 - $A_\lambda = \mathbb{Z}^3$ with multiplication determined by $\lambda = (a, b, c, d) \in \mathbb{Z}^4$

What are Fourier coefficients?

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ modular form of weight $k \geq 2$

$$c_\lambda(\varphi) := \int_{[N]} \psi_\lambda^{-1}(n) \varphi(n) dn$$

- φ invariant by $P(\widehat{\mathbb{Z}}) \implies c_\lambda(\varphi)$ is supported on $\lambda \in \mathbb{Z}^4$ and depends only on the **cubic ring** A_λ
 - $A_\lambda = \mathbb{Z}^3$ with multiplication determined by $\lambda = (a, b, c, d) \in \mathbb{Z}^4$
- More generally have a cubic algebra E_λ/\mathbb{Q}

What are Fourier coefficients?

$\varphi : G_2(\mathbb{Q}) \backslash G_2(\mathbb{A}) \rightarrow \mathbb{C}$ modular form of weight $k \geq 2$

$$c_\lambda(\varphi) := \int_{[N]} \psi_\lambda^{-1}(n) \varphi(n) dn$$

- φ invariant by $P(\widehat{\mathbb{Z}}) \implies c_\lambda(\varphi)$ is supported on $\lambda \in \mathbb{Z}^4$ and depends only on the **cubic ring** A_λ
 - $A_\lambda = \mathbb{Z}^3$ with multiplication determined by $\lambda = (a, b, c, d) \in \mathbb{Z}^4$
- More generally have a cubic algebra E_λ/\mathbb{Q}
- Turns out $c_\lambda(\varphi) = 0$ unless E_λ is totally real

Proof sketch

Periods for $\Theta(\pi)$ often reduce to periods for π .

Proof sketch

Periods for $\Theta(\pi)$ often reduce to periods for π .

For $\varphi \in \mathcal{A}_{GG}(f_\chi)$ (any level),

$$c_\lambda(\varphi) \sim \int_{[T_E]} \rho(t) dt$$

where:

- $E = E_\lambda$ is totally real cubic étale algebra corresponding to λ .
- $T_E \hookrightarrow \mathrm{PU}_3$ is a torus embedding coming from $E \hookrightarrow \mathrm{Herm}_{3 \times 3}(\mathbb{Q})$.
- ρ = thing you're lifting on PU_3 (Howe-PS CAP forms).

Proof sketch

Periods for $\Theta(\pi)$ often reduce to periods for π .

For $\varphi \in \mathcal{A}_{GG}(f_\chi)$ (any level),

$$c_\lambda(\varphi) \sim \int_{[T_E]} \rho(t) dt$$

ρ is itself a Θ lift from $U(1)$

Proof sketch

Periods for $\Theta(\pi)$ often reduce to periods for π .

For $\varphi \in \mathcal{A}_{GG}(f_\chi)$ (any level),

$$c_\lambda(\varphi) \sim \int_{[T_E]} \rho(t) dt$$

ρ is itself a Θ lift from $U(1)$

$$\left| \int_{[T_E]} \rho(t) dt \right|^2 \sim L(1/2, f_\chi \otimes \rho_E) L(1/2, \chi) \Delta_E^{1/2}$$

cf. Yang 97, Borade-Franzel-Girsch-Yao-Yu-Zelingher 24

Proof sketch

$$|c_\lambda(\varphi)|^2 = L(1/2, f_\chi \otimes \rho_E) L(1/2, \chi) \Delta_E^{1/2} \prod_v |I_v(\varphi_v, \lambda)|^2$$

Proof sketch

$$|c_\lambda(\varphi)|^2 = L(1/2, f_\chi \otimes \rho_E) L(1/2, \chi) \Delta_E^{1/2} \prod_v |I_v(\varphi_v, \lambda)|^2$$

- Additional $\Delta_E^{k/2-1}$ comes from evaluating I_∞ (using Pollack's explicit Whittaker model of minimal representation of E_6)

Proof sketch

$$|c_\lambda(\varphi)|^2 = L(1/2, f_\chi \otimes \rho_E) L(1/2, \chi) \Delta_E^{1/2} \prod_v |I_v(\varphi_v, \lambda)|^2$$

- Additional $\Delta_E^{k/2-1}$ comes from evaluating I_∞ (using Pollack's explicit Whittaker model of minimal representation of E_6)
- $I_\ell(\varphi_\ell, \lambda)$ is hard to compute in general
 - For $\ell \nmid N$, we show $I_\ell(\varphi_\ell^{\text{sph}}, \lambda) = 1$ if $A_\lambda \otimes \mathbb{Z}_\ell$ is a maximal order in a cubic étale algebra

Proof sketch

$$|c_\lambda(\varphi)|^2 = L(1/2, f_\chi \otimes \rho_E) L(1/2, \chi) \Delta_E^{1/2} \prod_v |I_v(\varphi_v, \lambda)|^2$$

- Additional $\Delta_E^{k/2-1}$ comes from evaluating I_∞ (using Pollack's explicit Whittaker model of minimal representation of E_6)
- $I_\ell(\varphi_\ell, \lambda)$ is hard to compute in general
 - For $\ell \nmid N$, we show $I_\ell(\varphi_\ell^{\text{sph}}, \lambda) = 1$ if $A_\lambda \otimes \mathbb{Z}_\ell$ is a maximal order in a cubic étale algebra
 - For $\ell | N$, we show you can “rig” φ_ℓ so $I_\ell(\varphi_\ell, \lambda)$ is the indicator function of $A_\lambda \otimes \mathbb{Z}_\ell = \text{any fixed } A_\ell$

Thanks!