


# BZSV Duality and Relative Langlands Program

Wee Teck Gan

January 22, 2024

(Joint with Bryan P.J. Wang)



# Local Langlands Program

- $F$  local field,
- $G$  reductive group over  $F$ ; set  $G = G(F)$
- $\text{Irr}(G) = \{\text{isom. classes of irred. smooth reps of } G\}.$

# Local Langlands Program

- $F$  local field,
- $G$  reductive group over  $F$ ; set  $G = G(F)$
- $\text{Irr}(G) = \{\text{isom. classes of irred. smooth reps of } G\}$ .

**Main Problem:** Classify  $\text{Irr}(G)$ .

# Local Langlands Program

- $F$  local field,
- $G$  reductive group over  $F$ ; set  $G = G(F)$
- $\text{Irr}(G) = \{\text{isom. classes of irred. smooth reps of } G\}.$

**Main Problem:** Classify  $\text{Irr}(G)$ .

**Main Conjecture:** There is a surjective map

$$\mathcal{L}_G : \text{Irr}(G) \longrightarrow \Phi(G) = \{\text{L-parameters of } G\}$$

with finite fibers which can also be explicitly parametrized.

# Local Langlands Program

- $F$  local field,
- $G$  reductive group over  $F$ ; set  $G = G(F)$
- $\text{Irr}(G) = \{\text{isom. classes of irred. smooth reps of } G\}$ .

**Main Problem:** Classify  $\text{Irr}(G)$ .

**Main Conjecture:** There is a surjective map

$$\mathcal{L}_G : \text{Irr}(G) \longrightarrow \Phi(G) = \{\text{L-parameters of } G\}$$

with finite fibers which can also be explicitly parametrized.

An L-parameter is an equivalence class of maps

$$\phi : \Gamma_F = \text{Gal}(\overline{F}/F) = \pi_1^{et}(\text{Spec}(F)) \longrightarrow G^\vee$$

where  $G^\vee$  is the Langlands dual group of  $G$  and equivalence is up to  $G^\vee$ -conjugacy. So (roughly)

$$\Phi(G) = \text{Hom}(\pi_1^{et}(\text{Spec}(F)), G^\vee) / G^\vee$$

# Global Langlands Program

- $F$  global field and  $G$  reductive group over  $F$
- $G(F) \subset G(\mathbb{A}) = \prod'_v G(F_v)$ ; set  $[G] = G(F) \backslash G(\mathbb{A})$ ;
- $\mathcal{A}_2(G) = \{\text{square-integrable automorphic forms of } G\} = \{f : [G] \rightarrow \mathbb{C}\}.$

# Global Langlands Program

- $F$  global field and  $G$  reductive group over  $F$
- $G(F) \subset G(\mathbb{A}) = \prod'_v G(F_v)$ ; set  $[G] = G(F) \backslash G(\mathbb{A})$ ;
- $\mathcal{A}_2(G) = \{\text{square-integrable automorphic forms of } G\} = \{f : [G] \rightarrow \mathbb{C}\}.$

**Main Problem:** Classify the irreducible constituents of  $\mathcal{A}_2(G)$ .

**Main Conjecture:** The irreducible constituents of  $\mathcal{A}_2(G)$  can be parametrized by A-parameters:

$$\Psi : \text{Gal}(\overline{F}/F) \times \text{SL}_2(\mathbb{C}) \longrightarrow G^\vee$$

# Relative Langlands?

- $H \subset G$  a subgroup over  $F$
- $\chi$  a 1-dimensional (automorphic) unitary character of  $H$ .



# Relative Langlands?

- $H \subset G$  a subgroup over  $F$
- $\chi$  a 1-dimensional (automorphic) unitary character of  $H$ .

**Local:** Classify the set

$$\mathrm{Irr}_{H,\chi}(G) := \{\pi : \mathrm{Hom}_H(\pi, \chi) \neq 0\} \subset \mathrm{Irr}(G)$$

of  $(H, \chi)$ -distinguished representations.

# Relative Langlands?

- $H \subset G$  a subgroup over  $F$
- $\chi$  a 1-dimensional (automorphic) unitary character of  $H$ .

**Local:** Classify the set

$$\mathrm{Irr}_{H,\chi}(G) := \{\pi : \mathrm{Hom}_H(\pi, \chi) \neq 0\} \subset \mathrm{Irr}(G)$$

of  $(H, \chi)$ -distinguished representations. By Frobenius reciprocity,

$$\mathrm{Irr}_{H,\chi}(G) = \{\pi : \mathrm{Hom}_G(\pi, \mathrm{Ind}_H^G \chi) \neq 0\},$$

so it is the set of irreducible submodules of  $C^\infty(H, \chi \backslash G)$ .

# Relative Langlands?

- $H \subset G$  a subgroup over  $F$
- $\chi$  a 1-dimensional (automorphic) unitary character of  $H$ .

**Local:** Classify the set

$$\mathrm{Irr}_{H,\chi}(G) := \{\pi : \mathrm{Hom}_H(\pi, \chi) \neq 0\} \subset \mathrm{Irr}(G)$$

of  $(H, \chi)$ -distinguished representations. By Frobenius reciprocity,

$$\mathrm{Irr}_{H,\chi}(G) = \{\pi : \mathrm{Hom}_G(\pi, \mathrm{Ind}_H^G \chi) \neq 0\},$$

so it is the set of irreducible submodules of  $C^\infty(H, \chi \backslash G)$ .

**$L^2$ -version:** describe the spectral decomposition of  $L^2(H, \chi \backslash G)$ .

# Relative Langlands?

- $H \subset G$  a subgroup over  $F$
- $\chi$  a 1-dimensional (automorphic) unitary character of  $H$ .

**Local:** Classify the set

$$\mathrm{Irr}_{H,\chi}(G) := \{\pi : \mathrm{Hom}_H(\pi, \chi) \neq 0\} \subset \mathrm{Irr}(G)$$

of  $(H, \chi)$ -distinguished representations. By Frobenius reciprocity,

$$\mathrm{Irr}_{H,\chi}(G) = \{\pi : \mathrm{Hom}_G(\pi, \mathrm{Ind}_H^G \chi) \neq 0\},$$

so it is the set of irreducible submodules of  $C^\infty(H, \chi \backslash G)$ .

**$L^2$ -version:** describe the spectral decomposition of  $L^2(H, \chi \backslash G)$ .

**Expect:**  $\mathrm{Irr}_{H,\chi}(G)$  corresponds to L-parameters which factor through some  $J^\vee \rightarrow G^\vee$ . So  $(H, \chi)$ -dist. reps are functorial lifts from another group  $J$ ).

# Relative Langlands?

**Global:** Have global period integral

$$\mathcal{P}_{H,\chi} : \mathcal{A}_2(G) \longrightarrow \mathbb{C}$$

defined by

$$\mathcal{P}_{H,\chi}(f) = \int_{[H]} \overline{\chi(h)} \cdot f(h) dh.$$

# Relative Langlands?

**Global:** Have global period integral

$$\mathcal{P}_{H,\chi} : \mathcal{A}_2(G) \longrightarrow \mathbb{C}$$

defined by

$$\mathcal{P}_{H,\chi}(f) = \int_{[H]} \overline{\chi(h)} \cdot f(h) dh.$$

Classify those  $\pi \subset \mathcal{A}_2(G)$  such that

$$\mathcal{P}_{H,\chi,\pi} := \mathcal{P}_{H,\chi}|_{\pi} \neq 0 \in \mathrm{Hom}_{H(\mathbb{A})}(\pi, \chi).$$

# Relative Langlands?

**Global:** Have global period integral

$$\mathcal{P}_{H,\chi} : \mathcal{A}_2(G) \longrightarrow \mathbb{C}$$

defined by

$$\mathcal{P}_{H,\chi}(f) = \int_{[H]} \overline{\chi(h)} \cdot f(h) dh.$$

Classify those  $\pi \subset \mathcal{A}_2(G)$  such that

$$\mathcal{P}_{H,\chi,\pi} := \mathcal{P}_{H,\chi}|_{\pi} \neq 0 \in \mathrm{Hom}_{H(\mathbb{A})}(\pi, \chi).$$

Expect:

- $\mathcal{P}_{H,\chi,\pi}$  can be factored as product of local  $H(F_v)$ -invariant functionals
- $\mathcal{P}_{H,\chi,\pi}$  to be related to an L-function on  $J$ .

# Classical Examples

Periods	$(G, H, \chi)$
Whittaker	$G \supset U$ (maximal unipotent) $\chi = \psi$ generic character
Symplectic	$GL_{2n} \supset Sp_{2n}$
Shalika	$GL_{2n} \supset P_{n,n} \supset GL_n^\Delta U$ $\chi = 1_{GL_n} \otimes \psi(\text{Tr}(-))$
Basic Gross-Prasad	$SO_{2n} \times SO_{2n+1} \supset SO_{2n}^\Delta$
General GP $n < m$ opp. parity	$SO_n \times SO_m \supset SO_n^\Delta U$ $\chi$ generic character



# Questions and Issues

- Which subgroups  $H \subset G$  to consider?

# Questions and Issues

- Which subgroups  $H \subset G$  to consider?
- For which  $(H, \chi)$  is  $L^2(H, \chi \backslash G)$  multiplicity-free?

# Questions and Issues

- Which subgroups  $H \subset G$  to consider?
- For which  $(H, \chi)$  is  $L^2(H, \chi \backslash G)$  multiplicity-free?
- For a given  $(H, \chi)$ , from which group  $J$  are  $(H, \chi)$ -dist. reps lifted?

# Questions and Issues

- Which subgroups  $H \subset G$  to consider?
- For which  $(H, \chi)$  is  $L^2(H, \chi \backslash G)$  multiplicity-free?
- For a given  $(H, \chi)$ , from which group  $J$  are  $(H, \chi)$ -dist. reps lifted?
- For a given  $(H, \chi)$ , which L-function on  $J$  is relevant for the global period?

# Questions and Issues

- Which subgroups  $H \subset G$  to consider?
- For which  $(H, \chi)$  is  $L^2(H, \chi \backslash G)$  multiplicity-free?
- For a given  $(H, \chi)$ , from which group  $J$  are  $(H, \chi)$ -dist. reps lifted?
- For a given  $(H, \chi)$ , which L-function on  $J$  is relevant for the global period?

**Upshot** The subject of RLP is based on a number of examples but there is no systematic framework,

# Questions and Issues

- Which subgroups  $H \subset G$  to consider?
- For which  $(H, \chi)$  is  $L^2(H, \chi \backslash G)$  multiplicity-free?
- For a given  $(H, \chi)$ , from which group  $J$  are  $(H, \chi)$ -dist. reps lifted?
- For a given  $(H, \chi)$ , which L-function on  $J$  is relevant for the global period?

**Upshot** The subject of RLP is based on a number of examples but there is no systematic framework, that is, until the publication of the Asterisque volume [SV]:



# Relative Langlands according to [SV]

Objects	spherical $G$ -variety $X = H \backslash G$
Local Qn	spectral decomposition of $L^2(X)$ or $C^\infty(X)$
Global Qn	global H-periods and related L-values
Dual Data	(i) $\iota_X : X^\vee \times \mathrm{SL}_2 \rightarrow G^\vee$ (ii) (graded symplectic) representation $V_X$ of $X^\vee$
Conjecture	(local) H-dist. $\pi$ are functorial lifts from $X^\vee$ via $\iota_X$ (global) H-period of $\pi$ given by $L(1/2, \sigma_\pi, V_X)$

# Issues with [SV]

Despite providing a uniform framework for the RLP and constructing the Langlands dual data for a spherical variety  $X$ , there are still some lingering issues with [SV]:



# Issues with [SV]

Despite providing a uniform framework for the RLP and constructing the Langlands dual data for a spherical variety  $X$ , there are still some lingering issues with [SV]:

- the definition of  $V_X$  from  $X$  follows a combinatorial algorithm which is neither so transparent nor conceptual.

# Issues with [SV]

Despite providing a uniform framework for the RLP and constructing the Langlands dual data for a spherical variety  $X$ , there are still some lingering issues with [SV]:

- the definition of  $V_X$  from  $X$  follows a combinatorial algorithm which is neither so transparent nor conceptual.
- it is not clear what L-functions (or  $V_X$ ) will arise from spherical varieties.

# Issues with [SV]

Despite providing a uniform framework for the RLP and constructing the Langlands dual data for a spherical variety  $X$ , there are still some lingering issues with [SV]:

- the definition of  $V_X$  from  $X$  follows a combinatorial algorithm which is neither so transparent nor conceptual.
- it is not clear what L-functions (or  $V_X$ ) will arise from spherical varieties.
- there are certain natural  $G$ -modules which are multiplicity-free and whose spectral decomposition can be described in the style of [SV], but which nonetheless do not fall into the framework of [SV].

# Issues with [SV]

Despite providing a uniform framework for the RLP and constructing the Langlands dual data for a spherical variety  $X$ , there are still some lingering issues with [SV]:

- the definition of  $V_X$  from  $X$  follows a combinatorial algorithm which is neither so transparent nor conceptual.
- it is not clear what L-functions (or  $V_X$ ) will arise from spherical varieties.
- there are certain natural  $G$ -modules which are multiplicity-free and whose spectral decomposition can be described in the style of [SV], but which nonetheless do not fall into the framework of [SV].

An example of this last point is the theta correspondence, i.e. the spectral decomposition of the Weil representation under the action of a dual pair.


The above issues are (to some extent) resolved in the 400-page preprint [BZSV] of Ben-Zvi, Sakellaridis and Venkatesh:

#### RELATIVE LANGLANDS DUALITY

DAVID BEN-ZVI, YIANNIS SAKELLARIDIS AND AKSHAY VENKATESH

**ABSTRACT.** We propose a duality in the relative Langlands program. This duality pairs a Hamiltonian space for a group  $G$  with a Hamiltonian space under its dual group  $\check{G}$ , and recovers at a numerical level the relationship between a period on  $G$  and an  $L$ -function attached to  $\check{G}$ ; it is an arithmetic analog of the electric-magnetic duality of boundary conditions in four-dimensional supersymmetric Yang-Mills theory.

This is a draft. We anticipate making another round of changes before submitting it for publication. All comments are very welcome! In particular, if we have failed to attribute or properly reference a work it is most likely due to either ignorance or forgetfulness - please tell us.



# Relative Langlands according to [BZSV]

Objects	Hyperspherical Hamiltonian $G$ -variety $M$
Local Qn	spectral decomp. of quantization $\Pi_M$ of $M$
Global Qn	spectral decomp. of theta function of $\Pi_M$
Dual Data	Hyperspherical Hamiltonian $G^\vee$ -variety $M^\vee$
Conjecture	(local) Galois action has fixed point on $M^\vee$ (global) L-function arises fro Galois rep. on tangent spaces of fixed points

# Comparing [BZSV] with [SV]

	[SV]	[BZSV]
Objects	spherical $X$	hyperspherical $M$
Spectral Qn	$L^2(X)$	Quantization of $M$
Dual Data	$(X^\vee, \iota_X, V_X)$	hyperspherical $M^\vee$
Conj.	Factor through $\iota_X$	Galois-fixed points on $M^\vee$

# Hyperspherical Hamiltonian $G$ -varieties $M$

(a) **symplectic  $G$ -variety**:  $M$  is a symplectic variety with  $G$  acting as symplectomorphisms. Then  $\mathcal{O}(M)$  is a  $G$ -equivariant Poisson algebra.



# Hyperspherical Hamiltonian $G$ -varieties $M$

- (a) **symplectic  $G$ -variety**:  $M$  is a symplectic variety with  $G$  acting as symplectomorphisms. Then  $\mathcal{O}(M)$  is a  $G$ -equivariant Poisson algebra.
- (b) **Hamiltonian  $G$ -variety**: a symplectic  $G$ -variety  $M$  equipped with a  $G$ -equivariant moment map  $\mu_M : M \rightarrow \mathfrak{g}^*$ .

# Hyperspherical Hamiltonian $G$ -varieties $M$

- (a) **symplectic  $G$ -variety**:  $M$  is a symplectic variety with  $G$  acting as symplectomorphisms. Then  $\mathcal{O}(M)$  is a  $G$ -equivariant Poisson algebra.
- (b) **Hamiltonian  $G$ -variety**: a symplectic  $G$ -variety  $M$  equipped with a  $G$ -equivariant moment map  $\mu_M : M \rightarrow \mathfrak{g}^*$ .
- (c) **Hyperspherical**: a Hamiltonian  $G$ -variety which is in addition:
  - affine, smooth and graded (with a commuting  $\mathbb{G}_m$ -action)

# Hyperspherical Hamiltonian $G$ -varieties $M$

- (a) **symplectic  $G$ -variety**:  $M$  is a symplectic variety with  $G$  acting as symplectomorphisms. Then  $\mathcal{O}(M)$  is a  $G$ -equivariant Poisson algebra.
- (b) **Hamiltonian  $G$ -variety**: a symplectic  $G$ -variety  $M$  equipped with a  $G$ -equivariant moment map  $\mu_M : M \rightarrow \mathfrak{g}^*$ .
- (c) **Hyperspherical**: a Hamiltonian  $G$ -variety which is in addition:
- affine, smooth and graded (with a commuting  $\mathbb{G}_m$ -action)
  - coisotropic: generic  $G$ -orbits are coisotropic i.e. for each orbit  $\mathcal{O}$  and  $x \in \mathcal{O}$ ,

# Hyperspherical Hamiltonian $G$ -varieties $M$

- (a) **symplectic  $G$ -variety**:  $M$  is a symplectic variety with  $G$  acting as symplectomorphisms. Then  $\mathcal{O}(M)$  is a  $G$ -equivariant Poisson algebra.
- (b) **Hamiltonian  $G$ -variety**: a symplectic  $G$ -variety  $M$  equipped with a  $G$ -equivariant moment map  $\mu_M : M \rightarrow \mathfrak{g}^*$ .
- (c) **Hyperspherical**: a Hamiltonian  $G$ -variety which is in addition:
- affine, smooth and graded (with a commuting  $\mathbb{G}_m$ -action)
  - coisotropic: generic  $G$ -orbits are coisotropic i.e. for each orbit  $\mathcal{O}$  and  $x \in \mathcal{O}$ ,

$$T_x \mathcal{O} \subset T_x M \text{ satisfies } T_x \mathcal{O}^\perp \subset T_x \mathcal{O}.$$

Equivalently, the subring  $\mathcal{O}(M)^G$  is Poisson-commutative.

- connected generic stabilizers (plus a couple of other technical conditions)

# Hyperspherical Hamiltonian $G$ -varieties $M$

- (a) **symplectic  $G$ -variety**:  $M$  is a symplectic variety with  $G$  acting as symplectomorphisms. Then  $\mathcal{O}(M)$  is a  $G$ -equivariant Poisson algebra.
- (b) **Hamiltonian  $G$ -variety**: a symplectic  $G$ -variety  $M$  equipped with a  $G$ -equivariant moment map  $\mu_M : M \rightarrow \mathfrak{g}^*$ .
- (c) **Hyperspherical**: a Hamiltonian  $G$ -variety which is in addition:
- affine, smooth and graded (with a commuting  $\mathbb{G}_m$ -action)
  - coisotropic: generic  $G$ -orbits are coisotropic i.e. for each orbit  $\mathcal{O}$  and  $x \in \mathcal{O}$ ,

$$T_x \mathcal{O} \subset T_x M \text{ satisfies } T_x \mathcal{O}^\perp \subset T_x \mathcal{O}.$$

Equivalently, the subring  $\mathcal{O}(M)^G$  is Poisson-commutative.

- connected generic stabilizers (plus a couple of other technical conditions)

If  $X$  is a spherical  $G$ -variety (affine smooth), then  $M = T^*X$  is hyperspherical.

# Structure Theory of Hyperspherical Varieties

Suppose one is given:

- $\iota : H \times \mathrm{SL}_2 \longrightarrow G$ , with  $H \subset Z_G(\iota(\mathrm{SL}_2))$  a spherical subgroup;
- $S$  a (graded) symplectic (finite dim) representation of  $H$ .

# Structure Theory of Hyperspherical Varieties

Suppose one is given:

- $\iota : H \times \mathrm{SL}_2 \longrightarrow G$ , with  $H \subset Z_G(\iota(\mathrm{SL}_2))$  a spherical subgroup;
- $S$  a (graded) symplectic (finite dim) representation of  $H$ .

Let  $\{h, e, f\}$  be  $\mathfrak{sl}_2$ -triple associated to  $\iota|_{\mathrm{SL}_2}$ . Then define a  $G$ -variety by:

$$M = (S \times_{\mathfrak{h}^*} (f + \mathfrak{g}^e)) \times^H G.$$

# Structure Theory of Hyperspherical Varieties

Suppose one is given:

- $\iota : H \times \mathrm{SL}_2 \longrightarrow G$ , with  $H \subset Z_G(\iota(\mathrm{SL}_2))$  a spherical subgroup;
- $S$  a (graded) symplectic (finite dim) representation of  $H$ .

Let  $\{h, e, f\}$  be  $\mathfrak{sl}_2$ -triple associated to  $\iota|_{\mathrm{SL}_2}$ . Then define a  $G$ -variety by:

$$M = (S \times_{\mathfrak{h}^*} (f + \mathfrak{g}^e)) \times^H G.$$

## Theorem (BZSV)

*Every hyperspherical  $G$ -variety is built in the above way (Whittaker induction).*



# Structure Theory of Hyperspherical Varieties

Suppose one is given:

- $\iota : H \times \mathrm{SL}_2 \longrightarrow G$ , with  $H \subset Z_G(\iota(\mathrm{SL}_2))$  a spherical subgroup;
- $S$  a (graded) symplectic (finite dim) representation of  $H$ .

Let  $\{h, e, f\}$  be  $\mathfrak{sl}_2$ -triple associated to  $\iota|_{\mathrm{SL}_2}$ . Then define a  $G$ -variety by:

$$M = (S \times_{\mathfrak{h}^*} (f + \mathfrak{g}^e)) \times^H G.$$

## Theorem (BZSV)

*Every hyperspherical  $G$ -variety is built in the above way (Whittaker induction).*

Observe that the above data is of the same type as the dual data from [SV]:

$$\iota_X : X^\vee \times \mathrm{SL}_2 \longrightarrow G^\vee$$

and  $V_X$  a (graded symplectic) rep. of  $X^\vee$ .

### 3 Basic Examples

M	Data for construction
$T^*(H \backslash G)$	$\iota : H \rightarrow G, S = 0$
$T_e^*(U \backslash G)$	$\iota : \mathrm{SL}_2 \rightarrow G \text{ (regular } \mathrm{SL}_2), S = 0$
$(W, G \subset \mathrm{Sp}(W))$	$\iota : G \rightarrow G, S = W$

### 3 Basic Examples

M	Data for construction
$T^*(H \backslash G)$	$\iota : H \rightarrow G, S = 0$
$T_e^*(U \backslash G)$	$\iota : \mathrm{SL}_2 \rightarrow G \text{ (regular } \mathrm{SL}_2), S = 0$
$(W, G \subset \mathrm{Sp}(W))$	$\iota : G \rightarrow G, S = W$

A mixed example:  $S = 0$  and

$$\iota : \mathrm{GL}_n \times \mathrm{SL}_2 \longrightarrow \mathrm{GL}_{2n} \quad (\text{tensor product}).$$

Then

$$M = \left\{ \begin{pmatrix} 0 & B \\ I & 0 \end{pmatrix} : B \in M_n \right\} \times^{\mathrm{GL}_n^\Delta} \mathrm{GL}_{2n} = T_e(\mathrm{GL}_n^\Delta U \backslash \mathrm{GL}_{2n})$$

Call this the Shalika variety since it gives rise to the Shalika period.

# Quantization

Quantization refers to the following philosophy:

- to a symplectic variety  $M$ , one can attach an associated Hilbert space  $\Pi_M$ .
- If a symplectic  $G$ -variety is Hamiltonian, then  $\Pi_M$  is a unitary rep. of  $G$ .

# Quantization

Quantization refers to the following philosophy:

- to a symplectic variety  $M$ , one can attach an associated Hilbert space  $\Pi_M$ .
- If a symplectic  $G$ -variety is Hamiltonian, then  $\Pi_M$  is a unitary rep. of  $G$ .

Ideally one would like a functor

$$\text{Symplectic } G\text{-varieties} \longrightarrow \text{Unitary } G\text{-modules}$$

# Quantization

Quantization refers to the following philosophy:

- to a symplectic variety  $M$ , one can attach an associated Hilbert space  $\Pi_M$ .
- If a symplectic  $G$ -variety is Hamiltonian, then  $\Pi_M$  is a unitary rep. of  $G$ .

Ideally one would like a functor

$$\text{Symplectic } G\text{-varieties} \longrightarrow \text{Unitary } G\text{-modules}$$

One does not have this, but many standard constructions in symplectic geometry can be quantized, i.e. have natural representation theoretic counterparts, as realized by Kirillov, Guillemin-Sternberg, Kazhdan etc.

# Classical vs. Quantization

Classical	Quantum
$M$	$(\rho_M, V_M)$
$C(M, \mathbb{R}) \subset C(M, \mathbb{C})$	$\text{Herm}(V_M) \subset \text{End}(V_M)$
$\mu : M \rightarrow \mathfrak{g}^*$	?
$\mu^* : C(\mathfrak{g}^*) \rightarrow C(M, \mathbb{C})$ (pullback)	$\rho_M : C^*(G) \rightarrow \text{End}(V_M)$ ( $C^*$ -alg. module)
Coisotropic	Multiplicity-free
$(M \times_{\mathfrak{g}^*}^G \{0\})$ (Symplectic reduction)	$(V_M)_G$ ( $G$ -coinvariants)

# Classical vs. Quantization

Classical	Quantum
$M$	$(\rho_M, V_M)$
$C(M, \mathbb{R}) \subset C(M, \mathbb{C})$	$\text{Herm}(V_M) \subset \text{End}(V_M)$
$\mu : M \rightarrow \mathfrak{g}^*$	?
$\mu^* : C(\mathfrak{g}^*) \rightarrow C(M, \mathbb{C})$ (pullback)	$\rho_M : C^*(G) \rightarrow \text{End}(V_M)$ ( $C^*$ -alg. module)
Coisotropic	Multiplicity-free
$(M \times_{\mathfrak{g}^*}^G \{0\})$ (Symplectic reduction)	$(V_M)_G$ ( $G$ -coinvariants)

The last line is often expressed as:

*Quantization commutes with Reduction.*



# Examples of Quantization

M	$\Pi_M$
$T^*(X),$ $X$ affine smooth spherical	$L^2(X)$
$T_e^*(U \backslash G) = (e + \mathfrak{g}^f) \times G$ (e regular nilpotent)	$L^2(U, \psi \backslash G)$ (Whittaker/Gelfand-Graev module)
$T_e^*(\mathrm{GL}_n^\Delta U \backslash \mathrm{GL}_{2n})$ (Shalika variety)	$L^2(\mathrm{GL}_n U, \psi \backslash \mathrm{GL}_{2n})$ (Shalika module)
symplectic vector space $W = X + X^*$	Weil representation $L^2(X)$
$(V \otimes W, \mathrm{O}(V) \times \mathrm{Sp}(W))$	Theta correspondence for $\mathrm{O}(V) \times \mathrm{Sp}(W)$

# Summary

	[SV]	[BZSV]
Objects	spherical $X$	hyperspherical $M$
Spectral Qn	$L^2(X)$	Quantization of $M$
Dual Data	$(X^\vee, \iota_X, V_X)$	hyperspherical $M^\vee$
Conj.	Factor though $\iota_X$	Galois-fixed points on $M^\vee$

# Summary

	[SV]	[BZSV]
Objects	spherical $X$	hyperspherical $M$
Spectral Qn	$L^2(X)$	Quantization of $M$
Dual Data	$(X^\vee, \iota_X, V_X)$	hyperspherical $M^\vee$
Conj.	Factor through $\iota_X$	Galois-fixed points on $M^\vee$

What is gained from [SV] to [BZSV]:

- Scope of RLP expanded (e.g. to include theta correspondence)
- there is now a clear symmetry between the basic object  $M$  and the dual data  $M^\vee$

## Two invariants associated to $M$

The symmetry between the automorphic and Galois side implies that to a hyperspherical  $G$ -variety  $M$ , one can attach 2 invariants:

# Two invariants associated to $M$

The symmetry between the automorphic and Galois side implies that to a hyperspherical  $G$ -variety  $M$ , one can attach 2 invariants:

- **Period invariant:**  $M$  gives rise to a theta function  $\Theta_M \in \mathcal{A}(G)$ . Each  $\pi \in \mathcal{A}_2(G)$  then gives:

$$\mathcal{P}_M(\pi) := \sum_{\{\phi_i\} \subset \pi} |\langle \Theta_M, \phi_i \rangle_{\text{Pet}}|^2.$$

So

$$\mathcal{P}_M : \{L^2\text{-automorphic reps of } G\} \longrightarrow \mathbb{C}.$$

# Two invariants associated to $M$

The symmetry between the automorphic and Galois side implies that to a hyperspherical  $G$ -variety  $M$ , one can attach 2 invariants:

- **Period invariant:**  $M$  gives rise to a theta function  $\Theta_M \in \mathcal{A}(G)$ . Each  $\pi \in \mathcal{A}_2(G)$  then gives:

$$\mathcal{P}_M(\pi) := \sum_{\{\phi_i\} \subset \pi} |\langle \Theta_M, \phi_i \rangle_{\text{Pet}}|^2.$$

So

$$\mathcal{P}_M : \{L^2\text{-automorphic reps of } G\} \longrightarrow \mathbb{C}.$$

- **$L$ -function-invariant:** through Galois action on tangent spaces of Galois-fixed points, get an invariant by considering special  $L$ -value

$$\mathcal{L}_M : \{\text{A-parameters valued in } G\} \longrightarrow \mathbb{C}$$

# Duality Conjecture

The above discussion led [BZSV] to the following

## Conjecture

*There is an involutive duality*

$$\{\text{hyperspherical } G\text{-var.}\} \longleftrightarrow \{\text{hyperspherical } G^\vee\text{-var.}\}$$

*such that if  $M \longleftrightarrow M^\vee$ , then*

$$\mathcal{P}_M = \mathcal{L}_{M^\vee}$$

*and*

$$\mathcal{P}_{M^\vee} = \mathcal{L}_M.$$

# Duality Conjecture

The above discussion led [BZSV] to the following

## Conjecture

*There is an involutive duality*

$$\{\text{hyperspherical } G\text{-var.}\} \longleftrightarrow \{\text{hyperspherical } G^\vee\text{-var.}\}$$

*such that if  $M \longleftrightarrow M^\vee$ , then*

$$\mathcal{P}_M = \mathcal{L}_{M^\vee}$$

*and*

$$\mathcal{P}_{M^\vee} = \mathcal{L}_M.$$

Note that in both equations, the domains of the LHS and RHS are identified by the Langlands correspondence.



# Examples of Duality [BZSV]

M	$M^\vee$
point	$T_e(U \backslash G)$
$T^*(X)$	$V_X \times^{X^\vee} G^\vee$
$T^*(\mathrm{Sp}_{2n} \backslash \mathrm{GL}_{2n})$ (symplectic period)	$T_e(\mathrm{GL}_n^\Delta U \backslash \mathrm{GL}_{2n})$ (Shalika period)
$T^*(\mathrm{SO}_{2n}^\Delta \backslash (\mathrm{SO}_{2n} \times \mathrm{SO}_{2n+1}))$ (Basic Gross-Prasad)	$V_{2n} \otimes W_{2n}$ (Equal Rank Theta Corr.)
$(V \otimes W, \mathrm{O}(V) \times \mathrm{Sp}(W))$	General GP-varieties

# Special Case (Joint with Bryan Wang)

We consider special cases of the data:

$$(\iota_X : H \times \mathrm{SL}_2 \rightarrow G, S)$$

such that

# Special Case (Joint with Bryan Wang)

We consider special cases of the data:

$$(\iota_X : H \times \mathrm{SL}_2 \rightarrow G, S)$$

such that

- $G$  is a symplectic or orthogonal group;

# Special Case (Joint with Bryan Wang)

We consider special cases of the data:

$$(\iota_X : H \times \mathrm{SL}_2 \rightarrow G, S)$$

such that

- $G$  is a symplectic or orthogonal group;
- $H = Z_G(\iota(\mathrm{SL}_2))$ ;

# Special Case (Joint with Bryan Wang)

We consider special cases of the data:

$$(\iota_X : H \times \mathrm{SL}_2 \rightarrow G, S)$$

such that

- $G$  is a symplectic or orthogonal group;
- $H = Z_G(\iota(\mathrm{SL}_2))$ ;
- $S = 0$ .

# Special Case (Joint with Bryan Wang)

We consider special cases of the data:

$$(\iota_X : H \times \mathrm{SL}_2 \rightarrow G, S)$$

such that

- $G$  is a symplectic or orthogonal group;
- $H = Z_G(\iota(\mathrm{SL}_2))$ ;
- $S = 0$ .

The associated  $G$ -variety  $M$  just depends on a unipotent conjugacy class  $e \in G$ . The quantization of  $M_e$  is a generalized Whittaker/Gelfand-Graev  $G$ -module:

$$\Pi_e = \mathrm{Ind}_{H.U}^G 1_H \otimes \psi.$$

# Results I

Our first result addresses the question: for which  $e$  is  $M_e$  hyperspherical? Note that for classical groups, nilpotent orbits are classified by partitions or Young diagrams with parity constraints (plus extra data).

Our first result addresses the question: for which  $e$  is  $M_e$  hyperspherical? Note that for classical groups, nilpotent orbits are classified by partitions or Young diagrams with parity constraints (plus extra data).

## Proposition

*Assume  $G = O_{2n}$  for simplicity. Then  $M_e$  is hyperspherical if and only if  $e$  belongs to the following list:*

- $e = [2n - r, 1^r]$ ,  $r$  odd (hook type)
- $e = (2^n)$  (Shalika type)
- $e = (3, 3)$ ,  $(4, 4)$  or  $(6, 6)$  (sporadic type)



# Results II

For those  $e$ 's of hook type or of sporadic type, our second result determines the hyperspherical dual  $M_e^\vee$ .

# Results II

For those  $e$ 's of hook type or of sporadic type, our second result determines the hyperspherical dual  $M_e^\vee$ .

## Theorem

*For  $e \in G = O_{2n}$  of hook type.*

$$M_e^\vee = M_{e^\vee},$$

*where  $e^\vee \in G^\vee = O_{2n}$  is also a nilpotent element of hook type. More precisely, the relation  $e \longleftrightarrow e^\vee$  is depicted by the following diagram.*

# Results II


For those  $e$ 's of hook type or of sporadic type, our second result determines the hyperspherical dual  $M_e^\vee$ .

## Theorem

For  $e \in G = O_{2n}$  of hook type.

$$M_e^\vee = M_{e^\vee},$$

where  $e^\vee \in G^\vee = O_{2n}$  is also a nilpotent element of hook type. More precisely, the relation  $e \longleftrightarrow e^\vee$  is depicted by the following diagram.



# Remarks on the proof

- The proof of the theorem involves resolving two spectral decomposition problems (for  $M$  and  $M^\vee$  resp.) , and showing that the answer can be described in terms of the dual variety ( $M^\vee$  and  $M$  resp.) as dictated by the BZSV conjecture.

# Remarks on the proof

- The proof of the theorem involves resolving two spectral decomposition problems (for  $M$  and  $M^\vee$  resp.) , and showing that the answer can be described in terms of the dual variety ( $M^\vee$  and  $M$  resp.) as dictated by the BZSV conjecture.
- For the hook type partitions, the solution of these spectral decomposition problems is achieved by using theta correspondence. For the sporadic types, one uses exceptional theta correspondences.

# Remarks on the proof

- The proof of the theorem involves resolving two spectral decomposition problems (for  $M$  and  $M^\vee$  resp.) , and showing that the answer can be described in terms of the dual variety ( $M^\vee$  and  $M$  resp.) as dictated by the BZSV conjecture.
- For the hook type partitions, the solution of these spectral decomposition problems is achieved by using theta correspondence. For the sporadic types, one uses exceptional theta correspondences.
- In particular, we use the results of Gomez and Zhu on the transfer of generalized Whittaker models under theta correspondence.

# Remarks on the proof

- The proof of the theorem involves resolving two spectral decomposition problems (for  $M$  and  $M^\vee$  resp.) , and showing that the answer can be described in terms of the dual variety ( $M^\vee$  and  $M$  resp.) as dictated by the BZSV conjecture.
- For the hook type partitions, the solution of these spectral decomposition problems is achieved by using theta correspondence. For the sporadic types, one uses exceptional theta correspondences.
- In particular, we use the results of Gomez and Zhu on the transfer of generalized Whittaker models under theta correspondence.
- Bryan has extended these local results to the global setting and the  $L^2$ -setting, allowing us to resolve the  $L^2$  and global version of the BZSV conjecture.

# A Geometric Principle

The proof of these spectral results have an underlying geometric incarnation, which suggests the following principle:



# A Geometric Principle

The proof of these spectral results have an underlying geometric incarnation, which suggests the following principle:

If  $\tilde{M}$  is a hyperspherical  $G \times H$ -var., with dual  $\tilde{M}^\vee$  (a  $G^\vee \times H^\vee$ -var.), then one should have a commutative diagram:

# A Geometric Principle

The proof of these spectral results have an underlying geometric incarnation, which suggests the following principle:

If  $\tilde{M}$  is a hyperspherical  $G \times H$ -var., with dual  $\tilde{M}^\vee$  (a  $G^\vee \times H^\vee$ -var.), then one should have a commutative diagram:

$$\begin{array}{ccc} \tilde{M} & \xrightarrow{\text{duality}} & \tilde{M}^\vee \\ \text{Reduction wrt } (H, pt) \downarrow & & \downarrow \text{Whittaker reduction wrt } H^\vee \\ M & \xrightarrow{\text{duality}} & M^\vee \end{array}$$

# A Geometric Principle

The proof of these spectral results have an underlying geometric incarnation, which suggests the following principle:

If  $\tilde{M}$  is a hyperspherical  $G \times H$ -var., with dual  $\tilde{M}^\vee$  (a  $G^\vee \times H^\vee$ -var.), then one should have a commutative diagram:

$$\begin{array}{ccc} \tilde{M} & \xrightarrow{\text{duality}} & \tilde{M}^\vee \\ \text{Reduction wrt } (H, pt) \downarrow & & \downarrow \text{Whittaker reduction wrt } H^\vee \\ M & \xrightarrow{\text{duality}} & M^\vee \end{array}$$

In other words,

*Hyperspherical Duality “commutes” with Reduction.*

Thank You for Your Attention!