

Explicit construction of Ramanujan bigraphs


Kathrin Maurischat, RWTH Aachen University,
on joint work with
Cristina Ballantine, Shai Evra, Brooke Feigon, Ori Parzanchevski.

2020 November 25

Biregular bigraphs

Graph $X = (L_x \cup R_x, E_x)$ satisfying

L_x R_x



bipartite : edges only between L_x - and R_x -vertices

biregular : # edges from each left vertex the same
right

Biregular bigraphs

We consider undirected graphs $X = (V_X, E_X)$ which are

- ▶ bipartite: $V_X = L_X \sqcup R_X$ such that if $(v_1, v_2) = (v_2, v_1) \in E_X$, then either $(v_1 \in L_X \text{ and } v_2 \in R_X)$ or $(v_1 \in R_X \text{ and } v_2 \in L_X)$.
- ▶ $(q_L + 1, q_R + 1)$ -biregular: $\forall v \in L_X: \#\{(v, v_2) \in E_X\} = q_L + 1$ and $\forall v \in R_X: \#\{(v_1, v) \in E_X\} = q_R + 1$.

Biregular bigraphs

We consider undirected graphs $X = (V_X, E_X)$ which are

- ▶ bipartite: $V_X = L_X \sqcup R_X$ such that if $(v_1, v_2) = (v_2, v_1) \in E_X$, then either $(v_1 \in L_X \text{ and } v_2 \in R_X)$ or $(v_1 \in R_X \text{ and } v_2 \in L_X)$.
- ▶ $(q_L + 1, q_R + 1)$ -biregular: $\forall v \in L_X: \#\{(v, v_2) \in E_X\} = q_L + 1$ and $\forall v \in R_X: \#\{(v_1, v) \in E_X\} = q_R + 1$.
- ▶ satisfying $q_L \geq q_R$, so $n_L = \#L_X \leq n_R = \#R_X$.

The adjacency matrix has got shape $\begin{pmatrix} 0 & A \\ A^t & 0 \end{pmatrix}$ with $A \in M_{n_L, n_R}(\{0, 1\})$.

Trivial eigenvalues: $\pm\sqrt{(q_L + 1)(q_R + 1)}$, 0 with multiplicity $(n_R - n_L)$.

Biregular bigraphs

We consider undirected graphs $X = (V_X, E_X)$ which are

- ▶ bipartite: $V_X = L_X \sqcup R_X$ such that if $(v_1, v_2) = (v_2, v_1) \in E_X$, then either $(v_1 \in L_X \text{ and } v_2 \in R_X)$ or $(v_1 \in R_X \text{ and } v_2 \in L_X)$.
- ▶ $(q_L + 1, q_R + 1)$ -biregular: $\forall v \in L_X: \#\{(v, v_2) \in E_X\} = q_L + 1$ and $\forall v \in R_X: \#\{(v_1, v) \in E_X\} = q_R + 1$.
- ▶ satisfying $q_L \geq q_R$, so $n_L = \#L_X \leq n_R = \#R_X$.

The adjacency matrix has got shape $\begin{pmatrix} 0 & A \\ A^t & 0 \end{pmatrix}$ with $A \in M_{n_L, n_R}(\{0, 1\})$.

Trivial eigenvalues: $\pm\sqrt{(q_L + 1)(q_R + 1)}$, 0 with multiplicity $(n_R - n_L)$.

Non-trivial eigenvalues: $\pm\lambda_2, \dots, \pm\lambda_n$, with $n \leq n_L$.

Ramanujan bigraphs

The Zeta function of a graph X is that of its fundamental group.

Ramanujan bigraphs

The Zeta function of a graph X is that of its fundamental group.

Proposition [Hashimoto 1989] For $(q_L + 1, q_R + 1)$ -biregular graphs,

$$Z_X(u)^{-1} = (1-u)^{\#E_X - n_L - n_R} (1+q_R u)^{n_L - n_R} \prod_{j=1}^n (1 - (\lambda_j^2 - q_L - q_R)u + q_L q_R u^2),$$

with trivial zeros $1, (q_L q_R)^{-1}, -q_R^{-1}$.

Ramanujan bigraphs

The Zeta function of a graph X is that of its fundamental group.

Proposition [Hashimoto 1989] For $(q_L + 1, q_R + 1)$ -biregular graphs,

$$Z_X(u)^{-1} = (1-u)^{\#E_X - n_L - n_R} (1+q_R u)^{n_L - n_R} \prod_{j=1}^n (1 - (\lambda_j^2 - q_L - q_R)u + q_L q_R u^2),$$

with trivial zeros $1, (q_L q_R)^{-1}, -q_R^{-1}$.

Riemann hypothesis The non-trivial zeros satisfy the property: If $\operatorname{Re}(s) \in (0, 1)$ and $Z_X((q_L q_R)^{-s})^{-1} = 0$, then $\operatorname{Re}(s) = 1/2$.

Ramanujan bigraphs

The Zeta function of a graph X is that of its fundamental group.

Proposition [Hashimoto 1989] For $(q_L + 1, q_R + 1)$ -biregular graphs,

$$Z_X(u)^{-1} = (1-u)^{\#E_X - n_L - n_R} (1+q_R u)^{n_L - n_R} \prod_{j=1}^n (1 - (\lambda_j^2 - q_L - q_R)u + q_L q_R u^2),$$

with trivial zeros $1, (q_L q_R)^{-1}, -q_R^{-1}$.

Riemann hypothesis The non-trivial zeros satisfy the property: If $\operatorname{Re}(s) \in (0, 1)$ and $Z_X((q_L q_R)^{-s})^{-1} = 0$, then $\operatorname{Re}(s) = 1/2$.

Definition A $(q_L + 1, q_R + 1)$ -biregular graph is a *Ramanujan bigraph* if it satisfies the Riemann hypothesis.

Alternative characterization

Proposition A finite, connected $(q_L + 1, q_R + 1)$ -bigregular bipartite graph is a Ramanujan bigraph, iff for all non-trivial eigenvalues λ

$$|\lambda^2 - q_L - q_R| \leq 2\sqrt{q_L q_R}.$$

Alternative characterization

Proposition A finite, connected $(q_L + 1, q_R + 1)$ -bigregular bipartite graph is a Ramanujan bigraph, iff for all non-trivial eigenvalues λ

$$|\lambda^2 - q_L - q_R| \leq 2\sqrt{q_L q_R}.$$

Remarks

- ▶ Interlacing families of Ramanujan bigraphs are known to satisfy extremely good expander properties. They are at the same time highly connected and sparse.

Alternative characterization

Proposition A finite, connected $(q_L + 1, q_R + 1)$ -bigregular bipartite graph is a Ramanujan bigraph, iff for all non-trivial eigenvalues λ

$$|\lambda^2 - q_L - q_R| \leq 2\sqrt{q_L q_R}.$$

Remarks

- ▶ Interlacing families of Ramanujan bigraphs are known to satisfy extremely good expander properties. They are at the same time highly connected and sparse.
- ▶ Interlacing families of Ramanujan bigraphs exist for all $q_L, q_R \geq 2$ [Marcus, Spielman, Srivastava '14].

Bruhat-Tits trees for unitary groups

Let E_p/\mathbb{Q}_p be an unramified quadratic field extension, $\Phi = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}$,
 $U_3 = U_3(E_p/\mathbb{Q}_p, \Phi)$ unitary group defined over \mathbb{Z}_p .

Bruhat-Tits trees for unitary groups

Let E_p/\mathbb{Q}_p be an unramified quadratic field extension, $\Phi = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}$, $U_3 = U_3(E_p/\mathbb{Q}_p, \Phi)$ unitary group defined over \mathbb{Z}_p . There are two conjugacy classes of maximal compact subgroups in $U_3(\mathbb{Q}_p)$,

$$K_L = U_3(\mathbb{Z}_p) \text{ and } K_R = U_3(\mathbb{Q}_p) \cap \begin{pmatrix} \mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} & p^{-1}\mathfrak{o}_{E_p} \\ \mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} & p^{-1}\mathfrak{o}_{E_p} \\ p\mathfrak{o}_{E_p} & p\mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} \end{pmatrix}.$$

Iwahori subgroup $I = K_L \cap K_R$.

Bruhat-Tits trees for unitary groups

Let E_p/\mathbb{Q}_p be an unramified quadratic field extension, $\Phi = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}$, $U_3 = U_3(E_p/\mathbb{Q}_p, \Phi)$ unitary group defined over \mathbb{Z}_p . There are two conjugacy classes of maximal compact subgroups in $U_3(\mathbb{Q}_p)$,

$$K_L = U_3(\mathbb{Z}_p) \text{ and } K_R = U_3(\mathbb{Q}_p) \cap \begin{pmatrix} \mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} & p^{-1}\mathfrak{o}_{E_p} \\ \mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} & p^{-1}\mathfrak{o}_{E_p} \\ p\mathfrak{o}_{E_p} & p\mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} \end{pmatrix}.$$

Iwahori subgroup $I = K_L \cap K_R$.

$L_{\mathcal{T}} = \{\text{conjugates of } K_L\}$,

$R_{\mathcal{T}} = \{\text{conjugates of } K_R\}$, and

$E_{\mathcal{T}} = \{(V_L, V_R) \mid V_L \cap V_R \text{ is conjugate to } I\}$.

Bruhat-Tits trees for unitary groups

Let E_p/\mathbb{Q}_p be an unramified quadratic field extension, $\Phi = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}$, $U_3 = U_3(E_p/\mathbb{Q}_p, \Phi)$ unitary group defined over \mathbb{Z}_p . There are two conjugacy classes of maximal compact subgroups in $U_3(\mathbb{Q}_p)$,

$$K_L = U_3(\mathbb{Z}_p) \text{ and } K_R = U_3(\mathbb{Q}_p) \cap \begin{pmatrix} \mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} & p^{-1}\mathfrak{o}_{E_p} \\ \mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} & p^{-1}\mathfrak{o}_{E_p} \\ p\mathfrak{o}_{E_p} & p\mathfrak{o}_{E_p} & \mathfrak{o}_{E_p} \end{pmatrix}.$$

Iwahori subgroup $I = K_L \cap K_R$.

$L_{\mathcal{T}} = \{\text{conjugates of } K_L\}$,

$R_{\mathcal{T}} = \{\text{conjugates of } K_R\}$, and

$E_{\mathcal{T}} = \{(V_L, V_R) \mid V_L \cap V_R \text{ is conjugate to } I\}$.

The graph $\mathcal{T} = \mathcal{T}_p$ is the Bruhat-Tits building (tree) of $U_3(\mathbb{Q}_p)$. It is $(p^3 + 1, p + 1)$ -biregular.

Ramanujan bigraphs from Bruhat-Tits trees

Theorem [Ballantine, Ciubotaru '11] Let Γ be a discrete, cocompact subgroup of $U_3(\mathbb{Q}_p)$ acting on \mathcal{T}_p without fixed points. Then the quotient \mathcal{T}_p/Γ is a Ramanujan bigraph **iff** $L^2(U_3(\mathbb{Q}_p)/\Gamma)^I$ satisfies the *Ramanujan hypothesis*,

Ramanujan bigraphs from Bruhat-Tits trees

Theorem [Ballantine, Ciubotaru '11] Let Γ be a discrete, cocompact subgroup of $U_3(\mathbb{Q}_p)$ acting on \mathcal{T}_p without fixed points. Then the quotient \mathcal{T}_p/Γ is a Ramanujan bigraph **iff** $L^2(U_3(\mathbb{Q}_p)/\Gamma)'$ satisfies the *Ramanujan hypothesis*, i.e.

iff every irreducible unitary representation occurring non-trivially in $L^2(U_3(\mathbb{Q}_p)/\Gamma)'$ is *tempered*.

Ramanujan bigraphs from Bruhat-Tits trees

Theorem [Ballantine, Ciubotaru '11] Let Γ be a discrete, cocompact subgroup of $U_3(\mathbb{Q}_p)$ acting on \mathcal{T}_p without fixed points. Then the quotient \mathcal{T}_p/Γ is a Ramanujan bigraph **iff** $L^2(U_3(\mathbb{Q}_p)/\Gamma)'$ satisfies the *Ramanujan hypothesis*, i.e.

iff every irreducible unitary representation occurring non-trivially in $L^2(U_3(\mathbb{Q}_p)/\Gamma)'$ is *tempered*.

Idea of proof: Identify the Zeta-function of \mathcal{T}_p/Γ with the characteristic polynomial of $L^2(U_3(\mathbb{Q}_p)/\Gamma)'$ as module of the Iwahori-Hecke algebra.

Ramanujan bigraphs from Bruhat-Tits trees

Theorem [Ballantine, Ciubotaru '11] Let Γ be a discrete, cocompact subgroup of $U_3(\mathbb{Q}_p)$ acting on \mathcal{T}_p without fixed points. Then the quotient \mathcal{T}_p/Γ is a Ramanujan bigraph **iff** $L^2(U_3(\mathbb{Q}_p)/\Gamma)'$ satisfies the *Ramanujan hypothesis*, i.e.

iff every irreducible unitary representation occurring non-trivially in $L^2(U_3(\mathbb{Q}_p)/\Gamma)'$ is *tempered*.

Idea of proof: Identify the Zeta-function of \mathcal{T}_p/Γ with the characteristic polynomial of $L^2(U_3(\mathbb{Q}_p)/\Gamma)'$ as module of the Iwahori-Hecke algebra.

For $U_3(\mathbb{Q}_p)$, the inspection of [Rogawski 1990]'s A -packets provides a tool to prove/disprove temperedness.

Ramanujan bigraphs from Bruhat-Tits trees

Theorem [BC'11] Let E/\mathbb{Q} be an imaginary quadratic extension. Let G be the unitary group associated to a division algebra D of degree three over its center E with involution of the second kind (i.e. reducing to the Galois-automorphism on E). Assume $G(\mathbb{R})$ is compact. Let p be a prime inert in E , and $(n, p) = 1$. Let

$$\Gamma_p(n) = \ker(G(\mathbb{Z}[p^{-1}]) \rightarrow G(\mathbb{Z}[p^{-1}]/n\mathbb{Z}[p^{-1}])) \subset G(\mathbb{Q}).$$

Then, the quotient $\mathcal{T}_p/\Gamma_p(n)$ is a Ramanujan bigraph.

Ramanujan property for unitary groups

Let E/\mathbb{Q} be an imaginary quadratic extension, $\Phi \in M_3(E)$ a definite hermitian matrix, $G = U_3(E/\mathbb{Q}, \Phi)$.

Ramanujan property for unitary groups

Let E/\mathbb{Q} be an imaginary quadratic extension, $\Phi \in M_3(E)$ a definite hermitian matrix, $G = U_3(E/\mathbb{Q}, \Phi)$.

Let $K = \otimes_v K_v \subset G(\mathbb{A})$ be a compact open subgroup with $K_\infty = G_\infty$, $K_p \leq K_{L,p} = G(\mathbb{Z}_p)$ of finite index and $K_p = K_{L,p}$ almost everywhere. Let $Ram(K) = \{p \mid K_p \neq K_{L,p}\}$.

Definition K is said to satisfy the *Ramanujan property* if every irreducible subrepresentations of $L^2(G(\mathbb{Q}) \backslash G(\mathbb{A}))^K$ is tempered at all local places.

Ramanujan property for unitary groups

Let E/\mathbb{Q} be an imaginary quadratic extension, $\Phi \in M_3(E)$ a definite hermitian matrix, $G = U_3(E/\mathbb{Q}, \Phi)$.

Let $K = \otimes_v K_v \subset G(\mathbb{A})$ be a compact open subgroup with $K_\infty = G_\infty$, $K_p \leq K_{L,p} = G(\mathbb{Z}_p)$ of finite index and $K_p = K_{L,p}$ almost everywhere. Let $Ram(K) = \{p \mid K_p \neq K_{L,p}\}$.

Definition K is said to satisfy the *Ramanujan property* if every irreducible subrepresentations of $L^2(G(\mathbb{Q}) \backslash G(\mathbb{A}))^K$ is tempered at all local places.

Theorem [BEFMP'20] The Ramanujan property is satisfied by K when one of the following holds.

- (1) [EP'18/20] There exists a prime $p \in Ram(E)$ such that $p \notin Ram(K)$.
- (2) $K' \subset K$ for a compact open subgroup K' satisfying the Ramanujan property.
- (3) $K \subset K'$ for a compact open subgroup K' satisfying the Ramanujan property, and for all p such that $K_p \neq K'_p$, the group K_p is a parahoric subgroup.

Specialization

Specialize to $E = \mathbb{Q}(\sqrt{-3})$ and $\Phi = \mathbf{1}_3$.

Specialization

Specialize to $E = \mathbb{Q}(\sqrt{-3})$ and $\Phi = \mathbf{1}_3$.

Let $K_3(H) = \{g \in K_{L,3} \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.

Specialization

Specialize to $E = \mathbb{Q}(\sqrt{-3})$ and $\Phi = \mathbf{1}_3$.

Let $K_3(H) = \{g \in K_{L,3} \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.

Theorem A [BEFMP'20] K satisfies the Ramanujan property in each of the following cases.

- (1) $3 \notin \text{Ram}(K)$.
- (2) $\text{Ram}(K) = \{3\}$, $K_3 = K_3(H)$.
- (3) $3 \in \text{Ram}(K)$, $K_3 = K_3(H)$, and for all $3 \neq p \in \text{Ram}(K)$ there is a parahoric subgroup contained in K_p .

Specialization

Specialize to $E = \mathbb{Q}(\sqrt{-3})$ and $\Phi = \mathbf{1}_3$.

Let $K_3(H) = \{g \in K_{L,3} \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.

Theorem A [BEFMP'20] K satisfies the Ramanujan property in each of the following cases.

- (1) $3 \notin \text{Ram}(K)$.
- (2) $\text{Ram}(K) = \{3\}$, $K_3 = K_3(H)$.
- (3) $3 \in \text{Ram}(K)$, $K_3 = K_3(H)$, and for all $3 \neq p \in \text{Ram}(K)$ there is a parahoric subgroup contained in K_p .

Theorem B [BEFMP'20] K **doesn't** satisfy the Ramanujan property in each of the following cases.

- (1) $\text{Ram}(K) = \{3\}$, $K_3 \subset K_3(3)$ (principal congruence subgroup).
- (2) $\text{Ram}(K) = \{3, q\}$, $K_3 = K_3(H)$ and $K_q = K_q(q)$.
- (3) $3 \in \text{Ram}(K)$, $K_3 = K_3(H)$, for some $3 \neq q \in \text{Ram}(K)$ it holds $K_q \subset K_q(q)$.

Recall: Cayley graphs

For a group G with set of generators S such that $k = \#S < \infty$, $S = S^{-1}$, $e \notin S$, define the **Cayley graph** $X(G, S) = (V_X, E_X)$ by

$$V_X = G \text{ and } E_X = \{(g, gs) \mid g \in G, s \in S\}.$$

Recall: Cayley graphs

For a group G with set of generators S such that $k = \#S < \infty$, $S = S^{-1}$, $e \notin S$, define the **Cayley graph** $X(G, S) = (V_X, E_X)$ by

$$V_X = G \text{ and } E_X = \{(g, gs) \mid g \in G, s \in S\}.$$

- ▶ $X(G, S)$ is k -regular, connected.
- ▶ Cycles correspond to relations in the group.
- ▶ G acts on $X(G, S)$ by multiplication from the left.

Recall: Cayley graphs

For a group G with set of generators S such that $k = \#S < \infty$, $S = S^{-1}$, $e \notin S$, define the **Cayley graph** $X(G, S) = (V_X, E_X)$ by

$$V_X = G \text{ and } E_X = \{(g, gs) \mid g \in G, s \in S\}.$$

- ▶ $X(G, S)$ is k -regular, connected.
- ▶ Cycles correspond to relations in the group.
- ▶ G acts on $X(G, S)$ by multiplication from the left.
- ▶ (Regular) Ramanujan graphs arising as quotients of SL_2 -trees admit an explicit description by Cayley graphs [Lubotzky, Philips, Sarnak, Montenegro; Cartwright, Steger, Mantero, Zappa]

Bi-Cayley graphs [BEFMP'20]

Bi-Cayley graphs [BEFMP'20]

Definition In a group G let S^1, \dots, S^L be pairwise disjoint finite subsets of equal size. The pair $(G, \sqcup_{j=1}^L S^j)$ satisfies the **bi-Cayley axioms** if:

- (1) $S = \sqcup_{j=1}^L S^j$ is a generating set for G , $S = S^{-1}$, and $e \notin S$.
- (2) For all $s, t \in S$, $s \neq t$, we have
 $s, t \in S^i$ for some $i \in \{1, \dots, L\} \iff s^{-1}t, s^{-1} \in S^j$ for some $j \in \{1, \dots, L\}$.

Bi-Cayley graphs [BEFMP'20]

Definition In a group G let S^1, \dots, S^L be pairwise disjoint finite subsets of equal size. The pair $(G, \sqcup_{j=1}^L S^j)$ satisfies the **bi-Cayley axioms** if:

- (1) $S = \sqcup_{j=1}^L S^j$ is a generating set for G , $S = S^{-1}$, and $e \notin S$.
- (2) For all $s, t \in S$, $s \neq t$, we have
 $s, t \in S^i$ for some $i \in \{1, \dots, L\} \iff s^{-1}t, s^{-1} \in S^j$ for some $j \in \{1, \dots, L\}$.

Def/Prop Let the pair $(G, \sqcup_{j=1}^L S^j)$ satisfy the bi-Cayley axioms.

- (1) Define the following equivalence relation on $G \times \{1, \dots, L\}$:

$$(g_1, j_1) \sim (g_2, j_2) \iff \begin{cases} \text{either } g_1 = g_2 \text{ and } j_1 = j_2 \\ \text{or } g_1^{-1}g_2 \in S^{j_1} \text{ and } g_2^{-1}g_1 \in S^{j_2}. \end{cases}$$

Bi-Cayley graphs [BEFMP'20]

Definition In a group G let S^1, \dots, S^L be pairwise disjoint finite subsets of equal size. The pair $(G, \sqcup_{j=1}^L S^j)$ satisfies the **bi-Cayley axioms** if:


- (1) $S = \sqcup_{j=1}^L S^j$ is a generating set for G , $S = S^{-1}$, and $e \notin S$.
- (2) For all $s, t \in S$, $s \neq t$, we have
 $s, t \in S^i$ for some $i \in \{1, \dots, L\} \iff s^{-1}t, s^{-1} \in S^j$ for some $j \in \{1, \dots, L\}$.

Def/Prop Let the pair $(G, \sqcup_{j=1}^L S^j)$ satisfy the bi-Cayley axioms.

- (1) Define the following equivalence relation on $G \times \{1, \dots, L\}$:
$$(g_1, j_1) \sim (g_2, j_2) \iff \begin{cases} \text{either } g_1 = g_2 \text{ and } j_1 = j_2 \\ \text{or } g_1^{-1}g_2 \in S^{j_1} \text{ and } g_2^{-1}g_1 \in S^{j_2}. \end{cases}$$
- (2) The associated **bi-Cayley graph** $X = BCay(G, \sqcup_{j=1}^L S^j)$ is the $(L, \#[g, i])$ -biregular bigraph with vertices $L_X = G$, $R_X = G \times \{1, \dots, L\} / \sim$ and edges $E_X = \{(g, [g, i]) \in L_X \times R_X \mid g \in G, i \in \{1, \dots, L\}\}$.

Nice example

$$G = S_3$$



$$\blacklozenge : S = S^1 \cup S^2$$

with $S^1 = \{(12)\}$, $S^2 = \{(23)\}$

$$\text{Cay}(G, S) = (2, 2) - \text{biregular},$$

6 left, 6 right
= 12-Cycle

[Whereas $\text{Cay}(S_3, S)$
= 6-Cycle]

$$\text{Cay}(G, \tilde{S}) : (3, 2) - \text{biregular}, \left\{ \begin{array}{l} 6 \text{ left, } 3 \text{ right} \end{array} \right.$$

$$\blacklozenge \cup \blacksquare : \tilde{S} = S^1 \cup S^2 \cup S^3,$$

with $S^1 = \{(12)\}$, $S^2 = \{(23)\}$, $S^3 = \{(13)\}$

Instructive example

- ▶ Let $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$.

Instructive example

- ▶ Let $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$.
- ▶ For $p \neq 3$ let $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.

Instructive example

- ▶ Let $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$.
- ▶ For $p \neq 3$ let $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.
- ▶ Let $H_p = \Lambda_p \pmod{3}$.

Instructive example

- ▶ Let $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$.
- ▶ For $p \neq 3$ let $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.
- ▶ Let $H_p = \Lambda_p \pmod{3}$.

In particular for $p = 2$, denote for $j = 0, \dots, 5$,

$$A_j = -\frac{1}{2} \begin{pmatrix} 1 & 0 & \sqrt{-3}\zeta_6^j \\ 0 & -2 & 0 \\ \sqrt{-3}\zeta_6^{-j} & 0 & 1 \end{pmatrix}, \quad B_j = -\frac{1}{2} \begin{pmatrix} 1 & \sqrt{-3}\zeta_6^j & 0 \\ \sqrt{-3}\zeta_6^{-j} & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix},$$
$$C_j = -\frac{1}{2} \begin{pmatrix} -2 & 0 & 0 \\ 0 & 1 & \sqrt{-3}\zeta_6^j \\ 0 & \sqrt{-3}\zeta_6^{-j} & 1 \end{pmatrix}.$$

Instructive example

- ▶ Let $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$.
- ▶ For $p \neq 3$ let $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.
- ▶ Let $H_p = \Lambda_p \pmod{3}$.

In particular for $p = 2$, denote for $j = 0, \dots, 5$,

$$A_j = -\frac{1}{2} \begin{pmatrix} 1 & 0 & \sqrt{-3}\zeta_6^j \\ 0 & -2 & 0 \\ \sqrt{-3}\zeta_6^{-j} & 0 & 1 \end{pmatrix}, \quad B_j = -\frac{1}{2} \begin{pmatrix} 1 & \sqrt{-3}\zeta_6^j & 0 \\ \sqrt{-3}\zeta_6^{-j} & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix},$$
$$C_j = -\frac{1}{2} \begin{pmatrix} -2 & 0 & 0 \\ 0 & 1 & \sqrt{-3}\zeta_6^j \\ 0 & \sqrt{-3}\zeta_6^{-j} & 1 \end{pmatrix}.$$

- ▶ Set $S^1 = \{A_0, A_3\}$, $S^2 = \{A_1, A_4\}$, $S^3 = \{A_2, A_5\}$, $S^4 = \{B_0, B_3\}$, $S^5 = \{B_1, B_4\}$, $S^6 = \{B_2, B_5\}$, $S^7 = \{C_0, C_3\}$, $S^8 = \{C_1, C_4\}$, $S^9 = \{C_2, C_5\}$.
- ▶ Then $(H_2, \sqcup S^j)$ satisfies the bi-Cayley axioms.

Instructive example

- ▶ Let $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$.
- ▶ For $p \neq 3$ let $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.
- ▶ Let $H_p = \Lambda_p \pmod{3}$.

In particular for $p = 2$, denote for $j = 0, \dots, 5$,

$$A_j = -\frac{1}{2} \begin{pmatrix} 1 & 0 & \sqrt{-3}\zeta_6^j \\ 0 & -2 & 0 \\ \sqrt{-3}\zeta_6^{-j} & 0 & 1 \end{pmatrix}, \quad B_j = -\frac{1}{2} \begin{pmatrix} 1 & \sqrt{-3}\zeta_6^j & 0 \\ \sqrt{-3}\zeta_6^{-j} & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix},$$
$$C_j = -\frac{1}{2} \begin{pmatrix} -2 & 0 & 0 \\ 0 & 1 & \sqrt{-3}\zeta_6^j \\ 0 & \sqrt{-3}\zeta_6^{-j} & 1 \end{pmatrix}.$$

- ▶ Set $S^1 = \{A_0, A_3\}$, $S^2 = \{A_1, A_4\}$, $S^3 = \{A_2, A_5\}$, $S^4 = \{B_0, B_3\}$, $S^5 = \{B_1, B_4\}$, $S^6 = \{B_2, B_5\}$, $S^7 = \{C_0, C_3\}$, $S^8 = \{C_1, C_4\}$, $S^9 = \{C_2, C_5\}$.
- ▶ Then $(H_2, \sqcup S^j)$ satisfies the bi-Cayley axioms.
The bi-Cayley graph $BCay(H_2, \sqcup S^j)$ is $(2^3 + 1, 2 + 1)$ -biregular with $\#L_X = 2016$ and $\#R_X = 6048$.

Instructive example

- ▶ Let $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$.
- ▶ For $p \neq 3$ let $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.
- ▶ Let $H_p = \Lambda_p \pmod{3}$.

In particular for $p = 2$, denote for $j = 0, \dots, 5$,

$$A_j = -\frac{1}{2} \begin{pmatrix} 1 & 0 & \sqrt{-3}\zeta_6^j \\ 0 & -2 & 0 \\ \sqrt{-3}\zeta_6^{-j} & 0 & 1 \end{pmatrix}, \quad B_j = -\frac{1}{2} \begin{pmatrix} 1 & \sqrt{-3}\zeta_6^j & 0 \\ \sqrt{-3}\zeta_6^{-j} & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix},$$
$$C_j = -\frac{1}{2} \begin{pmatrix} -2 & 0 & 0 \\ 0 & 1 & \sqrt{-3}\zeta_6^j \\ 0 & \sqrt{-3}\zeta_6^{-j} & 1 \end{pmatrix}.$$

- ▶ Set Setting
 $S^1 = \{A_0, A_3\}$, $S^2 = \{A_1, A_4\}$, $S^3 = \{A_2, A_5\}$, $S^4 = \{B_0, B_3\}$, $S^5 = \{B_1, B_4\}$, $S^6 = \{B_2, B_5\}$, $S^7 = \{C_0, C_3\}$, $S^8 = \{C_1, C_4\}$, $S^9 = \{C_2, C_5\}$.
- ▶ Then $(H_2, \sqcup S^j)$ satisfies the bi-Cayley axioms.
The bi-Cayley graph $BCay(H_2, \sqcup S^j)$ is $(2^3 + 1, 2 + 1)$ -biregular with $\#L_X = 2016$ and $\#R_X = 6048$.
- ▶ Works for each inert prime p like this.

Trees as bi-Cayley graphs

Proposition [BEFMP'20] For a $(q_L + 1, q_R + 1)$ -biregular tree \mathcal{T} , $q_L \geq q_R$, let Λ be a group acting isometrically on \mathcal{T} , s. th. the action is **simply transitive** on $L_{\mathcal{T}}$.

Trees as bi-Cayley graphs

Proposition [BEFMP'20] For a $(q_L + 1, q_R + 1)$ -biregular tree \mathcal{T} , $q_L \geq q_R$, let Λ be a group acting isometrically on \mathcal{T} , s. th. the action is **simply transitive** on $L_{\mathcal{T}}$.

Fix $v_0 \in L_{\mathcal{T}}$, let $v_1, \dots, v_{q_L+1} \in R_{\mathcal{T}}$ be its neighbors, define

$$S^i := \{g \in \Lambda \mid g \neq 1, \text{dist}(gv_0, v_i) = 1\}, \quad i = 1, \dots, q_L + 1.$$

Then $(\Lambda, \sqcup S^i)$ satisfies the bi-Cayley axioms, and the bi-Cayley graph $BCay(\Lambda, \sqcup S^i)$ is isomorphic to the tree \mathcal{T} .

(Simply) transitive actions

Lemma For a definite unitary group $G = U_3(E/\mathbb{Q}, \Phi)$ as before the following are equivalent.

(a) $G(\mathbb{A}) = G(\mathbb{Q}) \cdot G(\mathbb{R}) \cdot \prod_p G(\mathbb{Z}_p)$ (class number one)

(b) For each inert prime p ,

$$G(\mathbb{Q}_p) = G(\mathbb{Z}[p^{-1}]) \cdot G(\mathbb{Z}_p).$$

(c) For every inert prime p the action of $G(\mathbb{Z}[p^{-1}])$ is transitive on the left vertices $L_{\mathcal{T}_p}$ of the BT-tree.

(Simply) transitive actions

Lemma For a definite unitary group $G = U_3(E/\mathbb{Q}, \Phi)$ as before the following are equivalent.

(a) $G(\mathbb{A}) = G(\mathbb{Q}) \cdot G(\mathbb{R}) \cdot \prod_p G(\mathbb{Z}_p)$ (class number one)

(b) For each inert prime p ,

$$G(\mathbb{Q}_p) = G(\mathbb{Z}[p^{-1}]) \cdot G(\mathbb{Z}_p).$$

(c) For every inert prime p the action of $G(\mathbb{Z}[p^{-1}])$ is transitive on the left vertices $L_{\mathcal{T}_p}$ of the BT-tree.

The stabilizer of $K_{L,p} = G(\mathbb{Z}_p)$ in $G(\mathbb{Z}[p^{-1}])$ is

$$G(\mathbb{Z}) = G(\mathbb{Z}[p^{-1}]) \cap G(\mathbb{Z}_p).$$

Any subgroup Λ_p such that $G(\mathbb{Z}[p^{-1}]) = \Lambda_p \rtimes G(\mathbb{Z})$ then acts simply transitively.

Instructive example revisited

- ▶ For $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$, we have class number one.

Instructive example revisited

- ▶ For $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$, we have class number one.
For $p \neq 3$ have $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.

Instructive example revisited

- ▶ For $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$, we have class number one.
For $p \neq 3$ have $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.
- ▶ For each unramified prime p have $G(\mathbb{Z}[1/p]) = \Lambda_p \rtimes G(\mathbb{Z})$.

Instructive example revisited

- ▶ For $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$, we have class number one.
For $p \neq 3$ have $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.
- ▶ For each unramified prime p have $G(\mathbb{Z}[1/p]) = \Lambda_p \rtimes G(\mathbb{Z})$.
- ▶ So Λ_p acts simply transitively on the left vertices $L_{\mathcal{T}_p}$ of the BT-tree.

Instructive example revisited

- ▶ For $E = \mathbb{Q}(\sqrt{-3})$ and $G = U_3(E/\mathbb{Q}, \mathbf{1}_3)$, we have class number one. For $p \neq 3$ have $\Lambda_p = \{g \in G(\mathbb{Z}[1/p]) \mid g \equiv \begin{pmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{pmatrix} \pmod{3}\}$.
- ▶ For each unramified prime p have $G(\mathbb{Z}[1/p]) = \Lambda_p \rtimes G(\mathbb{Z})$.
- ▶ So Λ_p acts simply transitively on the left vertices $L_{\mathcal{T}_p}$ of the BT-tree.
- ▶ From the exact sequence $(\bmod 3)$

$$1 \rightarrow \Gamma_p \rightarrow \Lambda_p \rightarrow H_p \rightarrow 1$$

we find an explicit description of the quotient \mathcal{T}_p/Γ_p by the bi-Cayley graph $BCay(H_p, \sqcup S^j)$.

Remarks

- ▶ This example isn't a Ramanujan bigraph, but it fits onto slides.
- ▶ We have more than this one example for realizing quotients of BT-trees for U_3 by bi-Cayley graphs.
- ▶ Nevertheless, these examples are rare: First, class number one for unitary groups is exceptional. Second, the definition of H_p (i.e a congruence condition) is not obvious, but rather by chance.

Thank you!