

# Twisting moduli, meromorphy and zeros

Thomas Oliver  
School of Mathematical Sciences  
University of Nottingham

International Seminar on Automorphic Forms  
27 January 2020

# Contents

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

## 1 $L$ -functions

## 2 Converse Theorems

## 3 Cancellation of zeros

# $L$ -functions

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

- **Dirichlet series:** There are  $a_n \in \mathbb{C}$  such that  $L(s) = \sum_{n=1}^{\infty} a_n n^{-s}$ , converging absolutely for  $\operatorname{Re}(s) > 1$ .
- **Continuation:** There is an integer  $m \geq 0$  such that  $(s-1)^m L(s)$  continues to an entire function of finite order.
- **Functional equation:**  $\Lambda(s) = \overline{\omega \Lambda(1-\bar{s})}$  for some  $|\omega| = 1$  where  $\Lambda(s)$  is the completed  $L$ -function:

$$\Lambda(s) = Q^s L(s) \prod_{j=1}^k \Gamma(\lambda_j s + \mu_j), \quad Q, \lambda_j \in \mathbb{R}_{>0}, \quad \mu_j \in \mathbb{C}, \quad k \in \mathbb{Z}_{>0}.$$

- **Euler product:**  $a_1 = 1$  and  $\log L(s) = \sum_{n=2}^{\infty} b_n n^{-s}$  where  $b_n$  is supported on prime powers and  $b_n = O(n^{\theta})$  for some  $\theta < \frac{1}{2}$ .

# Degree (theory)

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

*L*-functions

Converse  
Theorems

Cancellation  
of zeros

Say  $\Lambda(s)$  is the completion of  $L(s)$ :

$$\Lambda(s) = Q^s L(s) \prod_{j=1}^k \Gamma(\lambda_j s + \mu_j).$$

- Roughly speaking, the degree of  $L$  is the number gamma functions appearing in  $\Lambda$ .
- This number is ambiguous due to the multiplication theorem for the gamma function.
- On the other hand, the sum  $d = 2 \sum_{j=1}^k \lambda_j$  is invariant and is known as the **degree**.

# Degree (practice)

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

## Completion

$$\Lambda(s) = Q^s L(s) \prod_{j=1}^k \Gamma(\lambda_j s + \mu_j).$$

## Degree

$$d = 2 \sum_{j=1}^k \lambda_j.$$

In all known examples:

- The  $\lambda_j$  can be taken to be  $\frac{1}{2}$  and  $d$  is an integer.
- The Euler factor at a good prime  $p$  is the reciprocal of a degree  $d$  polynomial in  $p^{-s}$ .

$$d = 1$$

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

We will use the following normalization of the gamma function:

$$\Gamma_{\mathbb{R}}(s) = \pi^{-s/2} \Gamma(s/2).$$

## Examples

- The Riemann zeta function  $\zeta(s)$  has gamma factor  $\Gamma_{\mathbb{R}}(s)$ .
- If  $\chi$  is an Dirichlet character of parity  $\epsilon$ , then  $L(s, \chi)$  has gamma factor  $\Gamma_{\mathbb{R}}(s + \epsilon)$ .

In 1999, Kaczorowski–Perelli proved that the degree 1 elements in the Selberg class are precisely the Dirichlet  $L$ -functions.

$$d > 1$$

If  $K$  is a number field, then  $\zeta_K(s)$  has degree  $d = [K : \mathbb{Q}]$ . In particular:

- If  $K$  is a real quadratic field, then the Dedekind zeta function  $\zeta_K(s)$  has gamma factor  $\Gamma_{\mathbb{R}}(s)^2$ .
- If  $K$  is a quadratic imaginary field, then  $\zeta_K(s)$  has gamma factor  $\Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s}\Gamma(s) = \Gamma_{\mathbb{R}}(s)\Gamma_{\mathbb{R}}(s+1)$ .

Other examples:

- If  $\rho$  is a  $d$ -dimensional Artin representation then  $L(s, \rho)$  has degree  $d$ .
- If  $\pi$  is a cuspidal automorphic representation of  $\mathrm{GL}_d(\mathbb{A}_{\mathbb{Q}})$ , then  $L(s, \pi)$  has degree  $d$ .

- Automorphic forms on  $GL(2)$  give examples of degree 2  $L$ -functions, but a complete classification remains unknown.
- We can simplify the classification problem by specifying the invariants, such as conductor and gamma factor.
- In 1936, Hecke proved that classical newforms  $f(z) = \sum_{n=1}^{\infty} a_n \exp(2\pi inz)$  on  $\Gamma_0(N)$  for  $N = 1, 2, 3, 4$  are characterized by their  $L$ -functions  $L_f(s) = \sum_{n=1}^{\infty} a_n n^{-s}$ .
- In 1949, Maass proved the analogous result for real-analytic forms.



$$N = 1$$

Recall that  $\Gamma_0(1)$  is generated by the matrices:

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Modularity with respect to the first generator follows from the Fourier expansion of  $f$ . By Mellin inversion, for  $y > 0$  we have:

$$f(iy) = \frac{1}{2\pi i} \int_{(c)} \Lambda_f(s) y^{-s} ds$$

Applying the functional equation  $\Lambda_f(s) = (-1)^{k/2} \Lambda_f(k-s)$  and shifting the contour of integration, we deduce modularity with respect to the second generator.

$$N \in \{2, 3, 4\}$$

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

For these  $N$ ,  $\Gamma_0(N)$  is generated by:

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ N & 1 \end{pmatrix}.$$

We handle the first generator as above. Modularity with respect to the second generator follows from the functional equation and the following identity:

$$\begin{pmatrix} 1 & 0 \\ N & 1 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ N & 0 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ N & 0 \end{pmatrix}^{-1}.$$

# Twisting

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- In order to extend Hecke's argument to every  $N$ , one could try writing down generators for  $\Gamma_0(N)$  and verifying modular transformation laws for each.
- In 1967, Weil discovered that this can be avoided by incorporating twisted functional equations.
- If  $L_f(s) = \sum_{n=1}^{\infty} a_n n^{-s}$  and  $\psi$  is a Dirichlet character, then we define  $L_f(s, \psi) = \sum_{n=1}^{\infty} a_n \psi(n) n^{-s}$ .
- If  $\psi$  is a primitive Dirichlet character mod  $q$  coprime to  $N$ , then the twisted functional equation may be written

$$\Lambda_f(s, \psi) = \omega(f, \psi) \Lambda_g(k - s, \bar{\psi}),$$

where  $g(z) = f(-1/Nz)$ .

## Weil, 1967

Let  $a_n, b_n$  be sequences of complex numbers such that  $a_n, b_n = O(n^\sigma)$  for some  $\sigma > 0$ , and let  $\mathcal{P}$  be a suitable set of moduli. For every primitive Dirichlet character  $\psi$  of modulus  $q \in \{1\} \cup \mathcal{P}$  assume that  $\sum_{n=1}^{\infty} a_n \psi(n) n^{-s}$  and  $\sum_{n=1}^{\infty} b_n \bar{\psi}(n) n^{-s}$  admit analytic continuation to  $\mathbb{C}$  and satisfy the twisted functional equations. Then  $f(z) = \sum_{n=1}^{\infty} a_n \exp(2\pi i n z)$  is a weight  $k$  cusp form on  $\Gamma_0(N)$  and  $g(z) = \sum_{n=1}^{\infty} b_n \exp(2\pi i n z) = f(-1/Nz)$ .

The proof is a bit tricky, and we won't review it here!

# Twisting moduli

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- Recall that Hecke's converse theorem (1936) did not require any twisting at all for  $1 \leq N \leq 4$ .
- Razar (1977) found explicit finite sets of twisting moduli, depending on  $N$ .
- Assuming an Euler product, Conrey–Farmer (1995) and Conrey–Farmer–Odgers–Snaith (2007) established twistless converse theorems for  $5 \leq N \leq 17$  and  $N = 23$ .
- Combining Conrey–Farmer and Weil, Diaconu–Perelli–Zaharescu (2002) showed that there is a single prime  $p$  such that one can take  $\mathcal{P} = \{p\}$
- Bedert–Cooper–O–Zhang (2020) established twistless converse theorems for  $N \in \{18, 20, 24\}$ .

# Modern context

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- A very well-known converse theorem for automorphic representations of  $GL(2)$  was established by Jacquet–Langlands in 1970.
- The Jacquet–Langlands converse theorem was generalised by Cogdell–Piatetski-Shapiro to  $GL(n)$  in the 1990s. Their results are central to the contemporary theory, and have been applied in the proofs of certain cases of Langlands functoriality.
- For  $n > 2$ , it is necessary to twist by representations of  $GL(n - 2)$ .
- We are interested in twistless converse theorems for  $GL(2)$ , and applications to critical zeros.

$$N = 5 \text{ (I)}$$

- The group  $\Gamma_0(5)$  is generated by the matrices

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 5 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 5 & 3 \end{pmatrix}.$$

- As before, modularity with respect to the first two matrices follows from the Fourier expansion and functional equation. It remains to verify modularity with respect to the final generator.
- Let  $\Omega_f$  denote the annihilator ideal of  $f$  in  $\mathbb{C}[\mathrm{GL}_2(\mathbb{R})^+]$ .
- The Euler product implies that  $T_2 f \equiv a_2 f \pmod{\Omega_f}$ , where  $T_2$  is the Hecke operator:

$$T_2 = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} + \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix} \in \mathbb{C}[\mathrm{GL}_2(\mathbb{R})^+].$$

## $N = 5$ (II)

We have the Heke relation:

$$\begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} + \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix} \equiv a_2 \pmod{\Omega_f}$$

Right-multiplying by  $\begin{pmatrix} 0 & -1 \\ N & 0 \end{pmatrix} \equiv \pm 1$ , we get:

$$\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} + \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 2 & 0 \\ 5 & 1 \end{pmatrix} \equiv a_2 \pmod{\Omega_f}.$$

Subtracting the above two equations gives:

$$\begin{pmatrix} 2 & 0 \\ 5 & 1 \end{pmatrix} \equiv \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix} \pmod{\Omega_f}.$$



# $N = 5$ (III)

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

We have:

$$\begin{pmatrix} 2 & 0 \\ 5 & 1 \end{pmatrix} \equiv \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix} \pmod{\Omega_f}.$$

Right-multiplying by  $\begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$  and applying  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \equiv 1$ , we get:

$$\begin{pmatrix} 2 & 1 \\ 5 & 3 \end{pmatrix} \equiv 1 \pmod{\Omega_f},$$

that is,  $f$  is modular with respect to the final generator of  $\Gamma_0(5)$ .

# Weil's lemma

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

## Definition

We say a matrix  $E \in \mathrm{SL}_2(\mathbb{R})$  is elliptic if  $|\mathrm{Tr}(E)| < 2$ .

- Elliptic matrices have a unique fixed point in the upper half-plane  $\mathcal{H}$ .
- An elliptic matrix with integer coefficients necessarily has finite order.

## Lemma

Say  $k \in \mathbb{Z}_{\geq 0}$ . If  $h : \mathcal{H} \rightarrow \mathbb{C}$  is holomorphic and satisfies  $h|_k E = h$  for some infinite order elliptic  $E$ , then  $h$  is constant.

# Proof of Weil's lemma (I)

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- Let  $P \in \mathcal{H}$  denote the unique fixed point of  $E$ .
- The **Cayley transform** given by the matrix

$$K = \frac{1}{\sqrt{P - \overline{P}}} \begin{pmatrix} 1 & -P \\ 1 & -\overline{P} \end{pmatrix}$$

takes  $\mathcal{H}$  to the open unit disk  $\mathcal{D}$  and  $P \in \mathcal{H}$  to  $0 \in \mathcal{D}$ .

- The hyperbolic circles centred at  $P$  correspond under  $K$  to the Euclidean circles

$$C_r = \{re^{it} : t \in [0, 2\pi)\} \subset \mathcal{D}, \quad r < 1.$$

.

# Proof of Weil's lemma (II)

The transformation  $R = KEK^{-1}$  on  $\mathbb{P}^1(\mathbb{C})$  has the form

$$\begin{pmatrix} e^{i\pi\theta} & 0 \\ 0 & e^{-i\pi\theta} \end{pmatrix}, \quad \theta \in \mathbb{R} \setminus \mathbb{Q}.$$

Because  $h = h|_k E$ , the function  $\tilde{h}(z) = h(K^{-1}z)$  satisfies:

$$\tilde{h}(z) = e^{ki\pi\theta} \tilde{h}(e^{2\pi i\theta} z).$$

For  $k > 0$ , this implies  $h = 0$ . For  $k = 0$ , we get:

$$\tilde{h}(Rz) = \tilde{h}(z).$$

Since the set  $\{e^{2\pi im\theta} : m \in \mathbb{Z}\}$  is dense on the unit circle, we have:

$$\tilde{h}(z) = \tilde{h}(|z|).$$

This means that  $h$  is constant on hyperbolic circles  $K^{-1}C_r$  centred at  $P$ . Because  $h$  is holomorphic, we deduce that  $h$  is constant on  $\mathcal{H}$ .

$$N = 11$$

- $\Gamma_0(11)$  is generated by the matrices:

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 11 & 6 \end{pmatrix}, \begin{pmatrix} 3 & 1 \\ 11 & 4 \end{pmatrix}.$$

- Modularity with respect to the first matrix follows from the Fourier expansion, and the second matrix follows from an argument similar to that given when  $N = 5$ .
- As for the third, Conrey–Farmer find the infinite order elliptic matrix  $\begin{pmatrix} 1 & -2/3 \\ 11/2 & -8/3 \end{pmatrix}$  which satisfies:

$$1 - \begin{pmatrix} 3 & 1 \\ 11 & 4 \end{pmatrix} \equiv \left( 1 - \begin{pmatrix} 3 & 1 \\ 11 & 4 \end{pmatrix} \right) \begin{pmatrix} 1 & -2/3 \\ 11/2 & -8/3 \end{pmatrix}.$$

- For  $k > 0$ , it therefore follows that  $f - f|_k \gamma = 0$  as required.

# Maass forms (weight 0)

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- A **weight 0 Maass form** is an eigenfunction of the hyperbolic Laplacian satisfying certain growth and automorphy properties.
- A cuspidal weight 0 Maass form with eigenvalue  $\frac{1}{4} - \nu^2$  has a Fourier expansion of the form

$$f(x + iy) = \sum_{n \neq 0} a_n \sqrt{y} K_\nu(2\pi ny) \exp(2\pi inx),$$

where  $K_\nu$  is the  $K$ -Bessel function:

$$K_\nu(u) = \frac{1}{2} \int_0^\infty e^{-|u|(t+t^{-1})/2} y^{\nu-1} dt$$

# Parity

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- A weight 0 Maass form  $f$  with Fourier expansion  $\sum_{n \neq 0} a_n \sqrt{y} K_\nu(2\pi ny) \exp(2\pi i n x)$ , is **even** (resp. **odd**) if  $a_n = a_{-n}$  (resp.  $a_n = -a_{-n}$ ).
- The **parity** of  $f$  is 0 if  $f$  is even and 1 if  $f$  is odd.
- A weight 0 Maass form with Laplace eigenvalue  $\frac{1}{4} - \nu^2$  and parity  $\epsilon$ , has gamma factor:

$$\Gamma_{\mathbb{R}}(s + \epsilon + \nu) \Gamma_{\mathbb{R}}(s + \epsilon - \nu).$$

# Maass forms (weight 1)

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- Similarly, a weight  $k$  Maass form is an eigenfunction of the weight  $k$  hyperbolic Laplacian with certain growth and automorphy properties.
- A weight  $k$  Maass form has Fourier expansion similar to those in weight 0, in which  $K_\nu$  is replaced by a more general Whittaker function.
- We say that a weight 1 Maass form  $f$  with eigenvalue  $\frac{1}{4} - \nu^2$  has parity  $\epsilon \in \{-1, 1\}$  if  $a_{-n} = \epsilon \nu a_n$ .
- A weight 1 Maass form of parity  $\epsilon$  and eigenvalue  $\frac{1}{4} - \nu^2$  has gamma factor:

$$\Gamma_{\mathbb{R}}\left(s + \frac{1 + \epsilon}{2} + \nu\right) \Gamma_{\mathbb{R}}\left(s + \frac{1 - \epsilon}{2} - \nu\right).$$

- We will only ever consider weight 0 and weight 1 Maass forms.



# Weil's lemma for real-analytic functions (I)

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- The proof of Weil's Lemma shows that it remains valid for real-analytic functions in weights  $> 0$ .
- The proof of Weil's lemma in weight 0 relied crucially on the holomorphy of  $h$ .
- In fact, there are real-analytic functions invariant under the weight 0 action of infinite order elliptic matrices that are non-constant.
- The proof of Weil's Lemma shows that if a continuous function is invariant under an infinite order elliptic then it is constant on hyperbolic circles around the fixed point.
- It follows that if a continuous function is invariant under two infinite order elliptic matrices with distinct fixed points, then it is constant.

# Weil's lemma for real-analytic functions (II)

In fact, one does not need to construct two infinite order elliptic matrices.

## Lemma

Let  $c : \mathcal{H} \rightarrow \mathbb{C}$  be a continuous function. Say there exist  $E_1, E_2 \in \mathrm{SL}_2(\mathbb{R})$  such that  $E_1$  is an elliptic matrix of infinite order with fixed point  $a \in \mathcal{H}$ ,  $a$  is not a fixed point of  $E_2$ , and  $c|_0 E_1 = c|_0 E_2 = c$ , then  $c$  is constant on  $\mathcal{H}$ .

## Proof

Indeed, since  $E_1$  is elliptic, we know that  $|\mathrm{tr}(E_1)| = |\mathrm{tr}(E_2 E_1 E_2^{-1})| < 2$ . Subsequently, we deduce that  $E_3 = E_2 E_1 E_2^{-1}$  is elliptic. Since  $E_1$  has infinite order, so does  $E_3$ .

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

# Conrey–Farmer–Maass

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

- The results of Conrey–Farmer apply immediately to weight 1 Maass forms, noting that the proof relies only on Weil’s lemma and the axiomatic properties of group actions.
- In Bedert–Cooper–O–Zhang (2020), a second matrix is found which works for weight 0 Maass forms on  $\Gamma_0(N)$  in the cases  $1 \leq N \leq 12$  and  $N \in \{16, 18\}$ .

# Critical zeros

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

For the remainder of this talk we will focus on the holomorphy assumptions of Weil's converse theorem. Our motivation for doing so is the non-trivial zeros of automorphic  $L$ -functions. Roughly speaking:

- The non-trivial zeros should be simple;
- Distinct  $L$ -functions should not have common zeros in the critical strip.

The **Grand Simplicity Hypothesis** asserts more precisely that non-real zeros in the critical strip should be linearly independent over  $\mathbb{Q}$ .

# Multiple real zeros

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

$L$ -functions may have multiple zeros on the real line, corresponding to interesting arithmetic.

## Examples

- 1 Let  $K/\mathbb{Q}$  be a number field such that  $\mathcal{O}_K$  has rank  $r > 1$ . The Dedekind zeta function  $\zeta_K(s)$  has a zero of order  $r$  at  $s = 0$ .
- 2 Let  $E/\mathbb{Q}$  be an elliptic curve of rank  $r > 1$ . The conjecture of Birch–Swinnerton-Dyer implies that the Hasse–Weil  $L$ -function  $L(E, s)$  has a zero of order  $r$  at  $s = 1$ .

# Multiple imaginary zeros

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

- Artin  $L$ -functions may have multiple zeros in the critical strip corresponding to multiplicity in the decomposition of a Galois representation.
- Indeed, let  $\rho_1$  and  $\rho_2$  be continuous complex representations of  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ . We have

$$L(s, \rho_1 \oplus \rho_2) = L(s, \rho_1)L(s, \rho_2).$$

- In the Selberg class, we say an  $L$ -function is primitive if it is not a product of lower degree  $L$ -functions.

# Common zeros

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

Given two distinct  $L$ -functions  $L_1$  (resp.  $L_2$ ) in the Selberg class, let  $d_1$  (resp.  $d_2$ ) denote the degree and let  $S_1$  (resp.  $S_2$ ) denote the set of zeros in the critical strip.

- A lower bound for the number of elements in the symmetric difference  $(S_1 \setminus S_2) \cup (S_2 \setminus S_1)$  up to a finite height was given by Murty–Murty (1994).
- Under certain orthogonality hypotheses, a lower bound for the asymmetric difference  $S_1 \setminus S_2$  when  $d_1 = d_2$  was established by Bombieri–Perelli (1998).
- When  $d_2 - d_1 \leq 0$ , a lower bound for  $S_1 \setminus S_2$  was proved by Srinivas (2003).
- Much less is known in the cases where  $d_2 - d_1 > 0$ .

$$d_2 - d_1 = 1$$

Booker, 2013

Let  $\pi_1$  (resp.  $\pi_2$ ) be a unitary cuspidal automorphic representation of  $\mathrm{GL}_{d_1}(\mathbb{A}_{\mathbb{Q}})$  (resp.  $\mathrm{GL}_{d_2}(\mathbb{A}_{\mathbb{Q}})$ ) such that  $d_2 - d_1 = 1$ , then  $\Lambda(s, \pi_2)/\Lambda(s, \pi_1)$  has infinitely many poles.

Proof idea

If the quotient had finitely many poles, then it would be a completed Dirichlet  $L$ -function. This violates the cuspidality of  $\pi_2$ , which gives rise to a primitive element in the Selberg class.

We can use degree 2 converse theorems to extend this argument to more general quotients, however we will encounter non-standard Euler factors and poles in the critical strip.



# Weil with poles

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

Booker–Krishnumurthy, 2013

Let  $\{a_n\}_{n=1}^{\infty}, \{b_n\}_{n=1}^{\infty} \subset \mathbb{C}$  satisfy  $|a_n|, |b_n| = O(n^{\sigma})$ , for some  $\sigma \in \mathbb{R}_{>0}$ . Let  $\mathcal{P}$  be a set of primes such that  $\{p \in \mathcal{P} : p \equiv u \pmod{v}\}$  is infinite for every  $u, v \in \mathbb{Z}_{>0}$  with  $(u, v) = 1$  and  $p \nmid N$  for any  $p \in \mathcal{P}$ . If, for all  $q \in \mathcal{P} \cup \{1\}$ ,  $\Lambda_f(s, \psi)$  and  $\Lambda_g(s, \bar{\psi})$  admit **meromorphic** continuation to  $\mathbb{C}$  and satisfy the functional equation, and there is a polynomial  $P(s) \in \mathbb{C}[s]$  such that  $P(s)\Lambda_f(s, \mathbf{1})$  is entire, then  $f(z) := \sum_{n=1}^{\infty} a_n \exp(2\pi i n z)$  is a modular form on  $\Gamma_0(N)$  of weight  $k$  and  $g(z) = f(-1/Nz)$ .

# Real-analytic version

Neururer–O (2018), Hochfilzer–O (2019)

Let  $\epsilon \in \{0, 1\}$ ,  $\nu \in \mathbb{C}$ , and  $a_n, b_n$  be sequences of complex numbers such that  $|a_n|, |b_n| = O(n^\sigma)$  for some  $\sigma \in \mathbb{R}_{>0}$ . Let  $\mathcal{P}$  be a set of primes such that  $\{p \in \mathcal{P} : p \equiv u \pmod{\nu}\}$  is infinite for every  $u, \nu \in \mathbb{Z}_{>0}$  with  $(u, \nu) = 1$  and  $p \nmid N$  for any  $p \in \mathcal{P}$ . If, for all  $q \in \mathcal{P} \cup \{1\}$ ,  $\Lambda_f(s, \psi)$  and  $\Lambda_g(s, \bar{\psi})$  admit **meromorphic** continuation to  $\mathbb{C}$  and satisfy the functional equation, and there is a polynomial  $P(s) \in \mathbb{C}[s]$  such that  $P(s)\Lambda_f(s, \mathbf{1})$  is entire, then  $f(z) := \sum_{n \neq 0} a_n \sqrt{y} K_\nu(2\pi ny) \exp(2\pi i nx)$  is a Maass form on  $\Gamma_0(N)$  of weight 0 and  $g(z) = f(-1/Nz)$ .

(NO) applies when  $\nu \neq 0$ ;

(HO) applies when  $\nu = 0$ .

# Sym<sup>2</sup>

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

Given a Maass form  $f$ , its symmetric square  $L$ -function is given by:

$$L(\mathrm{Sym}^2 f, s) = \zeta(2s) \sum_{n=1}^{\infty} \frac{a_{n^2}}{n^s}.$$

It's gamma factor is given by:

$$\pi^{-3s/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s+\nu}{2}\right) \Gamma\left(\frac{s-\nu}{2}\right),$$

and so  $\mathrm{Sym}^2 f$  has degree 3. Gelbart–Jacquet showed that  $\mathrm{Sym}^2 f$  defines an isobaric automorphic representation of  $\mathrm{GL}_3(\mathbb{A}_{\mathbb{Q}})$ .

$$d_2 - d_1 = 2 \text{ (I)}$$

## Neururer–O, 2018

Let  $\xi(s)$  denote the completed Riemann zeta function. If  $f$  is a (non-CM) Maass form on  $\Gamma_0(N)$ , then  $\Lambda(\text{Sym}^2 f, s)/\xi(s)$  has infinitely many poles.

## Proof idea

The quotient  $\Lambda_2(s)/\Lambda_1(s)$  has the functional equation of a Maass form with eigenvalue  $\frac{1}{4} - \nu^2$ . If it has finitely many poles, then the converse theorem of Neururer–O implies that it is the  $L$ -function of a Maass form. This gives an impossible linear dependence of Euler products.

Using different methods, Raghunathan (1999) obtained this result for modular forms  $f$ .

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

$$d_2 - d_1 = 2 \text{ (II)}$$

## Hochfilzer–O, 2019

Let  $\Lambda_1(s)$  (resp.  $\Lambda_2(s)$ ) be a completed primitive Artin  $L$ -function of degree  $d_1$  (resp.  $d_2$ ) such that  $d_2 - d_1 \leq 2$ . If the gamma factors cancel as required, then  $\Lambda_2(s)/\Lambda_1(s)$  has infinitely many poles.

## Proof idea

The quotient  $\Lambda_2(s)/\Lambda_1(s)$  has the functional equation of a Maass form with eigenvalue  $\frac{1}{4}$  or a weight 1 modular form. If the quotient has finitely many poles, then by the converse theorems of Hochfilzer–O and Booker–Krishnumurthy, it is the  $L$ -function of a Maass or modular form. We argue as before.

Let  $\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_d(\mathbb{C})$  be a continuous irreducible representation. The following are standard conjectures about the Artin  $L$ -function  $L(s, \rho)$ :

- **Weak.**  $L(s, \rho)$  continues to an entire function on  $\mathbb{C}$ .
- **Strong.** There is a cuspidal automorphic representation  $\pi$  on  $\text{GL}_n(\mathbb{A}_{\mathbb{Q}})$  such that  $L(s, \pi) = L(s, \rho)$ .

The strong Artin conjecture implies the weak.

# Twistless

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

$L$ -functions

Converse  
Theorems

Cancellation  
of zeros

- Say  $d = 2$ . If enough twists are holomorphic, then it is known that  $L(s, \rho)$  is automorphic.
- A general Artin  $L$ -function is known to be meromorphic.
- Combining this knowledge with the converse theorem of Hochfilzer–O, we deduce that if a single twist is holomorphic then  $L(s, \rho)$  is automorphic.
- If fact, if a single twist has only finitely many poles, then  $L(s, \rho)$  is automorphic.
- This reproves a result of Booker (2003).

# References

Twisting  
moduli,  
meromorphy  
and zeros

Thomas Oliver  
School of  
Mathematical  
Sciences  
University of  
Nottingham

L-functions

Converse  
Theorems

Cancellation  
of zeros

- Neururer–O, *Weil's converse theorem for Maass forms and cancellation of zeros*, Acta Arithmetica, Vol. 196, 2020.
- Bedert–Cooper–O–Zhang, *Twisting moduli for  $GL(2)$* , Journal of Number Theory, Vol. 217, pages 142–162, 2020.
- Bettin–Bober–Booker–Conrey–Lee–Molteni–O–Platt–Steiner, *A conjectural extension of Hecke's converse theorem*, Ramanujan Journal, Vol. 47, pages 659–684, 2018.
- Hochfilzer–O, *Ratios of Artin L-functions*, arXiv:1910.02821.