

# From Asai to Triple Product: Euler Systems and p-adic L-functions

Giada GROSSI

CNRS, Université Sorbonne Paris Nord and IAS, Princeton


2nd December 2025

**International Seminar on Automorphic Forms**


$T$   $p$ -adic Galois representation

$L$ -function for  $T$   $\xleftarrow{\text{Bloch-Kato}} \rightarrow$  Selmer group for  $T^*(1)$

$T$   $p$ -adic Galois representation




$T$   $p$ -adic Galois representation



(a) special value formulae/explicit reciprocity laws


$T$   $p$ -adic Galois representation



(a) special value formulae/explicit reciprocity laws


(b) bound of the Selmer group

$T$   $p$ -adic Galois representation




- (a) special value formulae/explicit reciprocity laws
- (b) bound of the Selmer group
- for  $T$  of “automorphic origin”, construction of classes inspired by *automorphic periods*

$T$   $p$ -adic Galois representation



- (a) special value formulae/explicit reciprocity laws
- (b) bound of the Selmer group
  - for  $T$  of “automorphic origin”, construction of classes inspired by *automorphic periods*
  - (a) is often a link with  $p$ -adic  $L$ -function outside interpolation region


$T$   $p$ -adic Galois representation



- (a) special value formulae/explicit reciprocity laws
- (b) bound of the Selmer group
  - for  $T$  of “automorphic origin”, construction of classes inspired by *automorphic periods*
  - (a) is often a link with  $p$ -adic  $L$ -function outside interpolation region (i.e. not a classical  $L$ -value)



$T$   $p$ -adic Galois representation



(a) special value formulae/explicit reciprocity laws

(b) bound of the Selmer group

- for  $T$  of “automorphic origin”, construction of classes inspired by *automorphic periods*
- (a) is often a link with  $p$ -adic  $L$ -function outside interpolation region (i.e. not a classical  $L$ -value)  
     $\rightsquigarrow$  often need  *$p$ -adic variation* in families of the  $p$ -adic  $L$ -function and of the classes

- Kato's ES:  $T = T_f$ ,  $f$  cuspform

- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms

- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms  
[Bertolini–Darmon–Rotger],

- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms  
[Bertolini–Darmon–Rotger],  
[Lei–Loeffler–Zerbes]

- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms  
[Bertolini–Darmon–Rotger],  
[Lei–Loeffler–Zerbes]
- diagonal cycles:  $T = T_{f_1} \otimes T_{f_2} \otimes T_g$ ,  $f_1, f_2, g$  cuspforms  
[Darmon–Rotger]

- Kato's ES:  $T = T_f, f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g, f, g$  cuspforms  
[Bertolini–Darmon–Rotger],  
[Lei–Loeffler–Zerbes]
- diagonal cycles:  $T = T_{f_1} \otimes T_{f_2} \otimes T_g, f_1, f_2, g$  cuspforms  
[Darmon–Rotger]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{GL}_2 \times \mathrm{GL}_2 (\times \mathrm{GL}_2)$$

- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms  
[Bertolini–Darmon–Rotger],  
[Lei–Loeffler–Zerbes]
- diagonal cycles:  $T = T_{f_1} \otimes T_{f_2} \otimes T_g$ ,  $f_1, f_2, g$  cuspforms  
[Darmon–Rotger]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{GL}_2 \times \mathrm{GL}_2 (\times \mathrm{GL}_2)$$

Obtain classes in the cohomology of products of modular curves where  $T = T_f \otimes T_g$  or  $T_{f_1} \otimes T_{f_2} \otimes T_g$  is realised:



- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms  
[Bertolini–Darmon–Rotger],  
[Lei–Loeffler–Zerbes]
- diagonal cycles:  $T = T_{f_1} \otimes T_{f_2} \otimes T_g$ ,  $f_1, f_2, g$  cuspforms  
[Darmon–Rotger]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{GL}_2 \times \mathrm{GL}_2 (\times \mathrm{GL}_2)$$

Obtain classes in the cohomology of products of modular curves where  $T = T_f \otimes T_g$  or  $T_{f_1} \otimes T_{f_2} \otimes T_g$  is realised:

$$\left\{ \iota_\star(\text{Eisenstein class}) \right.$$

- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms  
[Bertolini–Darmon–Rotger],  
[Lei–Loeffler–Zerbes]
- diagonal cycles:  $T = T_{f_1} \otimes T_{f_2} \otimes T_g$ ,  $f_1, f_2, g$  cuspforms  
[Darmon–Rotger]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{GL}_2 \times \mathrm{GL}_2 (\times \mathrm{GL}_2)$$

Obtain classes in the cohomology of products of modular curves where  $T = T_f \otimes T_g$  or  $T_{f_1} \otimes T_{f_2} \otimes T_g$  is realised:

$$\left\{ \begin{array}{l} \iota_\star(\text{Eisenstein class}) \\ \text{image of the modular curve in the Chow group} \end{array} \right.$$

- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms  
[Bertolini–Darmon–Rotger],  
[Lei–Loeffler–Zerbes]
- diagonal cycles:  $T = T_{f_1} \otimes T_{f_2} \otimes T_g$ ,  $f_1, f_2, g$  cuspforms  
[Darmon–Rotger]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{GL}_2 \times \mathrm{GL}_2 (\times \mathrm{GL}_2)$$

Obtain classes in the cohomology of products of modular curves where  $T = T_f \otimes T_g$  or  $T_{f_1} \otimes T_{f_2} \otimes T_g$  is realised:

$$\left\{ \begin{array}{l} \iota_\star(\text{Eisenstein class}) \\ \text{image of the modular curve in the Chow group} \end{array} \right. \quad \leftarrow \text{Rankin–Selberg integral}$$

- Kato's ES:  $T = T_f$ ,  $f$  cuspform
- Beilinson–Flach classes:  $T = T_f \otimes T_g$ ,  $f, g$  cuspforms  
[Bertolini–Darmon–Rotger],  
[Lei–Loeffler–Zerbes]
- diagonal cycles:  $T = T_{f_1} \otimes T_{f_2} \otimes T_g$ ,  $f_1, f_2, g$  cuspforms  
[Darmon–Rotger]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{GL}_2 \times \mathrm{GL}_2 (\times \mathrm{GL}_2)$$

Obtain classes in the cohomology of products of modular curves where  $T = T_f \otimes T_g$  or  $T_{f_1} \otimes T_{f_2} \otimes T_g$  is realised:

$$\begin{cases} \iota_\star(\text{Eisenstein class}) & \rightsquigarrow \text{Rankin–Selberg integral} \\ \text{image of the modular curve in the Chow group} & \rightsquigarrow \text{Ichino's formula} \end{cases}$$

# Hilbertification: $GL_2 \times GL_2 \rightsquigarrow \text{Res}_{F/\mathbb{Q}} GL_2$

# Hilbertification: $GL_2 \times GL_2 \rightsquigarrow \text{Res}_{F/\mathbb{Q}} GL_2$

$F$  real quadratic field

# Hilbertification: $GL_2 \times GL_2 \rightsquigarrow \operatorname{Res}_{F/\mathbb{Q}} GL_2$

$F$  real quadratic field

- Asai–Flach classes:  $T = T^{\text{As}}(\mathcal{F}) = \otimes \operatorname{Ind}_{F/\mathbb{Q}} T_{\mathcal{F}}$ ,  
 $\mathcal{F}$  Hilbert cuspform [Lei–Loeffler–Zerbes]

$F$  real quadratic field

- Asai–Flach classes:  $T = T^{\text{As}}(\mathcal{F}) = \otimes \text{Ind}_{F/\mathbb{Q}} T_{\mathcal{F}}$ ,  
 $\mathcal{F}$  Hilbert cuspform [Lei–Loeffler–Zerbes]
- Hirzebruch–Zagier cycles:  $T = T^{\text{As}}(\mathcal{F}) \otimes T_g$ ,  
 $\mathcal{F}$  Hilbert cuspform,  $g$  elliptic cuspform  
[Blanco-Chacón–Fornea, Fornea–Jin]



# Hilbertification: $GL_2 \times GL_2 \rightsquigarrow \text{Res}_{F/\mathbb{Q}} GL_2$

$F$  real quadratic field

- Asai–Flach classes:  $T = T^{\text{As}}(\mathcal{F}) = \otimes \text{Ind}_{F/\mathbb{Q}} T_{\mathcal{F}}$ ,  
 $\mathcal{F}$  Hilbert cuspform [Lei–Loeffler–Zerbes]
- Hirzebruch–Zagier cycles:  $T = T^{\text{As}}(\mathcal{F}) \otimes T_g$ ,  
 $\mathcal{F}$  Hilbert cuspform,  $g$  elliptic cuspform  
[Blanco-Chacón–Fornea, Fornea–Jin]

$$GL_2 \hookrightarrow GL_2 \times GL_2 (\times GL_2)$$

# Hilbertification: $\mathrm{GL}_2 \times \mathrm{GL}_2 \rightsquigarrow \mathrm{Res}_{F/\mathbb{Q}} \mathrm{GL}_2$

$F$  real quadratic field

- Asai–Flach classes:  $T = T^{\mathrm{As}}(\mathcal{F}) = \otimes \mathrm{Ind}_{F/\mathbb{Q}} T_{\mathcal{F}}$ ,  
 $\mathcal{F}$  Hilbert cuspform [Lei–Loeffler–Zerbes]
- Hirzebruch–Zagier cycles:  $T = T^{\mathrm{As}}(\mathcal{F}) \otimes T_g$ ,  
 $\mathcal{F}$  Hilbert cuspform,  $g$  elliptic cuspform  
[Blanco-Chacón–Fornea, Fornea–Jin]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{Res}_{F/\mathbb{Q}} \mathrm{GL}_2 \quad (\times \mathrm{GL}_2)$$

# Hilbertification: $\mathrm{GL}_2 \times \mathrm{GL}_2 \rightsquigarrow \mathrm{Res}_{F/\mathbb{Q}} \mathrm{GL}_2$

$F$  real quadratic field

- Asai–Flach classes:  $T = T^{\mathrm{As}}(\mathcal{F}) = \otimes \mathrm{Ind}_{F/\mathbb{Q}} T_{\mathcal{F}}$ ,  
 $\mathcal{F}$  Hilbert cuspform [Lei–Loeffler–Zerbes]
- Hirzebruch–Zagier cycles:  $T = T^{\mathrm{As}}(\mathcal{F}) \otimes T_g$ ,  
 $\mathcal{F}$  Hilbert cuspform,  $g$  elliptic cuspform  
[Blanco–Chacón–Fornea, Fornea–Jin]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{Res}_{F/\mathbb{Q}} \mathrm{GL}_2 \quad (\times \mathrm{GL}_2)$$

Obtain classes in the cohomology of the Hilbert modular surface (times the modular curve):

# Hilbertification: $\mathrm{GL}_2 \times \mathrm{GL}_2 \rightsquigarrow \mathrm{Res}_{F/\mathbb{Q}} \mathrm{GL}_2$

$F$  real quadratic field

- Asai–Flach classes:  $T = T^{\mathrm{As}}(\mathcal{F}) = \otimes \mathrm{Ind}_{F/\mathbb{Q}} T_{\mathcal{F}}$ ,  
 $\mathcal{F}$  Hilbert cuspform [Lei–Loeffler–Zerbes]
- Hirzebruch–Zagier cycles:  $T = T^{\mathrm{As}}(\mathcal{F}) \otimes T_g$ ,  
 $\mathcal{F}$  Hilbert cuspform,  $g$  elliptic cuspform  
[Blanco–Chacón–Fornea, Fornea–Jin]

$$\mathrm{GL}_2 \hookrightarrow \mathrm{Res}_{F/\mathbb{Q}} \mathrm{GL}_2 \quad (\times \mathrm{GL}_2)$$

Obtain classes in the cohomology of the Hilbert modular surface (times the modular curve):

$$\begin{cases} \iota_{\star}(\text{Eisenstein class}) & \rightsquigarrow \text{Asai integral} \\ \text{image of the modular curve in the Chow group} & \rightsquigarrow \text{Ichino's formula} \end{cases}$$

- $F$  real quadratic field,  $G = \operatorname{Res}_{F/\mathbb{Q}} \operatorname{GL}_2$

- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime

- $F$  real quadratic field,  $G = \operatorname{Res}_{F/\mathbb{Q}} \operatorname{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ ,

- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$



# Setting and notation

- $F$  real quadratic field,  $G = \operatorname{Res}_{F/\mathbb{Q}} \operatorname{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$

- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$

- $F$  real quadratic field,  $G = \operatorname{Res}_{F/\mathbb{Q}} \operatorname{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)} =$  automorphic sheaf of weight  $(k_1, k_2)$

- $F$  real quadratic field,  $G = \operatorname{Res}_{F/\mathbb{Q}} \operatorname{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)}$  = automorphic sheaf of weight  $(k_1, k_2) \Rightarrow \mathcal{F} \in H^0(X_G, \omega_G^{(k_1, k_2)}(-D))$ ,

- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)} = \text{automorphic sheaf of weight } (k_1, k_2) \Rightarrow \mathcal{F} \in H^0(X_G, \omega_G^{(k_1, k_2)}(-D))$ ,  
 $\mathcal{F}^{1-\text{ah}} \in H^1(X_G, \omega_G^{(2-k_1, k_2)}(-D))$

- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)}$  = automorphic sheaf of weight  $(k_1, k_2) \Rightarrow \mathcal{F} \in H^0(X_G, \omega_G^{(k_1, k_2)}(-D))$ ,  
 $\mathcal{F}^{1-\text{ah}} \in H^1(X_G, \omega_G^{(2-k_1, k_2)}(-D))$
- $V_{\mathcal{F}}$  associated 2-dimensional  $p$ -adic Galois representation  
(Blasius–Rogawski–Taylor) of  $G_F = \text{Gal}(\bar{\mathbb{Q}}/F)$

- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)}$  = automorphic sheaf of weight  $(k_1, k_2) \Rightarrow \mathcal{F} \in H^0(X_G, \omega_G^{(k_1, k_2)}(-D))$ ,  
 $\mathcal{F}^{1-\text{ah}} \in H^1(X_G, \omega_G^{(2-k_1, k_2)}(-D))$
- $V_{\mathcal{F}}$  associated 2-dimensional  $p$ -adic Galois representation  
 (Blasius–Rogawski–Taylor) of  $G_F = \text{Gal}(\bar{\mathbb{Q}}/F)$   
 $\rightsquigarrow V^{\text{As}}(\mathcal{F}) = \bigotimes -\text{Ind}_{G_F/G_{\mathbb{Q}}} V_{\mathcal{F}}$ , 4-dimensional rep of  $G_{\mathbb{Q}}$

- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)}$  = automorphic sheaf of weight  $(k_1, k_2) \Rightarrow \mathcal{F} \in H^0(X_G, \omega_G^{(k_1, k_2)}(-D))$ ,  
 $\mathcal{F}^{1-\text{ah}} \in H^1(X_G, \omega_G^{(2-k_1, k_2)}(-D))$
- $V_{\mathcal{F}}$  associated 2-dimensional  $p$ -adic Galois representation  
 (Blasius–Rogawski–Taylor) of  $G_F = \text{Gal}(\bar{\mathbb{Q}}/F)$   
 $\rightsquigarrow V^{\text{As}}(\mathcal{F}) = \bigotimes -\text{Ind}_{G_F/G_{\mathbb{Q}}} V_{\mathcal{F}}$ , 4-dimensional rep of  $G_{\mathbb{Q}}$   
 realised in the  $\Pi_f^{\vee}$ -isotypic part of  $H_{\text{ét},!}^2(X_{G,\bar{\mathbb{Q}}}, \mathcal{H}^{[k_1, k_2]})$



- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)} = \text{automorphic sheaf of weight } (k_1, k_2) \Rightarrow \mathcal{F} \in H^0(X_G, \omega_G^{(k_1, k_2)}(-D))$ ,  
 $\mathcal{F}^{1-\text{ah}} \in H^1(X_G, \omega_G^{(2-k_1, k_2)}(-D))$
- $V_{\mathcal{F}}$  associated 2-dimensional  $p$ -adic Galois representation  
 (Blasius–Rogawski–Taylor) of  $G_F = \text{Gal}(\bar{\mathbb{Q}}/F)$   
 $\rightsquigarrow V^{\text{As}}(\mathcal{F}) = \bigotimes -\text{Ind}_{G_F/G_{\mathbb{Q}}} V_{\mathcal{F}}$ , 4-dimensional rep of  $G_{\mathbb{Q}}$   
 realised in the  $\Pi_f^{\vee}$ -isotypic part of  $H_{\text{ét},!}^2(X_{G, \bar{\mathbb{Q}}}, \mathcal{H}^{[k_1, k_2]})$
- can think of the degenerate case  $F = \mathbb{Q} \oplus \mathbb{Q}$  as  $X_G = X_{\text{GL}_2} \times X_{\text{GL}_2}$ ,

- $F$  real quadratic field,  $G = \operatorname{Res}_{F/\mathbb{Q}} \operatorname{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)} = \text{automorphic sheaf of weight } (k_1, k_2) \Rightarrow \mathcal{F} \in H^0(X_G, \omega_G^{(k_1, k_2)}(-D))$ ,  
 $\mathcal{F}^{1-\text{ah}} \in H^1(X_G, \omega_G^{(2-k_1, k_2)}(-D))$
- $V_{\mathcal{F}}$  associated 2-dimensional  $p$ -adic Galois representation  
 (Blasius–Rogawski–Taylor) of  $G_F = \operatorname{Gal}(\bar{\mathbb{Q}}/F)$   
 $\rightsquigarrow V^{\text{As}}(\mathcal{F}) = \bigotimes -\operatorname{Ind}_{G_F/G_{\mathbb{Q}}} V_{\mathcal{F}}$ , 4-dimensional rep of  $G_{\mathbb{Q}}$   
 realised in the  $\Pi_f^{\vee}$ -isotypic part of  $H_{\text{ét},!}^2(X_{G, \bar{\mathbb{Q}}}, \mathcal{H}^{[k_1, k_2]})$
- can think of the degenerate case  $F = \mathbb{Q} \oplus \mathbb{Q}$  as  $X_G = X_{\operatorname{GL}_2} \times X_{\operatorname{GL}_2}$ ,  
 $\mathcal{F} = f_1 \otimes f_2$

- $F$  real quadratic field,  $G = \text{Res}_{F/\mathbb{Q}} \text{GL}_2$
- Fix neat level  $K \subset G(\mathbb{A}_f)$ , spherical at  $p$  fixed odd prime
- $X_G$  (a toroidal) compactification of the Hilbert modular surface of level  $K$ , the reflex field is  $\mathbb{Q}$
- $\mathcal{F}$  (cuspidal) automorphic form for  $G$  of weight  $(k_1, k_2)$ ,  $k_1 \geq k_2$ ,  $k_1 \equiv k_2 \pmod{2}$   
 $\rightsquigarrow$  generates cuspidal automorphic rep  $\Pi$  of  $G(\mathbb{A})$
- $\omega_G^{(k_1, k_2)} = \text{automorphic sheaf of weight } (k_1, k_2) \Rightarrow \mathcal{F} \in H^0(X_G, \omega_G^{(k_1, k_2)}(-D))$ ,  
 $\mathcal{F}^{1-\text{ah}} \in H^1(X_G, \omega_G^{(2-k_1, k_2)}(-D))$
- $V_{\mathcal{F}}$  associated 2-dimensional  $p$ -adic Galois representation  
 (Blasius–Rogawski–Taylor) of  $G_F = \text{Gal}(\bar{\mathbb{Q}}/F)$   
 $\rightsquigarrow V^{\text{As}}(\mathcal{F}) = \bigotimes -\text{Ind}_{G_F/G_{\mathbb{Q}}} V_{\mathcal{F}}$ , 4-dimensional rep of  $G_{\mathbb{Q}}$   
 realised in the  $\Pi_f^{\vee}$ -isotypic part of  $H_{\text{ét},!}^2(X_{G, \bar{\mathbb{Q}}}, \mathcal{H}^{[k_1, k_2]})$
- can think of the degenerate case  $F = \mathbb{Q} \oplus \mathbb{Q}$  as  $X_G = X_{\text{GL}_2} \times X_{\text{GL}_2}$ ,  
 $\mathcal{F} = f_1 \otimes f_2$  and  $\mathcal{F}^{1-\text{ah}} = f_1(-\bar{\tau}_1)f_2(\tau)$

## (A) The Rankin–Selberg/Asai case

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G$$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G$$

- Euler system classes:  $\iota_*(\text{Siegel units/Eis classes})$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G$$

- Euler system classes:  $\iota_*(\text{Siegel units/Eis classes}) \in H_{\mathrm{mot}}^3(X_G)(-j)$  for

$$0 \leq j \leq k_2 - 2$$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G$$

- Euler system classes:  $\iota_*(\text{Siegel units/Eis classes}) \in H_{\mathrm{mot}}^3(X_G)(-j)$  for

$$0 \leq j \leq k_2 - 2$$

- critical values of  $L(V^{\mathrm{As}}(\mathcal{F}), s)$ :  $s = j$  integer such that

$$k_2 \leq j \leq k_1 - 1,$$

## (A) The Rankin–Selberg/Asai case

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G$$

- Euler system classes:  $\iota_*(\text{Siegel units/Eis classes}) \in H_{\mathrm{mot}}^3(X_G)(-j)$  for

$$0 \leq j \leq k_2 - 2$$

- critical values of  $L(V^{\mathrm{As}}(\mathcal{F}), s)$ :  $s = j$  integer such that

$$k_2 \leq j \leq k_1 - 1,$$

obtained as

$$\int_{[\mathrm{GL}_2]} \iota^*(\mathcal{F}^{1-\mathrm{ah}}) \mathrm{Eis}^{(k_1-k_2)}(s)$$

(Rankin–Selberg/Asai integral)



## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G \times X_{\mathrm{GL}_2}$$

## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G \times X_{\mathrm{GL}_2}$$

- $\Delta$ -cycles: modified diagonal image of  $\mathcal{E}^{\frac{k_1+k_2+k}{2}-3} \rightarrow \mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2}$

## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G \times X_{\mathrm{GL}_2}$$

- $\Delta$ -cycles: modified diagonal image of  $\mathcal{E}^{\frac{k_1+k_2+k}{2}-3} \rightarrow \mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2}$   
 $\in \mathrm{CH}_0^{r+2}(\mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2})$

## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G \times X_{\mathrm{GL}_2}$$

- $\Delta$ -cycles: modified diagonal image of  $\mathcal{E}^{\frac{k_1+k_2+k}{2}-3} \rightarrow \mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2}$   
 $\in \mathrm{CH}_0^{r+2}(\mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2})$  for

$(k_1, k_2, k)$  balanced

## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G \times X_{\mathrm{GL}_2}$$

- $\Delta$ -cycles: modified diagonal image of  $\mathcal{E}^{\frac{k_1+k_2+k}{2}-3} \rightarrow \mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2}$   
 $\in \mathrm{CH}_0^{r+2}(\mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2})$  for

$$(k_1, k_2, k) \text{ balanced}$$

$$(\Rightarrow L(V^{\mathrm{As}}(\mathcal{F}) \otimes V_g, \frac{1}{2}) = 0)$$

## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G \times X_{\mathrm{GL}_2}$$

- $\Delta$ -cycles: modified diagonal image of  $\mathcal{E}^{\frac{k_1+k_2+k}{2}-3} \rightarrow \mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2}$   
 $\in \mathrm{CH}_0^{r+2}(\mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2})$  for

$$(k_1, k_2, k) \text{ balanced}$$

$$(\Rightarrow L(V^{\mathrm{As}}(\mathcal{F}) \otimes V_g, \frac{1}{2}) = 0)$$

- **Ichino's formula** for  $(\sqrt{\phantom{x}} \text{ of } L(V^{\mathrm{As}}(\mathcal{F}) \otimes V_g, \frac{1}{2}))$ :



## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G \times X_{\mathrm{GL}_2}$$

- $\Delta$ -cycles: modified diagonal image of  $\mathcal{E}^{\frac{k_1+k_2+k}{2}-3} \rightarrow \mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2}$   
 $\in \mathrm{CH}_0^{r+2}(\mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2})$  for

$$(k_1, k_2, k) \text{ balanced}$$

$$(\Rightarrow L(V^{\mathrm{As}}(\mathcal{F}) \otimes V_g, \frac{1}{2}) = 0)$$

- **Ichino's formula** for  $(\sqrt{\phantom{x}} \text{ of } L(V^{\mathrm{As}}(\mathcal{F}) \otimes V_g, \frac{1}{2}))$ :

$$\int_{[\mathrm{GL}_2]} \iota^*(\mathcal{F}^{1-\mathrm{ah}}) \delta_k^{k_1-k_2} g \quad \text{if } (k_1, k_2, k) \text{ F-unbalanced}$$

## (B) The (twisted) triple product case

$\mathcal{F}$  Hilbert cuspform of weight  $(k_1, k_2)$ ,  $g$  elliptic cuspform of weight  $k$

Assumptions on the levels such that  $\epsilon_v \cdot \eta_v(-1) = +1$  for all finite  $v$

$$\iota : X_{\mathrm{GL}_2} \hookrightarrow X_G \times X_{\mathrm{GL}_2}$$

- $\Delta$ -cycles: modified diagonal image of  $\mathcal{E}^{\frac{k_1+k_2+k}{2}-3} \rightarrow \mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2} \in \mathrm{CH}_0^{r+2}(\mathcal{A}^{(k_1-2, k_2-2)} \times \mathcal{E}^{k-2})$  for

$$(k_1, k_2, k) \text{ balanced}$$


$$(\Rightarrow L(V^{\mathrm{As}}(\mathcal{F}) \otimes V_g, \tfrac{1}{2}) = 0)$$

- **Ichino's formula** for  $(\sqrt{\phantom{x}} \text{ of } L(V^{\mathrm{As}}(\mathcal{F}) \otimes V_g, \tfrac{1}{2}))$ :

$$\int_{[\mathrm{GL}_2]} \iota^*(\mathcal{F}^{1-\mathrm{ah}}) \delta_k^{k_1-k_2} g \quad \text{if } (k_1, k_2, k) \text{ F-unbalanced}$$

$$\int_{[\mathrm{GL}_2]} \iota^*(\delta_{(k_1, k_2)}^k \mathcal{F}) g^{\mathrm{ah}} \quad \text{if } (k_1, k_2, k) \text{ } \mathbb{Q}\text{-unbalanced}$$

## (B) The (twisted) triple product case



**Unbalanced regions** ( $\epsilon = +1$ )

(F-unbalanced):  $k_1 \geq k + k_2$

(Q-unbalanced):  $k \geq k_1 + k_2$

In both cases

“Algebraic classes live in a region distinct from that in which the interesting  $L$ -values occur”

In both cases

“Algebraic classes live in a region distinct from that  
in which the interesting  $L$ -values occur”

$\rightsquigarrow$  vary  $\mathcal{F}$  and  $j$  (or  $\mathcal{F}$  and  $g$ ) in  $p$ -adic families

In both cases

“Algebraic classes live in a region distinct from that in which the interesting  $L$ -values occur”

$\rightsquigarrow$  vary  $\mathcal{F}$  and  $j$  (or  $\mathcal{F}$  and  $g$ ) in  $p$ -adic families

- 1 Construct  $p$ -adic  $L$ -functions interpolating  $L$ -values in “interesting regions”

In both cases

“Algebraic classes live in a region distinct from that in which the interesting  $L$ -values occur”

$\rightsquigarrow$  vary  $\mathcal{F}$  and  $j$  (or  $\mathcal{F}$  and  $g$ ) in  $p$ -adic families

- 1 Construct  $p$ -adic  $L$ -functions interpolating  $L$ -values in “interesting regions”
- 2 Vary algebraic classes in the  $p$ -adic family

In both cases

“Algebraic classes live in a region distinct from that in which the interesting  $L$ -values occur”

$\rightsquigarrow$  vary  $\mathcal{F}$  and  $j$  (or  $\mathcal{F}$  and  $g$ ) in  $p$ -adic families

- 1 Construct  $p$ -adic  $L$ -functions interpolating  $L$ -values in “interesting regions”
- 2 Vary algebraic classes in the  $p$ -adic family  
 $\rightsquigarrow$  obtain classes outside the algebraic region too



In both cases

“Algebraic classes live in a region distinct from that in which the interesting  $L$ -values occur”

$\rightsquigarrow$  vary  $\mathcal{F}$  and  $j$  (or  $\mathcal{F}$  and  $g$ ) in  $p$ -adic families

- 1 Construct  $p$ -adic  $L$ -functions interpolating  $L$ -values in “interesting regions”
- 2 Vary algebraic classes in the  $p$ -adic family  
 $\rightsquigarrow$  obtain classes outside the algebraic region too
- 3 Prove an explicit reciprocity law (often using ERL in fixed weights in algebraic region)

In both cases

“Algebraic classes live in a region distinct from that in which the interesting  $L$ -values occur”

$\rightsquigarrow$  vary  $\mathcal{F}$  and  $j$  (or  $\mathcal{F}$  and  $g$ ) in  $p$ -adic families

- 1 Construct  $p$ -adic  $L$ -functions interpolating  $L$ -values in “interesting regions”
- 2 Vary algebraic classes in the  $p$ -adic family  
 $\rightsquigarrow$  obtain classes outside the algebraic region too
- 3 Prove an explicit reciprocity law (often using ERL in fixed weights in algebraic region)
- 4 Specialise to obtain complex  $L$ -value  $\leftrightarrow \log_p / AJ_p$  of class in 2

# Main results (A)

Let  $\mathcal{U}_i \subset \mathcal{W}$  open affinoids of weight space

# Main results (A)

Let  $\mathcal{U}_i \subset \mathcal{W}$  open affinoids of weight space

$\prod$   $p$ -ordinary family over  $\mathcal{U}_1 \times \mathcal{U}_2$  of cuspidal automorphic unitary representations of  $G$

Let  $\mathcal{U}_i \subset \mathcal{W}$  open affinoids of weight space

$\underline{\Pi}$   $p$ -ordinary family over  $\mathcal{U}_1 \times \mathcal{U}_2$  of cuspidal automorphic unitary representations of  $G$

Theorem (Grossi–Loeffler–Zerbes II, 1)

Assume  $p$  splits in  $F$ . There exists  $\mathcal{L}_p^{\text{As}}(\underline{\Pi}) \in \mathcal{O}(\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times)$  such that for any  $(k_1, k_2, j) \in (\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times) \cap (\mathbb{Z}_{\geq 0}^3)$  such that  $k_2 \leq j \leq k_1 - 1$

Let  $\mathcal{U}_i \subset \mathcal{W}$  open affinoids of weight space

$\underline{\Pi}$   $p$ -ordinary family over  $\mathcal{U}_1 \times \mathcal{U}_2$  of cuspidal automorphic unitary representations of  $G$

Theorem (Grossi–Loeffler–Zerbes II, 1)

Assume  $p$  splits in  $F$ . There exists  $\mathcal{L}_p^{\text{As}}(\underline{\Pi}) \in \mathcal{O}(\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times)$  such that for any  $(k_1, k_2, j) \in (\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times) \cap (\mathbb{Z}_{\geq 0}^3)$  such that  $k_2 \leq j \leq k_1 - 1$  we have

$$\mathcal{L}_p^{\text{As}}(\underline{\Pi})(k_1, k_2, j) \sim L^{\text{imp}}(\text{As}(\underline{\Pi}[(k_1, k_2)]), j),$$

Let  $\mathcal{U}_i \subset \mathcal{W}$  open affinoids of weight space

$\Pi$   $p$ -ordinary family over  $\mathcal{U}_1 \times \mathcal{U}_2$  of cuspidal automorphic unitary representations of  $G$

## Theorem (Grossi–Loeffler–Zerbes II, 1)

Assume  $p$  splits in  $F$ . There exists  $\mathcal{L}_p^{\text{As}}(\Pi) \in \mathcal{O}(\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times)$  such that for any  $(k_1, k_2, j) \in (\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times) \cap (\mathbb{Z}_{\geq 0}^3)$  such that  $k_2 \leq j \leq k_1 - 1$  we have

$$\mathcal{L}_p^{\text{As}}(\Pi)(k_1, k_2, j) \sim L^{\text{imp}}(\text{As}(\Pi[(k_1, k_2)]), j),$$

- if  $\mathcal{F} \in \Pi[(k_1, k_2)]$ ,  $L^{\text{imp}}(\text{As}(\Pi[(k_1, k_2)]), s)$  is the previously denoted  $L(V^{\text{As}}(\mathcal{F}), s)$ , but with modified local factors at bad primes

Let  $\mathcal{U}_i \subset \mathcal{W}$  open affinoids of weight space

$\Pi$   $p$ -ordinary family over  $\mathcal{U}_1 \times \mathcal{U}_2$  of cuspidal automorphic unitary representations of  $G$

## Theorem (Grossi–Loeffler–Zerbes II, 1)

Assume  $p$  splits in  $F$ . There exists  $\mathcal{L}_p^{\text{As}}(\Pi) \in \mathcal{O}(\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times)$  such that for any  $(k_1, k_2, j) \in (\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times) \cap (\mathbb{Z}_{\geq 0}^3)$  such that  $k_2 \leq j \leq k_1 - 1$  we have

$$\mathcal{L}_p^{\text{As}}(\Pi)(k_1, k_2, j) \sim L^{\text{imp}}(\text{As}(\Pi[(k_1, k_2)]), j),$$

- if  $\mathcal{F} \in \Pi[(k_1, k_2)]$ ,  $L^{\text{imp}}(\text{As}(\Pi[(k_1, k_2)]), s)$  is the previously denoted  $L(V^{\text{As}}(\mathcal{F}), s)$ , but with modified local factors at bad primes
- it specialises to the one variable  $p$ -adic  $L$ -function  $L_p^{\text{As}}(\Pi[(k_1, k_2)], -)$  constructed in GLZ-I



Let  $\mathcal{U}_i \subset \mathcal{W}$  open affinoids of weight space

$\Pi$   $p$ -ordinary family over  $\mathcal{U}_1 \times \mathcal{U}_2$  of cuspidal automorphic unitary representations of  $G$

## Theorem (Grossi–Loeffler–Zerbes II, 1)

Assume  $p$  splits in  $F$ . There exists  $\mathcal{L}_p^{\text{As}}(\Pi) \in \mathcal{O}(\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times)$  such that for any  $(k_1, k_2, j) \in (\mathcal{U}_1 \times \mathcal{U}_2 \times \mathbb{Z}_p^\times) \cap (\mathbb{Z}_{\geq 0}^3)$  such that  $k_2 \leq j \leq k_1 - 1$  we have

$$\mathcal{L}_p^{\text{As}}(\Pi)(k_1, k_2, j) \sim L^{\text{imp}}(\text{As}(\Pi[(k_1, k_2)]), j),$$

- if  $\mathcal{F} \in \Pi[(k_1, k_2)]$ ,  $L^{\text{imp}}(\text{As}(\Pi[(k_1, k_2)]), s)$  is the previously denoted  $L(V^{\text{As}}(\mathcal{F}), s)$ , but with modified local factors at bad primes
- it specialises to the one variable  $p$ -adic  $L$ -function  $L_p^{\text{As}}(\Pi[(k_1, k_2)], -)$  constructed in GLZ-I
- the Rankin–Selberg case is due to Hida and Panchishkin

## Theorem (GLZ II - explicit reciprocity law in the geometric region)

*For each  $j \in \mathbb{Z}$  with  $0 \leq j \leq k_2 - 2$ , the Bloch–Kato logarithm of the Asai–Flach class*

$$\mathrm{AF}^{[\Pi, j]} \in H^1(\mathbb{Q}, V_p^{\mathrm{As}}(\Pi)^*(-j))$$

*can be expressed in terms of the  $p$ -adic  $L$ -value  $L_p^{\mathrm{As}}(\Pi, 1 + j)$ .*

## Theorem (GLZ II - explicit reciprocity law in the geometric region)

*For each  $j \in \mathbb{Z}$  with  $0 \leq j \leq k_2 - 2$ , the Bloch–Kato logarithm of the Asai–Flach class*

$$\mathrm{AF}^{[\Pi,j]} \in H^1(\mathbb{Q}, V_p^{\mathrm{As}}(\Pi)^*(-j))$$

*can be expressed in terms of the  $p$ -adic  $L$ -value  $L_p^{\mathrm{As}}(\Pi, 1 + j)$ .*

## Theorem (Lei–Loeffler–Zerbes, Loeffler–Sheth, 2)

*The Euler system with base class  $\mathrm{AF}_1 = \mathrm{AF}^{[\Pi,j]}$  varies in  $p$ -adic families to a system  $\mathrm{AF}_n^{[\Pi,j]}$*

## Theorem (GLZ II - explicit reciprocity law in the geometric region)

For each  $j \in \mathbb{Z}$  with  $0 \leq j \leq k_2 - 2$ , the Bloch–Kato logarithm of the Asai–Flach class

$$\mathrm{AF}^{[\Pi, j]} \in H^1(\mathbb{Q}, V_p^{\mathrm{As}}(\Pi)^*(-j))$$

can be expressed in terms of the  $p$ -adic  $L$ -value  $L_p^{\mathrm{As}}(\Pi, 1 + j)$ .

## Theorem (Lei–Loeffler–Zerbes, Loeffler–Sheth, 2)

The Euler system with base class  $\mathrm{AF}_1 = \mathrm{AF}^{[\Pi, j]}$  varies in  $p$ -adic families to a system  $\mathrm{AF}_n^{[\underline{\Pi}, j]}$

With these results, we get 3 (i.e. big log of  $\mathrm{AF}_1^{[\underline{\Pi}, j]}$  in terms of  $\mathcal{L}_p^{\mathrm{As}}(\underline{\Pi})$ )

## Theorem (GLZ II - explicit reciprocity law in the geometric region)

For each  $j \in \mathbb{Z}$  with  $0 \leq j \leq k_2 - 2$ , the Bloch–Kato logarithm of the Asai–Flach class

$$\mathrm{AF}^{[\Pi, j]} \in H^1(\mathbb{Q}, V_p^{\mathrm{As}}(\Pi)^*(-j))$$

can be expressed in terms of the  $p$ -adic  $L$ -value  $L_p^{\mathrm{As}}(\Pi, 1 + j)$ .

## Theorem (Lei–Loeffler–Zerbes, Loeffler–Sheth, 2)

The Euler system with base class  $\mathrm{AF}_1 = \mathrm{AF}^{[\Pi, j]}$  varies in  $p$ -adic families to a system  $\mathrm{AF}_n^{[\underline{\Pi}, j]}$

With these results, we get 3 (i.e. big log of  $\mathrm{AF}_1^{[\underline{\Pi}, j]}$  in terms of  $\mathcal{L}_p^{\mathrm{As}}(\underline{\Pi})$ )

## Theorem (GLZ II, 4 and Bloch–Kato)

$$\mathcal{L}_p^{\mathrm{As}}(\underline{\Pi})(k_1, k_2, 1 + j) \neq 0 \Rightarrow \mathrm{AF}_1^{[\underline{\Pi}, j]}(k_1, k_2, j) \neq 0.$$

## Theorem (GLZ II - explicit reciprocity law in the geometric region)

For each  $j \in \mathbb{Z}$  with  $0 \leq j \leq k_2 - 2$ , the Bloch–Kato logarithm of the Asai–Flach class

$$\mathrm{AF}^{[\Pi, j]} \in H^1(\mathbb{Q}, V_p^{\mathrm{As}}(\Pi)^*(-j))$$

can be expressed in terms of the  $p$ -adic  $L$ -value  $L_p^{\mathrm{As}}(\Pi, 1 + j)$ .

## Theorem (Lei–Loeffler–Zerbes, Loeffler–Sheth, 2)

The Euler system with base class  $\mathrm{AF}_1 = \mathrm{AF}^{[\Pi, j]}$  varies in  $p$ -adic families to a system  $\mathrm{AF}_n^{[\Pi, j]}$

With these results, we get 3 (i.e. big log of  $\mathrm{AF}_1^{[\Pi, j]}$  in terms of  $\mathcal{L}_p^{\mathrm{As}}(\Pi)$ )

## Theorem (GLZ II, 4 and Bloch–Kato)

$$\mathcal{L}_p^{\mathrm{As}}(\Pi)(k_1, k_2, 1 + j) \neq 0 \Rightarrow \mathrm{AF}_1^{[\Pi, j]}(k_1, k_2, j) \neq 0.$$

Under some technical assumptions on  $\mathcal{F}$ , for  $k_2 - 1 \leq j \leq k_1 - 2$  (since  $L(V^{\mathrm{As}}(\mathcal{F}), j + 1) \neq 0$ ),

the Bloch–Kato Selmer group of  $(V^{\mathrm{As}}(\mathcal{F}))^*(-j)$  is zero.

Let  $\Pi$   $p$ -ordinary family cuspidal automorphic unitary representations of  $G \times \mathrm{GL}_2$


Let  $\Pi$   $p$ -ordinary family cuspidal automorphic unitary representations of  $G \times \mathrm{GL}_2$

$$G = \mathrm{GL}_2 \times \mathrm{GL}_2$$



Let  $\Pi$   $p$ -ordinary family cuspidal automorphic unitary representations of  $G \times \mathrm{GL}_2$

$$G = \mathrm{GL}_2 \times \mathrm{GL}_2$$



**Theorem (Darmon–Rotger, 1)**

*There exist  $p$ -adic  $L$ -functions*

$$\mathcal{L}_p^{f_2}(\Pi), \mathcal{L}_p^{f_1}(\Pi), \mathcal{L}_p^g(\Pi)$$


*interpolating the value*

$$L(\Pi[(k_1, k_2, k)], \tfrac{1}{2})$$

*for  $(k_1, k_2, k)$  in the corresponding regions.*

Let  $\Pi$   $p$ -ordinary family cuspidal automorphic unitary representations of  $G \times \mathrm{GL}_2$

$$G = \mathrm{Res}_{F/\mathbb{Q}} \mathrm{GL}_2$$



**Theorem (Blanco-Chacón–Fornea, 1)**

*There exists a  $p$ -adic  $L$ -function*

$$\mathcal{L}_p^g(\Pi)$$


*interpolating the value*

$$L(\Pi[(k_1, k_2, k)], \tfrac{1}{2})$$

*for  $(k_1, k_2, k)$  in the  $\mathbb{Q}$ -dominant region.*


# Main results (B) - $\mathbb{Q}$ -dominant region

In both cases:



# Main results (B) - $\mathbb{Q}$ -dominant region


In both cases:



2 varying  $\Delta$  in family of parallel weights  
 $k_1 = k_2$

# Main results (B) - $\mathbb{Q}$ -dominant region


In both cases:



- 2 varying  $\Delta$  in family of parallel weights  
 $k_1 = k_2$
- 3  $p$ -adic Gross–Zagier formula

# Main results (B) - $\mathbb{Q}$ -dominant region


In both cases:



- 2 varying  $\Delta$  in family of parallel weights  
 $k_1 = k_2$
- 3  $p$ -adic Gross–Zagier formula
- 4 specialise in weight  $(1, 1, 2)$  (where  $\mathcal{L}_p^g(\Pi)$  interpolates classical value)

# Main results (B) - $\mathbb{Q}$ -dominant region


In both cases:



- 2 varying  $\Delta$  in family of parallel weights  
 $k_1 = k_2$
- 3  $p$ -adic Gross–Zagier formula
- 4 specialise in weight  $(1, 1, 2)$  (where  $\mathcal{L}_p^g(\Pi)$  interpolates classical value)  
 $\rightsquigarrow$  non-vanishing of  $L$ -value gives 4 linearly-independent non-Selmer classes

# Main results (B) - $\mathbb{Q}$ -dominant region

In both cases:




- 2 varying  $\Delta$  in family of parallel weights  
 $k_1 = k_2$
- 3  $p$ -adic Gross–Zagier formula
- 4 specialise in weight  $(1, 1, 2)$  (where  $\mathcal{L}_p^g(\Pi)$  interpolates classical value)  
 $\rightsquigarrow$  non-vanishing of  $L$ -value gives 4 linearly-independent non-Selmer classes

$$(\underline{\Delta}(f_1^\alpha, f_2^\alpha), \underline{\Delta}(f_1^\alpha, f_2^\beta), \underline{\Delta}(f_1^\beta, f_2^\alpha), \underline{\Delta}(f_1^\beta, f_2^\beta))$$



# Main results (B) - $\mathbb{Q}$ -dominant region

In both cases:



- 2 varying  $\Delta$  in family of parallel weights  
 $k_1 = k_2$
- 3  $p$ -adic Gross–Zagier formula
- 4 specialise in weight  $(1, 1, 2)$  (where  $\mathcal{L}_p^g(\Pi)$  interpolates classical value)  
 $\rightsquigarrow$  non-vanishing of  $L$ -value gives 4 linearly-independent non-Selmer classes


$$(\underline{\Delta}(f_1^\alpha, f_2^\alpha), \underline{\Delta}(f_1^\alpha, f_2^\beta), \underline{\Delta}(f_1^\beta, f_2^\alpha), \underline{\Delta}(f_1^\beta, f_2^\beta))$$

## Theorem (Darmon–Rotger, Fornea–Jin, rank 0-twisted BSD)

Let  $E/\mathbb{Q}$  an elliptic curve with good ordinary reduction at  $p$   
and  $\rho$  the Artin rep attached to  $(f_1, f_2)$  (resp.  $\mathcal{F}$ ) ordinary and of weight  $(1, 1)$   
factoring through  $H$

# Main results (B) - $\mathbb{Q}$ -dominant region

In both cases:



- 2 varying  $\Delta$  in family of parallel weights  
 $k_1 = k_2$
- 3  $p$ -adic Gross–Zagier formula
- 4 specialise in weight  $(1, 1, 2)$  (where  $\mathcal{L}_p^g(\Pi)$  interpolates classical value)  
 $\rightsquigarrow$  non-vanishing of  $L$ -value gives 4 linearly-independent non-Selmer classes

$$(\underline{\Delta}(f_1^\alpha, f_2^\alpha), \underline{\Delta}(f_1^\alpha, f_2^\beta), \underline{\Delta}(f_1^\beta, f_2^\alpha), \underline{\Delta}(f_1^\beta, f_2^\beta))$$


## Theorem (Darmon–Rotger, Fornea–Jin, rank 0-twisted BSD)

Let  $E/\mathbb{Q}$  an elliptic curve with good ordinary reduction at  $p$   
and  $\rho$  the Artin rep attached to  $(f_1, f_2)$  (resp.  $\mathcal{F}$ ) ordinary and of weight  $(1, 1)$   
factoring through  $H$

$$L(E, \rho, 1) \neq 0 \Rightarrow E(H)^p = \{0\}$$


## Main results (B) - $F$ -dominant region

If  $G = \mathrm{GL}_2 \times \mathrm{GL}_2$ , similar strategy:



## Main results (B) - $F$ -dominant region


If  $G = \mathrm{GL}_2 \times \mathrm{GL}_2$ , similar strategy:



Specialising in weight  $(2,1,1)$ :

## Main results (B) - $F$ -dominant region

If  $G = \mathrm{GL}_2 \times \mathrm{GL}_2$ , similar strategy:




Specialising in weight  $(2,1,1)$ :

$\mathcal{L}_p^{f_1}(\underline{\Pi})(2,1,1) \neq 0 \Rightarrow$  two linearly independent Selmer classes

## Main results (B) - $F$ -dominant region

If  $G = \mathrm{GL}_2 \times \mathrm{GL}_2$ , similar strategy:




Specialising in weight  $(2,1,1)$ :

$\mathcal{L}_p^{f_1}(\underline{\Pi})(2,1,1) \neq 0 \Rightarrow$  two linearly independent Selmer classes

$$(\underline{\Delta}(f_1^\alpha, f_2^\alpha), \underline{\Delta}(f_1^\alpha, f_2^\beta))$$

## Main results (B) - $F$ -dominant region

If  $G = \mathrm{GL}_2 \times \mathrm{GL}_2$ , similar strategy:



Specialising in weight  $(2, 1, 1)$ :

$\mathcal{L}_p^{f_1}(\Pi)(2, 1, 1) \neq 0 \Rightarrow$  two linearly independent Selmer classes


$$(\underline{\Delta}(f_1^\alpha, f_2^\alpha), \underline{\Delta}(f_1^\alpha, f_2^\beta))$$

### Theorem (Darmon–Rotger, rank $\geq 2$ -twisted BSD)

Let  $E/\mathbb{Q}$  an elliptic curve with good ordinary reduction at  $p$   
and  $\rho$  the Artin rep attached to  $(f_1, f_2)$  ordinary and of weight  $(1, 1)$

## Main results (B) - $F$ -dominant region

If  $G = \mathrm{GL}_2 \times \mathrm{GL}_2$ , similar strategy:



Specialising in weight  $(2,1,1)$ :

$\mathcal{L}_p^{f_1}(\Pi)(2,1,1) \neq 0 \Rightarrow$  two linearly independent Selmer classes

$$(\underline{\Delta}(f_1^\alpha, f_2^\alpha), \underline{\Delta}(f_1^\alpha, f_2^\beta))$$

### Theorem (Darmon–Rotger, rank $\geq 2$ -twisted BSD)


Let  $E/\mathbb{Q}$  an elliptic curve with good ordinary reduction at  $p$   
and  $\rho$  the Artin rep attached to  $(f_1, f_2)$  ordinary and of weight  $(1,1)$

$$L(E, \rho, 1) = 0 \text{ and } \mathcal{L}_p^{f_1}(\Pi)(2,1,1) \neq 0$$



## Main results (B) - $F$ -dominant region

If  $G = \mathrm{GL}_2 \times \mathrm{GL}_2$ , similar strategy:



Specialising in weight  $(2,1,1)$ :

$\mathcal{L}_p^{f_1}(\Pi)(2,1,1) \neq 0 \Rightarrow$  two linearly independent Selmer classes

$$(\underline{\Delta}(f_1^\alpha, f_2^\alpha), \underline{\Delta}(f_1^\alpha, f_2^\beta))$$

### Theorem (Darmon–Rotger, rank $\geq 2$ -twisted BSD)

Let  $E/\mathbb{Q}$  an elliptic curve with good ordinary reduction at  $p$   
and  $\rho$  the Artin rep attached to  $(f_1, f_2)$  ordinary and of weight  $(1, 1)$

$$L(E, \rho, 1) = 0 \text{ and } \mathcal{L}_p^{f_1}(\Pi)(2, 1, 1) \neq 0 \Rightarrow \mathrm{Sel}_\rho(E, \rho) \text{ has rank } \geq 2.$$

### Rank $\geq 2$ Hilbert-twisted BSD


Let  $E/\mathbb{Q}$  an elliptic curve with good ordinary reduction at  $p$   
and  $\rho$  the Artin rep attached to  $\mathcal{F}$  ordinary and of weight  $(1, 1)$

$L(E, \rho, 1) = 0$  and  $p$ -adic  $L$ -value non vanishing  $\stackrel{?}{\Rightarrow}$   $\text{Sel}_p(E, \rho)$  has rank  $\geq 2$ .

## Rank $\geq 2$ Hilbert-twisted BSD

Let  $E/\mathbb{Q}$  an elliptic curve with good ordinary reduction at  $p$   
and  $\rho$  the Artin rep attached to  $\mathcal{F}$  ordinary and of weight  $(1, 1)$


$L(E, \rho, 1) = 0$  and  $p$ -adic  $L$ -value non vanishing  $\stackrel{?}{\Rightarrow}$   $\text{Sel}_p(E, \rho)$  has rank  $\geq 2$ .



## Rank $\geq 2$ Hilbert-twisted BSD

Let  $E/\mathbb{Q}$  an elliptic curve with good ordinary reduction at  $p$   
and  $\rho$  the Artin rep attached to  $\mathcal{F}$  ordinary and of weight  $(1, 1)$

$L(E, \rho, 1) = 0$  and  $p$ -adic  $L$ -value non vanishing  $\stackrel{?}{\Rightarrow}$   $\text{Sel}_p(E, \rho)$  has rank  $\geq 2$ .



## Theorem (Grossi–Graham, in progress)

There exists a  $p$ -adic  $L$ -function

$$\mathcal{L}_p^{\mathcal{F}}(\Pi)$$

interpolating the value

$$L(\Pi[(k_1, k_2, k)], \tfrac{1}{2})$$

for  $(k_1, k_2, k)$  in the  $F$ -dominant region.

- $\underline{\Pi}$  is a  $p$ -ordinary family of unitary cuspidal automorphic representations of  $G \times \mathrm{GL}_2$  over open affinoid of (three variable) weight space

- $\underline{\Pi}$  is a  $p$ -ordinary family of unitary cuspidal automorphic representations of  $G \times \mathrm{GL}_2$  over open affinoid of (three variable) weight space
- $p$  split in  $F$  and conductor assumptions

- $\Pi$  is a  $p$ -ordinary family of unitary cuspidal automorphic representations of  $G \times \mathrm{GL}_2$  over open affinoid of (three variable) weight space
- $p$  split in  $F$  and conductor assumptions
- Recent work of Loeffler-Kazi constructs similar  $p$ -adic  $L$ -function for overconvergent families

- $\Pi$  is a  $p$ -ordinary family of unitary cuspidal automorphic representations of  $G \times \mathrm{GL}_2$  over open affinoid of (three variable) weight space
- $p$  split in  $F$  and conductor assumptions
- Recent work of Loeffler-Kazi constructs similar  $p$ -adic  $L$ -function for overconvergent families

More generally, we construct  $p$ -adic  $L$ -functions in  $3d$ -variables for ordinary families of car of

$$\mathrm{Res}_{E/L} \mathrm{GL}_2,$$

where  $L$  is totally real of degree  $d$  and  $E$  is an étale cubic algebra.



- $\Pi$  is a  $p$ -ordinary family of unitary cuspidal automorphic representations of  $G \times \mathrm{GL}_2$  over open affinoid of (three variable) weight space
- $p$  split in  $F$  and conductor assumptions
- Recent work of Loeffler-Kazi constructs similar  $p$ -adic  $L$ -function for overconvergent families

More generally, we construct  $p$ -adic  $L$ -functions in  $3d$ -variables for ordinary families of car of

$$\mathrm{Res}_{E/L} \mathrm{GL}_2,$$

where  $L$  is totally real of degree  $d$  and  $E$  is an étale cubic algebra. It interpolates the central value of  $L(\Pi[k], r, s)$  for

$$k_{\tau_1} \geq k_{\tau_2} + k_{\tau_3} \text{ for every } \tau \in \Sigma_{\infty, L}, \tau_1, \tau_2, \tau_3 \mid \tau.$$

where  $r : {}^L G \rightarrow (\mathbb{C}^2)^{\otimes 3}$  with  $\mathrm{Gal}(E/L)$  acting as  $\mathrm{Gal}(E/L) \rightarrow \mathfrak{S}_3$ .

- $\Pi$  is a  $p$ -ordinary family of unitary cuspidal automorphic representations of  $G \times \mathrm{GL}_2$  over open affinoid of (three variable) weight space
- $p$  split in  $F$  and conductor assumptions
- Recent work of Loeffler-Kazi constructs similar  $p$ -adic  $L$ -function for overconvergent families

More generally, we construct  $p$ -adic  $L$ -functions in  $3d$ -variables for ordinary families of car of

$$\mathrm{Res}_{E/L} \mathrm{GL}_2,$$

where  $L$  is totally real of degree  $d$  and  $E$  is an étale cubic algebra. It interpolates the central value of  $L(\Pi[k], r, s)$  for

$$k_{\tau_1} \geq k_{\tau_2} + k_{\tau_3} \text{ for every } \tau \in \Sigma_{\infty, L}, \tau_1, \tau_2, \tau_3 \mid \tau.$$

where  $r : {}^L G \rightarrow (\mathbb{C}^2)^{\otimes 3}$  with  $\mathrm{Gal}(E/L)$  acting as  $\mathrm{Gal}(E/L) \rightarrow \mathfrak{S}_3$ .

For arithmetic applications, need 3:

- $\Pi$  is a  $p$ -ordinary family of unitary cuspidal automorphic representations of  $G \times \mathrm{GL}_2$  over open affinoid of (three variable) weight space
- $p$  split in  $F$  and conductor assumptions
- Recent work of Loeffler-Kazi constructs similar  $p$ -adic  $L$ -function for overconvergent families

More generally, we construct  $p$ -adic  $L$ -functions in  $3d$ -variables for ordinary families of car of

$$\mathrm{Res}_{E/L} \mathrm{GL}_2,$$

where  $L$  is totally real of degree  $d$  and  $E$  is an étale cubic algebra. It interpolates the central value of  $L(\Pi[k], r, s)$  for

$$k_{\tau_1} \geq k_{\tau_2} + k_{\tau_3} \text{ for every } \tau \in \Sigma_{\infty, L}, \tau_1, \tau_2, \tau_3 \mid \tau.$$

where  $r : {}^L G \rightarrow (\mathbb{C}^2)^{\otimes 3}$  with  $\mathrm{Gal}(E/L)$  acting as  $\mathrm{Gal}(E/L) \rightarrow \mathfrak{S}_3$ .

For arithmetic applications, need 3:

- compute  $AJ_p(\Delta)(\omega_g \otimes \eta_{\mathcal{F}})$  (as opposed to  $AJ_p(\Delta)(\eta_g \otimes \omega_{\mathcal{F}})$  in [BCF])

- $\Pi$  is a  $p$ -ordinary family of unitary cuspidal automorphic representations of  $G \times \mathrm{GL}_2$  over open affinoid of (three variable) weight space
- $p$  split in  $F$  and conductor assumptions
- Recent work of Loeffler-Kazi constructs similar  $p$ -adic  $L$ -function for overconvergent families

More generally, we construct  $p$ -adic  $L$ -functions in  $3d$ -variables for ordinary families of car of

$$\mathrm{Res}_{E/L} \mathrm{GL}_2,$$

where  $L$  is totally real of degree  $d$  and  $E$  is an étale cubic algebra. It interpolates the central value of  $L(\Pi[k], r, s)$  for

$$k_{\tau_1} \geq k_{\tau_2} + k_{\tau_3} \text{ for every } \tau \in \Sigma_{\infty, L}, \tau_1, \tau_2, \tau_3 \mid \tau.$$

where  $r : {}^L G \rightarrow (\mathbb{C}^2)^{\otimes 3}$  with  $\mathrm{Gal}(E/L)$  acting as  $\mathrm{Gal}(E/L) \rightarrow \mathfrak{S}_3$ .

For arithmetic applications, need 3:

- compute  $AJ_p(\Delta)(\omega_g \otimes \eta_{\mathcal{F}})$  (as opposed to  $AJ_p(\Delta)(\eta_g \otimes \omega_{\mathcal{F}})$  in [BCF])
- relate this to values of  $p$ -adic  $L$ -function

# The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

# The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

$$\eta_{\Pi} = \mathcal{F}^{1-\text{ah}} d\bar{z}_1 dz_2 \in H^1(X_G, \omega_G^{(2-k_1, k_2)})(\Pi)$$

## The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

$$\eta_{\Pi} = \mathcal{F}^{1-\text{ah}} d\bar{z}_1 dz_2 \in H^1(X_G, \omega_G^{(2-k_1, k_2)})[\Pi]$$

If  $\phi$  is a (nearly holomorphic) automorphic form for  $\text{GL}_2$  of weight  $k_1 - k_2$ ,

## The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

$$\eta_\Pi = \mathcal{F}^{1-\text{ah}} d\bar{z}_1 dz_2 \in H^1(X_G, \omega_G^{(2-k_1, k_2)})[\Pi]$$

If  $\phi$  is a (nearly holomorphic) automorphic form for  $\text{GL}_2$  of weight  $k_1 - k_2$ , let

$$I(\phi) = \int_{[\text{GL}_2]} \iota^* \mathcal{F}^{1-\text{ah}}(g) \phi(g) dg$$



## The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

$$\eta_{\Pi} = \mathcal{F}^{1-\text{ah}} d\bar{z}_1 dz_2 \in H^1(X_G, \omega_G^{(2-k_1, k_2)})[\Pi]$$

If  $\phi$  is a (nearly holomorphic) automorphic form for  $\text{GL}_2$  of weight  $k_1 - k_2$ , let

$$I(\phi) = \int_{[\text{GL}_2]} \iota^* \mathcal{F}^{1-\text{ah}}(g) \phi(g) dg$$

(Harris)  $\phi = \text{Eis}^{(k_1-k_2)}(s)$  real analytic Eis series,

$$I(\phi) \sim L(\Pi, \text{As}, s), s \text{ critical}$$

## The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

$$\eta_\Pi = \mathcal{F}^{1-\text{ah}} d\bar{z}_1 dz_2 \in H^1(X_G, \omega_G^{(2-k_1, k_2)})[\Pi]$$

If  $\phi$  is a (nearly holomorphic) automorphic form for  $\text{GL}_2$  of weight  $k_1 - k_2$ , let

$$I(\phi) = \int_{[\text{GL}_2]} \iota^* \mathcal{F}^{1-\text{ah}}(g) \phi(g) dg$$

(Harris)  $\phi = \text{Eis}^{(k_1-k_2)}(s)$  real analytic Eis series,

$$I(\phi) \sim L(\Pi, \text{As}, s), s \text{ critical}$$

(Ichino)  $\phi = \delta_{k_3}^{k_1-k_2} f$  cuspform  $f$  of weight  $k_3 < k_1 - k_2$  generating  $\sigma$ ,

$$I(\phi)^2 \sim L(\Pi \otimes \sigma, r, \frac{1}{2})$$

# The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

$$\eta_\Pi = \mathcal{F}^{1-\text{ah}} d\bar{z}_1 dz_2 \in H^1(X_G, \omega_G^{(2-k_1, k_2)})[\Pi]$$

If  $\phi$  is a (nearly holomorphic) automorphic form for  $\text{GL}_2$  of weight  $k_1 - k_2$ , let

$$I(\phi) = \int_{[\text{GL}_2]} \iota^* \mathcal{F}^{1-\text{ah}}(g) \phi(g) dg$$

(Harris)  $\phi = \text{Eis}^{(k_1-k_2)}(s)$  real analytic Eis series,

$$I(\phi) \sim L(\Pi, \text{As}, s), \text{ } s \text{ critical}$$

(Ichino)  $\phi = \delta_{k_3}^{k_1-k_2} f$  cuspform  $f$  of weight  $k_3 < k_1 - k_2$  generating  $\sigma$ ,

$$I(\phi)^2 \sim L(\Pi \otimes \sigma, r, \frac{1}{2})$$

- $I(\phi)$  can be expressed as a pairing in coherent cohomology:  $\langle \iota^* \eta_\Pi, \omega_\phi \rangle$

# The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

$$\eta_\Pi = \mathcal{F}^{1-\text{ah}} d\bar{z}_1 dz_2 \in H^1(X_G, \omega_G^{(2-k_1, k_2)})[\Pi]$$

If  $\phi$  is a (nearly holomorphic) automorphic form for  $\text{GL}_2$  of weight  $k_1 - k_2$ , let

$$I(\phi) = \int_{[\text{GL}_2]} \iota^* \mathcal{F}^{1-\text{ah}}(g) \phi(g) dg$$

(Harris)  $\phi = \text{Eis}^{(k_1-k_2)}(s)$  real analytic Eis series,

$$I(\phi) \sim L(\Pi, \text{As}, s), s \text{ critical}$$

(Ichino)  $\phi = \delta_{k_3}^{k_1-k_2} f$  cuspform  $f$  of weight  $k_3 < k_1 - k_2$  generating  $\sigma$ ,

$$I(\phi)^2 \sim L(\Pi \otimes \sigma, r, \frac{1}{2})$$

- $I(\phi)$  can be expressed as a pairing in coherent cohomology:  $\langle \iota^* \eta_\Pi, \omega_\phi \rangle$
- $p$ -adic variation of  $\langle -, - \rangle$ ,  $\phi$  and  $\eta_\Pi$  to obtain  $p$ -adic  $L$ -function

# The construction in GLZ (A) and GG (B) $F$ -dominant

$\Pi$  cuspidal automorphic rep of  $G$  of weight  $(k_1, k_2)$ ,  $k_1 > k_2$ :

$$\eta_\Pi = \mathcal{F}^{1-\text{ah}} d\bar{z}_1 dz_2 \in H^1(X_G, \omega_G^{(2-k_1, k_2)})[\Pi]$$

If  $\phi$  is a (nearly holomorphic) automorphic form for  $\text{GL}_2$  of weight  $k_1 - k_2$ , let

$$I(\phi) = \int_{[\text{GL}_2]} \iota^* \mathcal{F}^{1-\text{ah}}(g) \phi(g) dg$$

(Harris)  $\phi = \text{Eis}^{(k_1-k_2)}(s)$  real analytic Eis series,

$$I(\phi) \sim L(\Pi, \text{As}, s), \text{ } s \text{ critical}$$

(Ichino)  $\phi = \delta_{k_3}^{k_1-k_2} f$  cuspform  $f$  of weight  $k_3 < k_1 - k_2$  generating  $\sigma$ ,

$$I(\phi)^2 \sim L(\Pi \otimes \sigma, r, \tfrac{1}{2})$$

- $I(\phi)$  can be expressed as a pairing in coherent cohomology:  $\langle \iota^* \eta_\Pi, \omega_\phi \rangle$
- $p$ -adic variation of  $\langle -, - \rangle$ ,  $\phi$  and  $\eta_\Pi$  to obtain  $p$ -adic  $L$ -function  
↪ higher Hida theory (Boxer–Pilloni)

- Rankin–Selberg/triple product and  $\mathbb{Q}$ -dominant twisted triple product:

$$\langle \iota^* \omega_{\Pi}, \eta_{\phi} \rangle, \quad \omega_{\Pi} \in H^0(X_G) \text{ and } \eta_{\phi} \in H^1(X_{\mathrm{GL}_2}) \simeq H^0(X_{\mathrm{GL}_2})^{\vee}$$

- Rankin–Selberg/triple product and  $\mathbb{Q}$ -dominant twisted triple product:

$$\langle \iota^* \omega_\Pi, \eta_\phi \rangle, \quad \omega_\Pi \in H^0(X_G) \text{ and } \eta_\phi \in H^1(X_{\mathrm{GL}_2}) \simeq H^0(X_{\mathrm{GL}_2})^\vee$$

$$\omega_\Pi \leftrightarrow \mathrm{Eis}^{k_1-k_2} f,$$

- Rankin–Selberg/triple product and  $\mathbb{Q}$ -dominant twisted triple product:

$$\langle \iota^* \omega_\Pi, \eta_\phi \rangle, \quad \omega_\Pi \in H^0(X_G) \text{ and } \eta_\phi \in H^1(X_{\mathrm{GL}_2}) \simeq H^0(X_{\mathrm{GL}_2})^\vee$$

$$\omega_\Pi \leftrightarrow \mathrm{Eis}^{k_1-k_2} f, \quad f_1 f_2,$$



- Rankin–Selberg/triple product and  $\mathbb{Q}$ -dominant twisted triple product:

$$\langle \iota^* \omega_\Pi, \eta_\phi \rangle, \quad \omega_\Pi \in H^0(X_G) \text{ and } \eta_\phi \in H^1(X_{\mathrm{GL}_2}) \simeq H^0(X_{\mathrm{GL}_2})^\vee$$

$$\omega_\Pi \leftrightarrow \mathrm{Eis}^{k_1-k_2} f, \quad f_1 f_2, \quad \delta^k \mathcal{F},$$

- Rankin–Selberg/triple product and  $\mathbb{Q}$ -dominant twisted triple product:

$$\langle \iota^* \omega_\Pi, \eta_\phi \rangle, \quad \omega_\Pi \in H^0(X_G) \text{ and } \eta_\phi \in H^1(X_{\mathrm{GL}_2}) \simeq H^0(X_{\mathrm{GL}_2})^\vee$$

$$\omega_\Pi \leftrightarrow \mathrm{Eis}^{k_1-k_2} f, \quad f_1 f_2, \quad \delta^k \mathcal{F}, \quad \eta_\phi \leftrightarrow g(-\bar{\tau})$$

- Rankin–Selberg/triple product and  $\mathbb{Q}$ -dominant twisted triple product:

$$\langle \iota^* \omega_\Pi, \eta_\phi \rangle, \quad \omega_\Pi \in H^0(X_G) \text{ and } \eta_\phi \in H^1(X_{\mathrm{GL}_2}) \simeq H^0(X_{\mathrm{GL}_2})^\vee$$

$$\omega_\Pi \leftrightarrow \mathrm{Eis}^{k_1-k_2} f, \quad f_1 f_2, \quad \delta^k \mathcal{F}, \quad \eta_\phi \leftrightarrow g(-\bar{\tau})$$

- Previous works with this strategy:

“Eis classes ES”:  $\mathrm{GSp}_4 \supset \mathrm{GL}_2 \times \mathrm{GL}_2$  [Loeffler–Pilloni–Skinner–Zerbes]

“cycles ES”:  $U(1, 2n-1) \supset U(1, n-1) \times U(0, n)$  [Graham]

# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1\left(\omega_G^{(2-\kappa_1, \kappa_2)}\right)$

# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})$
- choice of suitable Hecke operators at  $p_1, p_2$  to cut out ordinary part

# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1\left(\omega_G^{(2-\kappa_1, \kappa_2)}\right)$
- choice of suitable Hecke operators at  $p_1, p_2$  to cut out ordinary part  
     $\rightsquigarrow$  classicality

# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})$
- choice of suitable Hecke operators at  $p_1, p_2$  to cut out ordinary part  
 $\rightsquigarrow$  classicality

## Theorem

There exists a free rank-1  $\mathcal{O}(U)$ -submodule  $H_{\underline{\Pi}}$  of

$$H^1\left(R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})^{ord}\right) \otimes \mathcal{O}(U)$$

such that, if  $k_1, k_2 > 2$ ,

$$H_{\underline{\Pi}}[(k_1, k_2)]$$

# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})$
- choice of suitable Hecke operators at  $p_1, p_2$  to cut out ordinary part  
     $\rightsquigarrow$  classicality

## Theorem

There exists a free rank-1  $O(U)$ -submodule  $H_{\underline{\Pi}}$  of

$$H^1\left(R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})^{\text{ord}}\right) \otimes O(U)$$

such that, if  $k_1, k_2 > 2$ ,

$$H_{\underline{\Pi}}[(k_1, k_2)] \simeq H^1\left(X_G, \omega_G^{(2-k_1, k_2)}\right)^{\text{ord}} [\underline{\Pi}[(k_1, k_2)]] .$$



# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})$
- choice of suitable Hecke operators at  $p_1, p_2$  to cut out ordinary part  
 $\rightsquigarrow$  classicality

## Theorem

There exists a free rank-1  $\mathcal{O}(U)$ -submodule  $H_{\square}$  of

$$H^1\left(R\Gamma_c^1\left(\omega_G^{(2-\kappa_1, \kappa_2)}\right)^{\text{ord}}\right) \otimes \mathcal{O}(U)$$

such that, if  $k_1, k_2 > 2$ ,

$$H_{\square}[(k_1, k_2)] \simeq H^1\left(X_G, \omega_G^{(2-k_1, k_2)}\right)^{\text{ord}}[\square[(k_1, k_2)]].$$

- $\eta_{\square}$  basis of  $H_{\square}$

# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})$
- choice of suitable Hecke operators at  $p_1, p_2$  to cut out ordinary part  
 $\rightsquigarrow$  classicality

## Theorem

There exists a free rank-1  $O(U)$ -submodule  $H_{\underline{n}}$  of

$$H^1\left(R\Gamma_c^1\left(\omega_G^{(2-\kappa_1, \kappa_2)}\right)^{\text{ord}}\right) \otimes O(U)$$

such that, if  $k_1, k_2 > 2$ ,

$$H_{\underline{n}}[(k_1, k_2)] \simeq H^1\left(X_G, \omega_G^{(2-k_1, k_2)}\right)^{\text{ord}} [\underline{n}[(k_1, k_2)]].$$

- $\eta_{\underline{n}}$  basis of  $H_{\underline{n}}$
- (A) Katz'  $p$ -adic Eisenstein measure

# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})$
- choice of suitable Hecke operators at  $p_1, p_2$  to cut out ordinary part  
 $\rightsquigarrow$  classicality

## Theorem

There exists a free rank-1  $O(U)$ -submodule  $H_{\underline{n}}$  of

$$H^1\left(R\Gamma_c^1\left(\omega_G^{(2-\kappa_1, \kappa_2)}\right)^{\text{ord}}\right) \otimes O(U)$$

such that, if  $k_1, k_2 > 2$ ,

$$H_{\underline{n}}[(k_1, k_2)] \simeq H^1\left(X_G, \omega_G^{(2-k_1, k_2)}\right)^{\text{ord}} [\underline{n}[(k_1, k_2)]] .$$

- $\eta_{\underline{n}}$  basis of  $H_{\underline{n}}$
- (A) Katz'  $p$ -adic Eisenstein measure  
(B)  $\Delta^{\kappa'} \phi_{\underline{\sigma}}$ ,  $\underline{\sigma}$  family of car of  $\text{GL}_2$ ,  $\phi_{\underline{\sigma}}$  basis of  $H^0(\omega_{\text{GL}_2}^{\kappa})^{\text{ord}} [\underline{\sigma}]$

# Higher Hida theory (after Boxer–Pilloni) and $p$ -adic $L$ -functions

- higher Hida theory complex:  $R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})$
- choice of suitable Hecke operators at  $p_1, p_2$  to cut out ordinary part  
 $\rightsquigarrow$  classicality

## Theorem

There exists a free rank-1  $O(U)$ -submodule  $H_{\underline{\Pi}}$  of

$$H^1\left(R\Gamma_c^1(\omega_G^{(2-\kappa_1, \kappa_2)})^{\text{ord}}\right) \otimes O(U)$$

such that, if  $k_1, k_2 > 2$ ,

$$H_{\underline{\Pi}}[(k_1, k_2)] \simeq H^1\left(X_G, \omega_G^{(2-k_1, k_2)}\right)^{\text{ord}} [\underline{\Pi}[(k_1, k_2)]] .$$

- $\eta_{\underline{\Pi}}$  basis of  $H_{\underline{\Pi}}$
- (A) Katz'  $p$ -adic Eisenstein measure  
(B)  $\Delta^{\kappa'} \phi_{\underline{\sigma}}$ ,  $\underline{\sigma}$  family of car of  $\text{GL}_2$ ,  $\phi_{\underline{\sigma}}$  basis of  $H^0(\omega_{\text{GL}_2}^{\kappa})^{\text{ord}}[\underline{\sigma}]$
- $\iota_p^* \eta_{\underline{\Pi}} \in H^1(\omega_{\text{GL}_2}^{2-(\kappa_1-\kappa_2)})$