

Matrix Kloosterman sums

Joint work with Árpád Tóth, arXiv:2109.00762

Márton Erdélyi

`merdelyi@math.bme.hu`

Budapest University of Technology of Economics

International Seminar on Automorphic Forms

25.01.2022

Classical Kloosterman sums

Let $q = p^f$ be a prime power, \mathbb{F} the field with q elements and $\varphi : \mathbb{F} \rightarrow \mathbb{C}^*$ a nontrivial additive character.

Definition

For $\alpha, \beta \in \mathbb{F}$ let $K(\alpha, \beta) = \sum_{\gamma \in \mathbb{F}^*} \varphi(\alpha\gamma + \beta\gamma^{-1})$.

Classical Kloosterman sums

Let $q = p^f$ be a prime power, \mathbb{F} the field with q elements and $\varphi : \mathbb{F} \rightarrow \mathbb{C}^*$ a nontrivial additive character.

Definition

For $\alpha, \beta \in \mathbb{F}$ let $K(\alpha, \beta) = \sum_{\gamma \in \mathbb{F}^*} \varphi(\alpha\gamma + \beta\gamma^{-1})$.

Basic properties

- ① $K(\alpha, \beta) \in \mathbb{R}$,
- ② $K(\alpha, \beta) = K(\beta, \alpha)$ and
- ③ if $\delta \in \mathbb{F}^*$, then $K(\alpha, \delta\beta) = K(\alpha\delta, \beta)$. Thus $K(\alpha, \delta) = K(\alpha\delta) := K(\alpha\delta, 1)$.

Classical Kloosterman sums

Let $q = p^f$ be a prime power, \mathbb{F} the field with q elements and $\varphi : \mathbb{F} \rightarrow \mathbb{C}^*$ a nontrivial additive character.

Definition

For $\alpha, \beta \in \mathbb{F}$ let $K(\alpha, \beta) = \sum_{\gamma \in \mathbb{F}^*} \varphi(\alpha\gamma + \beta\gamma^{-1})$.

Basic properties

- ① $K(\alpha, \beta) \in \mathbb{R}$,
- ② $K(\alpha, \beta) = K(\beta, \alpha)$ and
- ③ if $\delta \in \mathbb{F}^*$, then $K(\alpha, \delta\beta) = K(\alpha\delta, \beta)$. Thus $K(\alpha, \delta) = K(\alpha\delta) := K(\alpha\delta, 1)$.

Theorem (Weil)

$|K(\alpha)| < 2\sqrt{q}$ for any $\alpha \in \mathbb{F}$.

Classical Kloosterman sums

Let $q = p^f$ be a prime power, \mathbb{F} the field with q elements and $\varphi : \mathbb{F} \rightarrow \mathbb{C}^*$ a nontrivial additive character.

Definition

For $\alpha, \beta \in \mathbb{F}$ let $K(\alpha, \beta) = \sum_{\gamma \in \mathbb{F}^*} \varphi(\alpha\gamma + \beta\gamma^{-1})$.

Basic properties

- ① $K(\alpha, \beta) \in \mathbb{R}$,
- ② $K(\alpha, \beta) = K(\beta, \alpha)$ and
- ③ if $\delta \in \mathbb{F}^*$, then $K(\alpha, \delta\beta) = K(\alpha\delta, \beta)$. Thus
 $K(\alpha, \delta) = K(\alpha\delta) := K(\alpha\delta, 1)$.

Theorem (Weil)

$|K(\alpha)| < 2\sqrt{q}$ for any $\alpha \in \mathbb{F}$.

In fact, if $\alpha \neq 0$ there exist $\lambda \in \mathbb{C}$, $|\lambda| = \sqrt{q}$ such that $K(\alpha) = -\lambda - \bar{\lambda}$.

Classical Kloosterman sums

Let $q = p^f$ be a prime power, \mathbb{F} the field with q elements and $\varphi : \mathbb{F} \rightarrow \mathbb{C}^*$ a nontrivial additive character.

Definition

For $\alpha, \beta \in \mathbb{F}$ let $K(\alpha, \beta) = \sum_{\gamma \in \mathbb{F}^*} \varphi(\alpha\gamma + \beta\gamma^{-1})$.

Basic properties

- ① $K(\alpha, \beta) \in \mathbb{R}$,
- ② $K(\alpha, \beta) = K(\beta, \alpha)$ and
- ③ if $\delta \in \mathbb{F}^*$, then $K(\alpha, \delta\beta) = K(\alpha\delta, \beta)$. Thus
 $K(\alpha, \delta) = K(\alpha\delta) := K(\alpha\delta, 1)$.

Theorem (Weil)

$|K(\alpha)| < 2\sqrt{q}$ for any $\alpha \in \mathbb{F}$.

In fact, if $\alpha \neq 0$ there exist $\lambda \in \mathbb{C}$, $|\lambda| = \sqrt{q}$ such that $K(\alpha) = -\lambda - \bar{\lambda}$.
 $K(0) = -1$ and $K(0, 0) = q - 1$.

GL_n Kloosterman sums

Let $n \in \mathbb{N}$, $GL_n(\mathbb{F})$ the group of $n \times n$ invertible matrices over \mathbb{F} , $\text{tr} : GL_n(\mathbb{F}) \rightarrow \mathbb{F}$ the matrix trace and $\psi = \text{tr} \circ \varphi$.

Definition

For $a, b \in \mathbb{F}^{n \times n}$ let $K_n(a, b) = \sum_{c \in GL_n(\mathbb{F})} \psi(ac + bc^{-1})$.

GL_n Kloosterman sums

Let $n \in \mathbb{N}$, $GL_n(\mathbb{F})$ the group of $n \times n$ invertible matrices over \mathbb{F} , $\text{tr} : GL_n(\mathbb{F}) \rightarrow \mathbb{F}$ the matrix trace and $\psi = \text{tr} \circ \varphi$.

Definition

For $a, b \in \mathbb{F}^{n \times n}$ let $K_n(a, b) = \sum_{c \in GL_n(\mathbb{F})} \psi(ac + bc^{-1})$.

Remarks

- 1 For $n = 1$ this is the classical sum.

GL_n Kloosterman sums

Let $n \in \mathbb{N}$, $GL_n(\mathbb{F})$ the group of $n \times n$ invertible matrices over \mathbb{F} ,
 $\text{tr} : GL_n(\mathbb{F}) \rightarrow \mathbb{F}$ the matrix trace and $\psi = \text{tr} \circ \varphi$.

Definition

For $a, b \in \mathbb{F}^{n \times n}$ let $K_n(a, b) = \sum_{c \in GL_n(\mathbb{F})} \psi(ac + bc^{-1})$.

Remarks

- 1 For $n = 1$ this is the classical sum.
- 2 The mod p^m analogue of this arises at the study of the horocyclic flow on GL_n (cf Lee, Marklof, Int. Math. Research Notices, Vol 2018 (21))

GL_n Kloosterman sums

Let $n \in \mathbb{N}$, $GL_n(\mathbb{F})$ the group of $n \times n$ invertible matrices over \mathbb{F} , $\text{tr} : GL_n(\mathbb{F}) \rightarrow \mathbb{F}$ the matrix trace and $\psi = \text{tr} \circ \varphi$.

Definition

For $a, b \in \mathbb{F}^{n \times n}$ let $K_n(a, b) = \sum_{c \in GL_n(\mathbb{F})} \psi(ac + bc^{-1})$.

Remarks

- 1 For $n = 1$ this is the classical sum.
- 2 The mod p^m analogue of this arises at the study of the horocyclic flow on GL_n (cf Lee, Marklof, Int. Math. Research Notices, Vol 2018 (21))
- 3 If b is invertible, then $K_n(a, b) = K_n(ab) := K_n(ab, e)$.

GL_n Kloosterman sums

Let $n \in \mathbb{N}$, $GL_n(\mathbb{F})$ the group of $n \times n$ invertible matrices over \mathbb{F} , $\text{tr} : GL_n(\mathbb{F}) \rightarrow \mathbb{F}$ the matrix trace and $\psi = \text{tr} \circ \varphi$.

Definition

For $a, b \in \mathbb{F}^{n \times n}$ let $K_n(a, b) = \sum_{c \in GL_n(\mathbb{F})} \psi(ac + bc^{-1})$.

Remarks

- 1 For $n = 1$ this is the classical sum.
- 2 The mod p^m analogue of this arises at the study of the horocyclic flow on GL_n (cf Lee, Marklof, Int. Math. Research Notices, Vol 2018 (21))
- 3 If b is invertible, then $K_n(a, b) = K_n(ab) := K_n(ab, e)$.

Theorem

$$K_n(a) \ll \begin{cases} q^{(3n^2-1)/4}, & \text{if } n \text{ is odd} \\ q^{3n^2/4}, & \text{if } n \text{ is even} \end{cases} \text{ for any } a \in \mathbb{F}^{n \times n}.$$

The outline of the talk

- ① Explicit evaluation of the sum $K_n(a)$ (using elementary methods)
 - the case $n = 2$
 - over a field extension

The outline of the talk

- ① Explicit evaluation of the sum $K_n(a)$ (using elementary methods)
 - the case $n = 2$
 - over a field extension
- ② Bounds (using cohomological arguments)
 - The general picture – by Weil, Grothendieck and Deligne
 - The bound on $K_n(a)$ – bounding the weights
 - Purity type results – an illustration of a theorem of Fouvry and Katz

The case $n = 2$

The sum is invariant under conjugation: for c invertible

$$K_n(a) = K_n(e, a) = K_n(c^{-1}, ca) = K_n(ca, c^{-1}) = K_n(cac^{-1}).$$

The case $n = 2$

The sum is invariant under conjugation: for c invertible

$$K_n(a) = K_n(e, a) = K_n(c^{-1}, ca) = K_n(ca, c^{-1}) = K_n(cac^{-1}).$$

Theorem

Let $\alpha \neq \beta \in \mathbb{F}$, $\alpha \neq 0$ and $\gamma \in \mathbb{F}_{q^2} - \mathbb{F}$.

$$a^2 = 0 \quad \implies \quad K_2(a) = q$$

$$a \sim \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \quad \implies \quad K_2(a) = qK(\alpha)K(\beta)$$

$$a \sim \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix} \quad \implies \quad K_2(a) = q^3 - q^2 + qK(\alpha)^2$$

$$a \sim \begin{pmatrix} \alpha & 1 \\ 0 & \alpha \end{pmatrix} \quad \implies \quad K_2(a) = -q^2 + qK(\alpha)^2$$

The case $n = 2$

The sum is invariant under conjugation: for c invertible

$$K_n(a) = K_n(e, a) = K_n(c^{-1}, ca) = K_n(ca, c^{-1}) = K_n(cac^{-1}).$$

Theorem

Let $\alpha \neq \beta \in \mathbb{F}$, $\alpha \neq 0$ and $\gamma \in \mathbb{F}_{q^2} - \mathbb{F}$.

$$a^2 = 0 \implies K_2(a) = q$$

$$a \sim \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \implies K_2(a) = qK(\alpha)K(\beta)$$

$$a \sim \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix} \implies K_2(a) = q^3 - q^2 + qK(\alpha)^2$$

$$a \sim \begin{pmatrix} \alpha & 1 \\ 0 & \alpha \end{pmatrix} \implies K_2(a) = -q^2 + qK(\alpha)^2$$

$$a \sim_{\mathbb{F}_{q^2}} \begin{pmatrix} \gamma & 0 \\ 0 & \gamma^q \end{pmatrix} \implies K_2(a) = -qK(\gamma),$$

where in the last line the Kloosterman sum is over \mathbb{F}_{q^2} .

Proposition

Assume that $a = \left(\begin{array}{c|c} a_k & b \\ \hline 0 & a_l \end{array} \right)$ with $a_k \in \mathbb{F}^{k \times k}$, $a_l \in \mathbb{F}^{l \times l}$, $b \in \mathbb{F}^{k \times l}$ for some $k, l \in \mathbb{N}$ such that $k + l = n$ and a_k and a_l have no common eigenvalue in $\overline{\mathbb{F}}$. Then $K_n(a) = q^{kl} K_k(a_k) K_l(a_l)$.

Proposition

Assume that $a = \left(\begin{array}{c|c} a_k & b \\ \hline 0 & a_l \end{array} \right)$ with $a_k \in \mathbb{F}^{k \times k}$, $a_l \in \mathbb{F}^{l \times l}$, $b \in \mathbb{F}^{k \times l}$ for some $k, l \in \mathbb{N}$ such that $k + l = n$ and a_k and a_l have no common eigenvalue in $\overline{\mathbb{F}}$. Then $K_n(a) = q^{kl} K_k(a_k) K_l(a_l)$.

Lemma

If a_k and a_l are as above, then the linear endomorphism of $\mathbb{F}^{k \times l}$ given by $v \mapsto va_l - a_k v$ is an isomorphism.

Proposition

Assume that $a = \left(\begin{array}{c|c} a_k & b \\ \hline 0 & a_l \end{array} \right)$ with $a_k \in \mathbb{F}^{k \times k}$, $a_l \in \mathbb{F}^{l \times l}$, $b \in \mathbb{F}^{k \times l}$ for some $k, l \in \mathbb{N}$ such that $k + l = n$ and a_k and a_l have no common eigenvalue in $\overline{\mathbb{F}}$. Then $K_n(a) = q^{kl} K_k(a_k) K_l(a_l)$.

Lemma

If a_k and a_l are as above, then the linear endomorphism of $\mathbb{F}^{k \times l}$ given by $v \mapsto va_l - a_k v$ is an isomorphism.

Proof of the Lemma. It suffices to prove that this map is injective.

Proposition

Assume that $a = \left(\begin{array}{c|c} a_k & b \\ \hline 0 & a_l \end{array} \right)$ with $a_k \in \mathbb{F}^{k \times k}$, $a_l \in \mathbb{F}^{l \times l}$, $b \in \mathbb{F}^{k \times l}$ for some $k, l \in \mathbb{N}$ such that $k + l = n$ and a_k and a_l have no common eigenvalue in $\overline{\mathbb{F}}$. Then $K_n(a) = q^{kl} K_k(a_k) K_l(a_l)$.

Lemma

If a_k and a_l are as above, then the linear endomorphism of $\mathbb{F}^{k \times l}$ given by $v \mapsto va_l - a_k v$ is an isomorphism.

Proof of the Lemma. It suffices to prove that this map is injective.

Assume that $va_l - a_k v = 0$ for some v . Then for any polynomial $p \in \mathbb{F}[t]$ we have $vp(a_l) = p(a_k)v$.

Proposition

Assume that $a = \left(\begin{array}{c|c} a_k & b \\ \hline 0 & a_l \end{array} \right)$ with $a_k \in \mathbb{F}^{k \times k}$, $a_l \in \mathbb{F}^{l \times l}$, $b \in \mathbb{F}^{k \times l}$ for some $k, l \in \mathbb{N}$ such that $k + l = n$ and a_k and a_l have no common eigenvalue in $\overline{\mathbb{F}}$. Then $K_n(a) = q^{kl} K_k(a_k) K_l(a_l)$.

Lemma

If a_k and a_l are as above, then the linear endomorphism of $\mathbb{F}^{k \times l}$ given by $v \mapsto va_l - a_k v$ is an isomorphism.

Proof of the Lemma. It suffices to prove that this map is injective.

Assume that $va_l - a_k v = 0$ for some v . Then for any polynomial $p \in \mathbb{F}[t]$ we have $vp(a_l) = p(a_k)v$.

Let $p_k, p_l \in \mathbb{F}_q[t]$ be the characteristic polynomials of a_k and a_l . The Cayley-Hamilton theorem implies $0 = p_k(a_k)v = vp_k(a_l)$ and $0 = vp_l(a_l)$.

Proposition

Assume that $a = \left(\begin{array}{c|c} a_k & b \\ \hline 0 & a_l \end{array} \right)$ with $a_k \in \mathbb{F}^{k \times k}$, $a_l \in \mathbb{F}^{l \times l}$, $b \in \mathbb{F}^{k \times l}$ for some $k, l \in \mathbb{N}$ such that $k + l = n$ and a_k and a_l have no common eigenvalue in $\overline{\mathbb{F}}$. Then $K_n(a) = q^{kl} K_k(a_k) K_l(a_l)$.

Lemma

If a_k and a_l are as above, then the linear endomorphism of $\mathbb{F}^{k \times l}$ given by $v \mapsto va_l - a_k v$ is an isomorphism.

Proof of the Lemma. It suffices to prove that this map is injective.

Assume that $va_l - a_k v = 0$ for some v . Then for any polynomial $p \in \mathbb{F}[t]$ we have $vp(a_l) = p(a_k)v$.

Let $p_k, p_l \in \mathbb{F}_q[t]$ be the characteristic polynomials of a_k and a_l . The Cayley-Hamilton theorem implies $0 = p_k(a_k)v = vp_k(a_l)$ and $0 = vp_l(a_l)$. By our assumption on the eigenvalues we have that $p_k(t)$ and $p_l(t)$ are relatively prime in $\mathbb{F}[t]$, thus there are polynomials such that $p_k r_k + p_l r_l = 1$ which implies $0 = v$, hence our claim.

Proof of the Proposition. Let $U_{[k,l]} = \left\{ \left(\frac{e_k}{0} \middle| \frac{v}{e_l} \right) \middle| v \in \mathbb{F}^{k \times l} \right\}$. Then

$$\begin{aligned}
 K_n(a) &= \sum_x \psi(ax + x^{-1}) = \\
 &\quad \frac{1}{q^{kl}} \sum_{u \in U_{[k,l]}} \sum_x \psi(a(u^{-1}xu) + (u^{-1}xu)^{-1}) = \\
 &\quad \frac{1}{q^{kl}} \sum_x \sum_{u \in U_{[k,l]}} \psi((uau^{-1})x + x^{-1}).
 \end{aligned}$$

Proof of the Proposition. Let $U_{[k,l]} = \left\{ \left(\begin{array}{c|c} e_k & v \\ 0 & e_l \end{array} \right) \middle| v \in \mathbb{F}^{k \times l} \right\}$. Then

$$\begin{aligned} K_n(a) &= \sum_x \psi(ax + x^{-1}) = \\ &= \frac{1}{q^{kl}} \sum_{u \in U_{[k,l]}} \sum_x \psi(a(u^{-1}xu) + (u^{-1}xu)^{-1}) = \\ &= \frac{1}{q^{kl}} \sum_x \sum_{u \in U_{[k,l]}} \psi((uau^{-1})x + x^{-1}). \end{aligned}$$

Now $uau^{-1} = \left(\begin{array}{c|c} a_k & b + va_l - a_kv \\ 0 & a_l \end{array} \right)$ and so

$$K_n(a) = \frac{1}{q^{kl}} \sum_x \psi(ax + x^{-1}) \left(\sum_{v \in \mathbb{F}^{k \times l}} \psi_k((va_l - a_kv)x') \right),$$

where x' is the $l \times k$ matrix which we get by deleting the first k rows and last l columns of x .

Proof of the Proposition - continued.

By the lemma we have

$$\sum_{v \in \mathbb{F}^{k \times l}} \psi_k((va_l - a_k v)x') = \sum_{v \in \mathbb{F}^{k \times l}} \psi_k(vx') = \begin{cases} 0, & \text{if } x' \neq 0 \\ q^{kl}, & \text{if } x' = 0. \end{cases}$$

Proof of the Proposition - continued.

By the lemma we have

$$\sum_{v \in \mathbb{F}^{k \times l}} \psi_k((va_l - a_k v)x') = \sum_{v \in \mathbb{F}^{k \times l}} \psi_k(vx') = \begin{cases} 0, & \text{if } x' \neq 0 \\ q^{kl}, & \text{if } x' = 0. \end{cases}$$

This yields

$$K_n(a) = \sum_{\substack{x \in \text{GL}_n(\mathbb{F}) \\ x' = 0}} \psi(ax + x^{-1}) = q^{kl} K_k(a_k) K_l(a_l). \quad \square$$

Proof of the Proposition - continued.

By the lemma we have

$$\sum_{v \in \mathbb{F}^{k \times l}} \psi_k((va_l - a_k v)x') = \sum_{v \in \mathbb{F}^{k \times l}} \psi_k(vx') = \begin{cases} 0, & \text{if } x' \neq 0 \\ q^{kl}, & \text{if } x' = 0. \end{cases}$$

This yields

$$K_n(a) = \sum_{\substack{x \in \text{GL}_n(\mathbb{F}) \\ x' = 0}} \psi(ax + x^{-1}) = q^{kl} K_k(a_k) K_l(a_l). \quad \square$$

Corollary

Taking a field extension that contains the eigenvalues of a allows us to reduce the sum to such matrices which have a unique eigenvalue and are in Jordan normal form.

Proof of the Proposition - continued.

By the lemma we have

$$\sum_{v \in \mathbb{F}^{k \times l}} \psi_k((va_l - a_k v)x') = \sum_{v \in \mathbb{F}^{k \times l}} \psi_k(vx') = \begin{cases} 0, & \text{if } x' \neq 0 \\ q^{kl}, & \text{if } x' = 0. \end{cases}$$

This yields

$$K_n(a) = \sum_{\substack{x \in \text{GL}_n(\mathbb{F}) \\ x' = 0}} \psi(ax + x^{-1}) = q^{kl} K_k(a_k) K_l(a_l). \quad \square$$

Corollary

Taking a field extension that contains the eigenvalues of a allows us to reduce the sum to such matrices which have a unique eigenvalue and are in Jordan normal form.

Denote $a \in JN_n(\alpha)$ if $a \in \mathbb{F}^{n \times n}$ is Jordan normal form with unique eigenvalue $\alpha \in \mathbb{F}$.

- 1 If $a \in JN_n(0)$, then $K_n(a) = (-1)^n q^{n(n-1)/2}$.

- 1 If $a \in JN_n(0)$, then $K_n(a) = (-1)^n q^{n(n-1)/2}$.
- 2 Let $\alpha \in \mathbb{F}^*$ and $e_k \in \mathbb{F}^{k \times k}$ the identity matrix. Then $K_n(\alpha e_n) = q^{n-1} K(\alpha) K_{n-1}(\alpha e_{n-1}) + q^{2n-2} (q^{n-1} - 1) K_{n-2}(\alpha e_{n-2})$. It shows that our bound for K_n is sharp.

- 1 If $a \in JN_n(0)$, then $K_n(a) = (-1)^n q^{n(n-1)/2}$.
- 2 Let $\alpha \in \mathbb{F}^*$ and $e_k \in \mathbb{F}^{k \times k}$ the identity matrix. Then $K_n(\alpha e_n) = q^{n-1} K(\alpha) K_{n-1}(\alpha e_{n-1}) + q^{2n-2} (q^{n-1} - 1) K_{n-2}(\alpha e_{n-2})$. It shows that our bound for K_n is sharp.
- 3 Let $\alpha \in \mathbb{F}^*$ and $a \in JN_n(\alpha)$. There is an explicit recursion formula for $K_n(a)$ depending on q , $K_{n-1}(a')$, $K_{n-2}(a'')$, $K_{n-2}(a''')$ and d , where $a' \in JN_{n-1}(\alpha)$, $a'', a''' \in JN_{n-2}(\alpha)$ and $d = \dim(\text{Ker}(a - \alpha e_n))$.

- 1 If $a \in JN_n(0)$, then $K_n(a) = (-1)^n q^{n(n-1)/2}$.
- 2 Let $\alpha \in \mathbb{F}^*$ and $e_k \in \mathbb{F}^{k \times k}$ the identity matrix. Then $K_n(\alpha e_n) = q^{n-1} K(\alpha) K_{n-1}(\alpha e_{n-1}) + q^{2n-2} (q^{n-1} - 1) K_{n-2}(\alpha e_{n-2})$. It shows that our bound for K_n is sharp.
- 3 Let $\alpha \in \mathbb{F}^*$ and $a \in JN_n(\alpha)$. There is an explicit recursion formula for $K_n(a)$ depending on q , $K_{n-1}(a')$, $K_{n-2}(a'')$, $K_{n-2}(a''')$ and d , where $a' \in JN_{n-1}(\alpha)$, $a'', a''' \in JN_{n-2}(\alpha)$ and $d = \dim(\text{Ker}(a - \alpha e_n))$.

With the insight of Will Sawin:

Let $K(\alpha) = -\lambda - \bar{\lambda}$ with $|\lambda| = \sqrt{q}$ and a as above, then

$$K_n(a) = q^{n(n-1)/2} \sum_{d=0}^n \#(V \leq \mathbb{F}^n \mid \dim(V) = d, aV \subseteq V) \lambda^d \bar{\lambda}^{n-d}.$$

The general picture – by Weil, Grothendieck and Deligne

Let $\mathbb{F}_m \subset \bar{\mathbb{F}}$ the degree m extension of \mathbb{F} – the field with q^m elements. Let $\text{Tr}_m = \text{Tr}_{\mathbb{F}_m|\mathbb{F}}$ the field trace map and $\varphi_m = \text{Tr}_m \circ \varphi$.

The general picture – by Weil, Grothendieck and Deligne

Let $\mathbb{F}_m \subset \bar{\mathbb{F}}$ the degree m extension of \mathbb{F} – the field with q^m elements. Let $\text{Tr}_m = \text{Tr}_{\mathbb{F}_m|\mathbb{F}}$ the field trace map and $\varphi_m = \text{Tr}_m \circ \varphi$.

Let \mathbb{A}^1 be the affine line and X a quasiprojective variety and $f : X \rightarrow \mathbb{A}^1$ be a regular morphism. Then by the Grothendieck trace formula

$$\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = \sum_{i=0}^{2 \dim(X)} (-1)^i \sum_{j=1}^{d_i} (\lambda_j^i)^m,$$

where $X(\mathbb{F}_m)$ is the set of \mathbb{F}_m -points of X and the $\lambda_j^i \in \mathbb{C}$ are the eigenvalues of the geometric Frobenius acting on certain cohomology complex.

The general picture – by Weil, Grothendieck and Deligne

Let $\mathbb{F}_m \subset \bar{\mathbb{F}}$ the degree m extension of \mathbb{F} – the field with q^m elements. Let $\text{Tr}_m = \text{Tr}_{\mathbb{F}_m|\mathbb{F}}$ the field trace map and $\varphi_m = \text{Tr}_m \circ \varphi$.

Let \mathbb{A}^1 be the affine line and X a quasiprojective variety and $f : X \rightarrow \mathbb{A}^1$ be a regular morphism. Then by the Grothendieck trace formula

$$\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = \sum_{i=0}^{2 \dim(X)} (-1)^i \sum_{j=1}^{d_i} (\lambda_j^i)^m,$$

where $X(\mathbb{F}_m)$ is the set of \mathbb{F}_m -points of X and the $\lambda_j^i \in \mathbb{C}$ are the eigenvalues of the geometric Frobenius acting on certain cohomology complex.

By Deligne's work on the Weil conjectures it is known that the eigenvalues are Weil numbers, pure of weight $w \leq i$, i.e. they are algebraic and $|\iota(\lambda_j^i)| = q^{w/2}$ for all embedding $\iota : \mathbb{Q}(\lambda_j^i) \rightarrow \mathbb{C}$.

Classical Kloosterman sums revisited

Let $X = \mathbb{A}^1 - \{0\}$ and f be the morphism which is given by $\gamma \mapsto \alpha\gamma + \beta\gamma^{-1}$. Then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = K(\alpha, \beta, \mathbb{F}_m)$.

Classical Kloosterman sums revisited

Let $X = \mathbb{A}^1 - \{0\}$ and f be the morphism which is given by $\gamma \mapsto \alpha\gamma + \beta\gamma^{-1}$. Then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = K(\alpha, \beta, \mathbb{F}_m)$.

- If $\alpha, \beta \neq 0$ then $d_0 = d_2 = 0$, $d_1 = 2$, $\lambda_1^1 = \lambda$ and $\lambda_2^1 = \bar{\lambda}$ are pure of weight 1, thus $K(\alpha, \beta, \mathbb{F}_m) = -\lambda^m - \bar{\lambda}^m$.

Classical Kloosterman sums revisited

Let $X = \mathbb{A}^1 - \{0\}$ and f be the morphism which is given by $\gamma \mapsto \alpha\gamma + \beta\gamma^{-1}$. Then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = K(\alpha, \beta, \mathbb{F}_m)$.

- If $\alpha, \beta \neq 0$ then $d_0 = d_2 = 0$, $d_1 = 2$, $\lambda_1^1 = \lambda$ and $\lambda_2^1 = \bar{\lambda}$ are pure of weight 1, thus $K(\alpha, \beta, \mathbb{F}_m) = -\lambda^m - \bar{\lambda}^m$.
- If $\alpha = 0$ and $\beta \neq 0$, then $d_0 = d_2 = 0$, $d_1 = 1$, $\lambda_1^1 = 1$ is pure of weight 0, thus $K(0, \beta, \mathbb{F}_m) = -1^m = -1$.

Classical Kloosterman sums revisited

Let $X = \mathbb{A}^1 - \{0\}$ and f be the morphism which is given by $\gamma \mapsto \alpha\gamma + \beta\gamma^{-1}$. Then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = K(\alpha, \beta, \mathbb{F}_m)$.

- If $\alpha, \beta \neq 0$ then $d_0 = d_2 = 0$, $d_1 = 2$, $\lambda_1^1 = \lambda$ and $\lambda_2^1 = \bar{\lambda}$ are pure of weight 1, thus $K(\alpha, \beta, \mathbb{F}_m) = -\lambda^m - \bar{\lambda}^m$.
- If $\alpha = 0$ and $\beta \neq 0$, then $d_0 = d_2 = 0$, $d_1 = 1$, $\lambda_1^1 = 1$ is pure of weight 0, thus $K(0, \beta, \mathbb{F}_m) = -1^m = -1$.
- If $\alpha = \beta = 0$, then $d_0 = 0$, $d_1 = d_2 = 1$, $\lambda_1^1 = 1$ is pure of weight 0 and $\lambda_1^2 = q$ is pure of weight 2, thus $K(0, 0, \mathbb{F}_m) = q^m - 1^m = q^m - 1$.

Classical Kloosterman sums revisited

Let $X = \mathbb{A}^1 - \{0\}$ and f be the morphism which is given by $\gamma \mapsto \alpha\gamma + \beta\gamma^{-1}$. Then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = K(\alpha, \beta, \mathbb{F}_m)$.

- If $\alpha, \beta \neq 0$ then $d_0 = d_2 = 0$, $d_1 = 2$, $\lambda_1^1 = \lambda$ and $\lambda_2^1 = \bar{\lambda}$ are pure of weight 1, thus $K(\alpha, \beta, \mathbb{F}_m) = -\lambda^m - \bar{\lambda}^m$.
- If $\alpha = 0$ and $\beta \neq 0$, then $d_0 = d_2 = 0$, $d_1 = 1$, $\lambda_1^1 = 1$ is pure of weight 0, thus $K(0, \beta, \mathbb{F}_m) = -1^m = -1$.
- If $\alpha = \beta = 0$, then $d_0 = 0$, $d_1 = d_2 = 1$, $\lambda_1^1 = 1$ is pure of weight 0 and $\lambda_2^2 = q$ is pure of weight 2, thus $K(0, 0, \mathbb{F}_m) = q^m - 1^m = q^m - 1$.

Definition

The sum corresponding to $f : X \rightarrow \mathbb{A}^1$ is pure if $d_i = 0$ for all $i \neq \dim(X)$ and $\lambda_j^{\dim(X)}$ is pure of maximal weight (that is $|\lambda_j^{\dim(X)}| = q^{\dim(X)/2}$) for all $1 \leq j \leq d_{\dim(X)}$.

Classical Kloosterman sums revisited

Let $X = \mathbb{A}^1 - \{0\}$ and f be the morphism which is given by $\gamma \mapsto \alpha\gamma + \beta\gamma^{-1}$. Then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = K(\alpha, \beta, \mathbb{F}_m)$.

- If $\alpha, \beta \neq 0$ then $d_0 = d_2 = 0$, $d_1 = 2$, $\lambda_1^1 = \lambda$ and $\lambda_2^1 = \bar{\lambda}$ are pure of weight 1, thus $K(\alpha, \beta, \mathbb{F}_m) = -\lambda^m - \bar{\lambda}^m$.
- If $\alpha = 0$ and $\beta \neq 0$, then $d_0 = d_2 = 0$, $d_1 = 1$, $\lambda_1^1 = 1$ is pure of weight 0, thus $K(0, \beta, \mathbb{F}_m) = -1^m = -1$.
- If $\alpha = \beta = 0$, then $d_0 = 0$, $d_1 = d_2 = 1$, $\lambda_1^1 = 1$ is pure of weight 0 and $\lambda_2^2 = q$ is pure of weight 2, thus $K(0, 0, \mathbb{F}_m) = q^m - 1^m = q^m - 1$.

Definition

The sum corresponding to $f : X \rightarrow \mathbb{A}^1$ is pure if $d_i = 0$ for all $i \neq \dim(X)$ and $\lambda_j^{\dim(X)}$ is pure of maximal weight (that is $|\lambda_j^{\dim(X)}| = q^{\dim(X)/2}$) for all $1 \leq j \leq d_{\dim(X)}$.

In this case there is square-root cancellation.

Bounding the weights of $K_n(a)$

Problem: Bounds over \mathbb{F}_{km} do not imply similar bounds on \mathbb{F} .

Bounding the weights of $K_n(a)$

Problem: Bounds over \mathbb{F}_{km} do not imply similar bounds on \mathbb{F} .

For example $X = (\mathbb{A}^1 \setminus \{1\}) \sqcup \mathbb{A}^0$ and the regular function $f : X \rightarrow \mathbb{A}^1$ defined by

$$f(x) = \begin{cases} x, & \text{if } x \in \mathbb{A}^1 \setminus \{1\}, \\ 0, & \text{if } x \in \mathbb{A}^0. \end{cases}$$

then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = 1 - \varphi(1)^m$ which equals 0 $\iff p|m$.

Bounding the weights of $K_n(a)$

Problem: Bounds over \mathbb{F}_{km} do not imply similar bounds on \mathbb{F} .

For example $X = (\mathbb{A}^1 \setminus \{1\}) \sqcup \mathbb{A}^0$ and the regular function $f : X \rightarrow \mathbb{A}^1$ defined by

$$f(x) = \begin{cases} x, & \text{if } x \in \mathbb{A}^1 \setminus \{1\}, \\ 0, & \text{if } x \in \mathbb{A}^0. \end{cases}$$

then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = 1 - \varphi(1)^m$ which equals 0 $\iff p|m$.

Theorem

Let $a \in JN_n(\alpha, \mathbb{F}_m)$ with $\alpha \in \mathbb{F}_m^*$.

Bounding the weights of $K_n(a)$

Problem: Bounds over \mathbb{F}_{km} do not imply similar bounds on \mathbb{F} .

For example $X = (\mathbb{A}^1 \setminus \{1\}) \sqcup \mathbb{A}^0$ and the regular function $f : X \rightarrow \mathbb{A}^1$ defined by

$$f(x) = \begin{cases} x, & \text{if } x \in \mathbb{A}^1 \setminus \{1\}, \\ 0, & \text{if } x \in \mathbb{A}^0. \end{cases}$$

then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = 1 - \varphi(1)^m$ which equals 0 $\iff p|m$.

Theorem

Let $a \in JN_n(\alpha, \mathbb{F}_m)$ with $\alpha \in \mathbb{F}_m^$.*

Let $B \subset GL_n$ the Borel subgroup of the upper triangular matrices, $w \in W$ an element of the Weyl group and $C_w = BwB$ the corresponding Bruhat cell.

Bounding the weights of $K_n(a)$

Problem: Bounds over \mathbb{F}_{km} do not imply similar bounds on \mathbb{F} .

For example $X = (\mathbb{A}^1 \setminus \{1\}) \sqcup \mathbb{A}^0$ and the regular function $f : X \rightarrow \mathbb{A}^1$ defined by

$$f(x) = \begin{cases} x, & \text{if } x \in \mathbb{A}^1 \setminus \{1\}, \\ 0, & \text{if } x \in \mathbb{A}^0. \end{cases}$$

then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = 1 - \varphi(1)^m$ which equals 0 $\iff p|m$.

Theorem

Let $a \in JN_n(\alpha, \mathbb{F}_m)$ with $\alpha \in \mathbb{F}_m^$.*

Let $B \subset GL_n$ the Borel subgroup of the upper triangular matrices, $w \in W$ an element of the Weyl group and $C_w = BwB$ the corresponding Bruhat cell. Consider the map $f : C_w \rightarrow \mathbb{A}^1$, $x \mapsto \text{tr}(ax + x^{-1})$.

Bounding the weights of $K_n(a)$

Problem: Bounds over \mathbb{F}_{km} do not imply similar bounds on \mathbb{F} .

For example $X = (\mathbb{A}^1 \setminus \{1\}) \sqcup \mathbb{A}^0$ and the regular function $f : X \rightarrow \mathbb{A}^1$ defined by

$$f(x) = \begin{cases} x, & \text{if } x \in \mathbb{A}^1 \setminus \{1\}, \\ 0, & \text{if } x \in \mathbb{A}^0. \end{cases}$$

then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = 1 - \varphi(1)^m$ which equals 0 $\iff p|m$.

Theorem

Let $a \in JN_n(\alpha, \mathbb{F}_m)$ with $\alpha \in \mathbb{F}_m^$.*

Let $B \subset GL_n$ the Borel subgroup of the upper triangular matrices, $w \in W$ an element of the Weyl group and $C_w = BwB$ the corresponding Bruhat cell. Consider the map $f : C_w \rightarrow \mathbb{A}^1, x \mapsto \text{tr}(ax + x^{-1})$.

Then in the corresponding cohomology complex the terms with index more than $n^2 + 2n(w)$ vanish, where

$$n(w) = |\{(i, j) \mid 1 \leq i < j \leq n, w(j) < w(i) \leq j\}|.$$

Bounding the weights of $K_n(a)$

Problem: Bounds over \mathbb{F}_{km} do not imply similar bounds on \mathbb{F} .

For example $X = (\mathbb{A}^1 \setminus \{1\}) \sqcup \mathbb{A}^0$ and the regular function $f : X \rightarrow \mathbb{A}^1$ defined by

$$f(x) = \begin{cases} x, & \text{if } x \in \mathbb{A}^1 \setminus \{1\}, \\ 0, & \text{if } x \in \mathbb{A}^0. \end{cases}$$

then $\sum_{x \in X(\mathbb{F}_m)} \varphi_m(f(x)) = 1 - \varphi(1)^m$ which equals 0 $\iff p|m$.

Theorem

Let $a \in JN_n(\alpha, \mathbb{F}_m)$ with $\alpha \in \mathbb{F}_m^*$.

Let $B \subset GL_n$ the Borel subgroup of the upper triangular matrices, $w \in W$ an element of the Weyl group and $C_w = BwB$ the corresponding Bruhat cell. Consider the map $f : C_w \rightarrow \mathbb{A}^1$, $x \mapsto \text{tr}(ax + x^{-1})$.

Then in the corresponding cohomology complex the terms with index more than $n^2 + 2n(w)$ vanish, where

$$n(w) = |\{(i, j) \mid 1 \leq i < j \leq n, w(j) < w(i) \leq j\}|.$$

Thus $d_i = 0$ if $i > n^2 + 2n(w)$ and all the weights are at most $n^2 + 2n(w)$.

The general bound

To finish the proof we need the following steps:

- If $a \in JN_n(\alpha)$ then $K_n(a) \ll \begin{cases} q^{(3n^2-1)/4}, & \text{if } n \text{ is odd} \\ q^{3n^2/4}, & \text{if } n \text{ is even} \end{cases}$.

The general bound

To finish the proof we need the following steps:

- If $a \in JN_n(\alpha)$ then $K_n(a) \ll \begin{cases} q^{(3n^2-1)/4}, & \text{if } n \text{ is odd} \\ q^{3n^2/4}, & \text{if } n \text{ is even} \end{cases}$.
- Using the (cohomological version) of the Proposition we have the bound for any a ,

The general bound

To finish the proof we need the following steps:

- If $a \in JN_n(\alpha)$ then $K_n(a) \ll \begin{cases} q^{(3n^2-1)/4}, & \text{if } n \text{ is odd} \\ q^{3n^2/4}, & \text{if } n \text{ is even} \end{cases}$.
- Using the (cohomological version) of the Proposition we have the bound for any a , as the Frobenius eigenvalues are independent from the field extension.

The general bound

To finish the proof we need the following steps:

- If $a \in JN_n(\alpha)$ then $K_n(a) \ll \begin{cases} q^{(3n^2-1)/4}, & \text{if } n \text{ is odd} \\ q^{3n^2/4}, & \text{if } n \text{ is even} \end{cases}$.
- Using the (cohomological version) of the Proposition we have the bound for any a , as the Frobenius eigenvalues are independent from the field extension.

Remarks:

- $K_n(\alpha e) = \begin{cases} 2K(\alpha)q^{(3n^2-3)/4} + O(q^{(3n^2-3)/4-1}), & \text{if } n \text{ is odd} \\ q^{3n^2/4} + O(q^{3n^2/4-1}), & \text{if } n \text{ is even} \end{cases}$

The general bound

To finish the proof we need the following steps:

- If $a \in JN_n(\alpha)$ then $K_n(a) \ll \begin{cases} q^{(3n^2-1)/4}, & \text{if } n \text{ is odd} \\ q^{3n^2/4}, & \text{if } n \text{ is even} \end{cases}$.
- Using the (cohomological version) of the Proposition we have the bound for any a , as the Frobenius eigenvalues are independent from the field extension.

Remarks:

- $K_n(\alpha e) = \begin{cases} 2K(\alpha)q^{(3n^2-3)/4} + O(q^{(3n^2-3)/4-1}), & \text{if } n \text{ is odd} \\ q^{3n^2/4} + O(q^{3n^2/4-1}), & \text{if } n \text{ is even} \end{cases}$
- $|K_n(a)| \leq |K_n(\alpha e)|$ for any $a \in JN_n(\alpha)$.

The theorem of Fouvry and Katz

Theorem

Let $X \subset \mathbb{A}^n$ be a affine variety and $f \in \mathbb{Z}[x_1, x_2, \dots, x_N]$ there exist subvarieties $\mathbb{A}^N \supset V_1 \supset V_2 \supset \dots \supset V_N$, $\dim(V_j) \leq N - j$ such that for all $a \in \mathbb{A}^N - V_j$ we have

$$\left| \sum_{x \in X(\mathbb{F}_m)} \varphi(f(x_1, x_2, \dots, x_N) + a_1 x_1 + \dots + a_N x_N) \right| \ll q^{m(\dim(X) + j - 1)/2}$$

The theorem of Fouvry and Katz

Theorem

Let $X \subset \mathbb{A}^n$ be a affine variety and $f \in \mathbb{Z}[x_1, x_2, \dots, x_N]$ there exist subvarieties $\mathbb{A}^N \supset V_1 \supset V_2 \supset \dots \supset V_N$, $\dim(V_j) \leq N - j$ such that for all $a \in \mathbb{A}^N - V_j$ we have

$$\left| \sum_{x \in X(\mathbb{F}_m)} \varphi(f(x_1, x_2, \dots, x_N) + a_1 x_1 + \dots + a_N x_N) \right| \ll q^{m(\dim(X) + j - 1)/2}$$

The case $j = 1$ means than in the generic case there is square-root cancellation.

The theorem of Fouvry and Katz

Theorem

Let $X \subset \mathbb{A}^n$ be a affine variety and $f \in \mathbb{Z}[x_1, x_2, \dots, x_N]$ there exist subvarieties $\mathbb{A}^N \supset V_1 \supset V_2 \supset \dots \supset V_N$, $\dim(V_j) \leq N - j$ such that for all $a \in \mathbb{A}^N - V_j$ we have

$$\left| \sum_{x \in X(\mathbb{F}_m)} \varphi(f(x_1, x_2, \dots, x_N) + a_1 x_1 + \dots + a_N x_N) \right| \ll q^{m(\dim(X) + j - 1)/2}$$

The case $j = 1$ means than in the generic case there is square-root cancellation.

Example – classical Kloosterman sums:

Let $X = V(xy - 1) \subset \mathbb{A}^2$ and $f(x, y) = 0$. Then the sum in the theorem is $K(a_1, a_2, \mathbb{F}_m)$.

The theorem of Fouvry and Katz

Theorem

Let $X \subset \mathbb{A}^n$ be a affine variety and $f \in \mathbb{Z}[x_1, x_2, \dots, x_N]$ there exist subvarieties $\mathbb{A}^N \supset V_1 \supset V_2 \supset \dots \supset V_N$, $\dim(V_j) \leq N - j$ such that for all $a \in \mathbb{A}^N - V_j$ we have

$$\left| \sum_{x \in X(\mathbb{F}_m)} \varphi(f(x_1, x_2, \dots, x_N) + a_1 x_1 + \dots + a_N x_N) \right| \ll q^{m(\dim(X) + j - 1)/2}$$

The case $j = 1$ means than in the generic case there is square-root cancellation.

Example – classical Kloosterman sums:

Let $X = V(xy - 1) \subset \mathbb{A}^2$ and $f(x, y) = 0$. Then the sum in the theorem is $K(a_1, a_2, \mathbb{F}_m)$. In this case $V_1 = V(x = 0, y = 0) = \{(0, 0)\}$ and $V_2 = \emptyset$ are such subvarieties.

Example – GL_n Kloosterman sums:

- Let $N = n^2 + 1$. Then $\mathrm{GL}_n \subset \mathbb{A}^N = V(\mathbb{Z}[x, y])$ is a closed subvariety $V(\det(x)y = 1)$. Let $f : x \mapsto \mathrm{tr}(x^{-1})$ – this is indeed a polynomial in $y\mathbb{Z}[x]$.

Example – GL_n Kloosterman sums:

- Let $N = n^2 + 1$. Then $\mathrm{GL}_n \subset \mathbb{A}^N = V(\mathbb{Z}[x, y])$ is a closed subvariety $V(\det(x)y = 1)$. Let $f : x \mapsto \mathrm{tr}(x^{-1})$ – this is indeed a polynomial in $y\mathbb{Z}[x]$. If $a_N = 0$, then the sum in the theorem is $K_n(a)$.

Example – GL_n Kloosterman sums:

- Let $N = n^2 + 1$. Then $\mathrm{GL}_n \subset \mathbb{A}^N = V(\mathbb{Z}[x, y])$ is a closed subvariety $V(\det(x)y = 1)$. Let $f : x \mapsto \mathrm{tr}(x^{-1})$ – this is indeed a polynomial in $y\mathbb{Z}[x]$. If $a_N = 0$, then the sum in the theorem is $K_n(a)$.
- We have seen that if a has no eigenvalues with multiplicity larger than one, then the sum is pure, thus $|K_n(a)| \ll q^{n^2/2}$.

Example – GL_n Kloosterman sums:

- Let $N = n^2 + 1$. Then $\mathrm{GL}_n \subset \mathbb{A}^N = V(\mathbb{Z}[x, y])$ is a closed subvariety $V(\det(x)y = 1)$. Let $f : x \mapsto \mathrm{tr}(x^{-1})$ – this is indeed a polynomial in $y\mathbb{Z}[x]$. If $a_N = 0$, then the sum in the theorem is $K_n(a)$.
- We have seen that if a has no eigenvalues with multiplicity larger than one, then the sum is pure, thus $|K_n(a)| \ll q^{n^2/2}$.
So V_1 can be chosen to be $V(\gcd(\det(x - \lambda e), \frac{\partial}{\partial \lambda} \det(x - \lambda e)) = 1)$.

Example – GL_n Kloosterman sums:

- Let $N = n^2 + 1$. Then $GL_n \subset \mathbb{A}^N = V(\mathbb{Z}[x, y])$ is a closed subvariety $V(\det(x)y = 1)$. Let $f : x \mapsto \text{tr}(x^{-1})$ – this is indeed a polynomial in $y\mathbb{Z}[x]$. If $a_N = 0$, then the sum in the theorem is $K_n(a)$.
- We have seen that if a has no eigenvalues with multiplicity larger than one, then the sum is pure, thus $|K_n(a)| \ll q^{n^2/2}$.

So V_1 can be chosen to be $V(\gcd(\det(x - \lambda e), \frac{\partial}{\partial \lambda} \det(x - \lambda e)) = 1)$.

- If the characteristic polynomial factorizes over $\bar{\mathbb{F}}$ as $\prod_{j=1}^r (\alpha_j - t)^{\beta_j}$, then

the weights are at most $n^2 + \sum_{j=1}^r \frac{\beta_j^2}{2} - \sum_{2 \nmid \beta_j} 1$.

Example – GL_n Kloosterman sums:

- Let $N = n^2 + 1$. Then $GL_n \subset \mathbb{A}^N = V(\mathbb{Z}[x, y])$ is a closed subvariety $V(\det(x)y = 1)$. Let $f : x \mapsto \operatorname{tr}(x^{-1})$ – this is indeed a polynomial in $y\mathbb{Z}[x]$. If $a_N = 0$, then the sum in the theorem is $K_n(a)$.
- We have seen that if a has no eigenvalues with multiplicity larger than one, then the sum is pure, thus $|K_n(a)| \ll q^{n^2/2}$.

So V_1 can be chosen to be $V(\gcd(\det(x - \lambda e), \frac{\partial}{\partial \lambda} \det(x - \lambda e)) = 1)$.

- If the characteristic polynomial factorizes over $\bar{\mathbb{F}}$ as $\prod_{j=1}^r (\alpha_j - t)^{\beta_j}$, then

the weights are at most $n^2 + \sum_{j=1}^r \frac{\beta_j^2}{2} - \sum_{2 \nmid \beta_j} 1$.

These conditions can also be formulated as polynomial equations in the variables, which makes possible to get certain V_j -s explicitly.

Thank you for your attention!