



PONTIFICIA  
UNIVERSIDAD  
CATÓLICA DE  
VALPARAÍSO

# There are at most finitely many singular moduli that are $S$ -units

Sebastián Herrero

(joint with Ricardo Menares and Juan Rivera-Letelier)

International Seminar on Automorphic Forms, May 2021

# Notation

$$\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im}(z) > 0\}$$

$$\Gamma = \operatorname{SL}_2(\mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{Z}, ad - bc = 1 \right\}$$

# Notation

$$\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im}(z) > 0\}$$

$$\Gamma = \operatorname{SL}_2(\mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{Z}, ad - bc = 1 \right\}$$

$j : \mathbb{H} \rightarrow \mathbb{C}$  modular function with

$$j(z) = q^{-1} + 744 + \sum_{n=1}^{\infty} c_n q^n, \quad \text{where } q := e^{2\pi iz}.$$

# Notation

$$\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im}(z) > 0\}$$

$$\Gamma = \operatorname{SL}_2(\mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{Z}, ad - bc = 1 \right\}$$

$j : \mathbb{H} \rightarrow \mathbb{C}$  modular function with

$$j(z) = q^{-1} + 744 + \sum_{n=1}^{\infty} c_n q^n, \quad \text{where } q := e^{2\pi iz}.$$

Facts:

- 1  $j$  is a Hauptmodul for  $\Gamma$ .
- 2  $j(z)$  is the  $j$ -invariant of the elliptic curve  $E_z \simeq \mathbb{C}/(\mathbb{Z} + z\mathbb{Z})$ .

# CM points and singular moduli

$z$  in  $\mathbb{H}$  is a CM point if  $\mathbb{Q}(z)$  is quadratic (imaginary) over  $\mathbb{Q}$ .

# CM points and singular moduli

$z$  in  $\mathbb{H}$  is a CM point if  $\mathbb{Q}(z)$  is quadratic (imaginary) over  $\mathbb{Q}$ .

## Theorem (CM theory)

*If  $z$  is CM then  $j(z)$  is an algebraic integer. Moreover,  $\mathbb{Q}(z, j(z))$  is an abelian unramified extension of  $\mathbb{Q}(z)$ .*

# CM points and singular moduli

$z$  in  $\mathbb{H}$  is a CM point if  $\mathbb{Q}(z)$  is quadratic (imaginary) over  $\mathbb{Q}$ .

## Theorem (CM theory)

*If  $z$  is CM then  $j(z)$  is an algebraic integer. Moreover,  $\mathbb{Q}(z, j(z))$  is an abelian unramified extension of  $\mathbb{Q}(z)$ .*

If  $z$  is CM, we call  $j(z)$  a singular modulus (following Kronecker).

# A question of Masser

Are there *only finitely many* singular moduli that are algebraic units?



# A question of Masser

Are there *only finitely many* singular moduli that are algebraic units?

**Motivation:** In

*An effective “Theorem of André” for CM-points on a plane curve* (2013)

Bilu, Masser and Zannier proved that there are no pairs  $(j_1, j_2)$  of singular moduli on  $X_1 \cdot X_2 = 1$ .

# Habegger's theorem

# Habegger's theorem

In

*Singular moduli that are algebraic units* (2015)

Habegger proved the there are at most finitely many singular moduli that are algebraic units.

# Habegger's theorem

In

*Singular moduli that are algebraic units* (2015)

Habegger proved the there are at most finitely many singular moduli that are algebraic units.

Habegger's proof does not give a numerical bound for the number of singular moduli that are algebraic units.

# Habegger's theorem

In

*Singular moduli that are algebraic units* (2015)

Habegger proved there are at most finitely many singular moduli that are algebraic units.

Habegger's proof does not give a numerical bound for the number of singular moduli that are algebraic units.

**Natural question:** Is there any such *singular unit*?

# Refinements

# Refinements

① In

*No singular modulus is a unit* (2018)

Bilu, Habegger and Kühne proved that there are no *singular units*.

① In

*No singular modulus is a unit* (2018)

Bilu, Habegger and Kühne proved that there are no *singular units*.

② Let  $\Phi_m(X, Y)$  denote the  $m$ -th modular polynomial ( $m \geq 1$  integer).  
In

*Singular units and isogenies between CM elliptic curves* (2019)

Y. Li proved that  $\Phi_m(j_1, j_2)$  is never an algebraic unit for  $j_1, j_2$  singular moduli.



# Differences of singular moduli

Habegger's work (2015) implies the following result: given an algebraic integer  $\alpha$  there are at most finitely many singular moduli  $j$  such that  $j - \alpha$  is an algebraic unit.

# Differences of singular moduli

Habegger's work (2015) implies the following result: given an algebraic integer  $\alpha$  there are at most finitely many singular moduli  $j$  such that  $j - \alpha$  is an algebraic unit.

**Example:** If  $\alpha = 1$  then

$$j\left(\frac{1+i\sqrt{3}}{2}\right) - \alpha = 0 - 1 = -1$$

is an algebraic unit.

# Differences of singular moduli

Habegger's work (2015) implies the following result: given an algebraic integer  $\alpha$  there are at most finitely many singular moduli  $j$  such that  $j - \alpha$  is an algebraic unit.

**Example:** If  $\alpha = 1$  then

$$j\left(\frac{1+i\sqrt{3}}{2}\right) - \alpha = 0 - 1 = -1$$

is an algebraic unit.

In the case  $\alpha = j_2$  is a singular modulus we have, by Y. Li's theorem with  $m = 1$ , that  $j_1 - j_2$  is never an algebraic unit.

Fact: Differences of singular moduli are very special.

## More on CM points

For  $z$  CM, define

$$D = \text{Disc}(z) = \text{discriminant of min. poly. of } z \text{ over } \mathbb{Z}.$$

## More on CM points

For  $z$  CM, define

$$D = \text{Disc}(z) = \text{discriminant of min. poly. of } z \text{ over } \mathbb{Z}.$$

Then  $z$  is of the form

$$\frac{-b + \sqrt{D}}{2a}$$

with  $a, b, c \in \mathbb{Z}$  coprime,  $a > 0$ ,  $D = b^2 - 4ac < 0$ .

## More on CM points

For  $z$  CM, define

$$D = \text{Disc}(z) = \text{discriminant of min. poly. of } z \text{ over } \mathbb{Z}.$$

Then  $z$  is of the form

$$\frac{-b + \sqrt{D}}{2a}$$

with  $a, b, c \in \mathbb{Z}$  coprime,  $a > 0$ ,  $D = b^2 - 4ac < 0$ .

$\Gamma$  acts on the set  $\text{CM}_D$  of CM points of discriminant  $D$  and we define

$$\Lambda_D = \Gamma \backslash \text{CM}_D.$$

## More on CM points

For  $z$  CM, define

$$D = \text{Disc}(z) = \text{discriminant of min. poly. of } z \text{ over } \mathbb{Z}.$$

Then  $z$  is of the form

$$\frac{-b + \sqrt{D}}{2a}$$

with  $a, b, c \in \mathbb{Z}$  coprime,  $a > 0$ ,  $D = b^2 - 4ac < 0$ .

$\Gamma$  acts on the set  $\text{CM}_D$  of CM points of discriminant  $D$  and we define

$$\Lambda_D = \Gamma \backslash \text{CM}_D.$$

### Theorem (CM theory)

$\Lambda_D$  is finite of cardinality  $h(D)$  (class number) and  $j(\Lambda_D)$  is a full Galois orbit.

# Norms of differences

Given  $\alpha$  in  $\overline{\mathbb{Q}}$  define

$$\mathrm{Nm}(\alpha) = \prod_{\sigma: \mathbb{Q}(\alpha) \hookrightarrow \overline{\mathbb{Q}}} \sigma(\alpha).$$



# Norms of differences

Given  $\alpha$  in  $\overline{\mathbb{Q}}$  define

$$\mathrm{Nm}(\alpha) = \prod_{\sigma: \mathbb{Q}(\alpha) \hookrightarrow \overline{\mathbb{Q}}} \sigma(\alpha).$$

Then  $j_1 - j_2$  is an algebraic unit if and only if  $\mathrm{Nm}(j_1 - j_2) = \pm 1$ .

# Norms of differences

Given  $\alpha$  in  $\overline{\mathbb{Q}}$  define

$$\mathrm{Nm}(\alpha) = \prod_{\sigma: \mathbb{Q}(\alpha) \hookrightarrow \overline{\mathbb{Q}}} \sigma(\alpha).$$

Then  $j_1 - j_2$  is an algebraic unit if and only if  $\mathrm{Nm}(j_1 - j_2) = \pm 1$ .

In

*On singular moduli* (1985)

Gross and Zagier gave an *arithmetic formula* for  $\mathrm{Nm}(j_1 - j_2)$  under certain hypotheses.

# Norms of differences

Given  $\alpha$  in  $\overline{\mathbb{Q}}$  define

$$\mathrm{Nm}(\alpha) = \prod_{\sigma: \mathbb{Q}(\alpha) \hookrightarrow \overline{\mathbb{Q}}} \sigma(\alpha).$$

Then  $j_1 - j_2$  is an algebraic unit if and only if  $\mathrm{Nm}(j_1 - j_2) = \pm 1$ .

In

*On singular moduli* (1985)

Gross and Zagier gave an *arithmetic formula* for  $\mathrm{Nm}(j_1 - j_2)$  under certain hypotheses.

It is not clear how to use Gross and Zagier's formula (or extensions of it) to prove *directly* that  $j_1 - j_2$  is never an algebraic unit.

# Singular $S$ -units

Fix  $S$  a finite set of prime numbers.

# Singular $S$ -units

Fix  $S$  a finite set of prime numbers.

An algebraic integer is an  $S$ -unit if no primes outside  $S$  divide  $N_{\mathbf{m}}(\alpha)$ .

# Singular $S$ -units

Fix  $S$  a finite set of prime numbers.

An algebraic integer is an  $S$ -unit if no primes outside  $S$  divide  $N_{\mathbf{m}}(\alpha)$ .

Theorem (H–Menares–Rivera-Letelier, 2021)

*There are at most finitely many singular moduli that are  $S$ -units.*

# Singular $S$ -units

Fix  $S$  a finite set of prime numbers.

An algebraic integer is an  $S$ -unit if no primes outside  $S$  divide  $N_{\mathbb{m}}(\alpha)$ .

Theorem (H–Menaes–Rivera–Letelier, 2021)

*There are at most finitely many singular moduli that are  $S$ -units.*

By Bilu, Habegger and Kühne, or by Y. Li, every singular modulus is an  $S$ -unit for some finite set  $S$ .

# Numerics: A. Sutherland's table<sup>1</sup>

$D$	$\prod_{z \in \Lambda_D} j(z)$	$D$	$\prod_{z \in \Lambda_D} j(z)$	$D$	$\prod_{z \in \Lambda_D} j(z)$
-3	0	-32	$2^6 5^6 23^3$	-63	$-3^6 5^{12} 17^3 41^3 47^3$
-4	$2^6 3^3$	-35	$-2^{30} 5^3$	-64	$-2^3 3^6 23^3 47^3$
-7	$-3^3 5^3$	-36	$-2^{12} 3^3 11^3 23^3$	-67	$-2^{15} 3^3 5^3 11^3$
-8	$2^6 5^3$	-39	$3^{15} 17^3 23^3 29^3$	-68	$-2^{24} 5^{12} 17^3 47^3$
-11	$-2^{15}$	-40	$2^{12} 3^6 5^3 29^3$	-71	$-11^9 17^6 23^3 41^3 47^3 53^3$
-12	$2^4 3^3 5^3$	-43	$-2^{18} 3^3 5^3$	-72	$2^{12} 5^6 29^3 53^3$
-15	$-3^6 5^3 11^3$	-44	$2^{12} 11^3 17^3 29^3$	-75	$2^{30} 3^6 5^{11} 11^3$
-16	$2^3 3^3 11^3$	-47	$-5^{15} 11^6 23^3 29^3$	-76	$2^{12} 3^9 41^3 53^3$
-19	$-2^{15} 3^3$	-48	$2^4 3^9 5^6 11^3$	-79	$-3^{15} 17^3 29^3 47^3 53^3 59^3$
-20	$-2^{12} 5^3 11^3$	-51	$2^{33} 3^6$	-80	$2^{12} 5^6 11^3 17^6 59^3$
-23	$-5^9 11^3 17^3$	-52	$-2^{12} 3^6 5^6 23^3$	-83	$-2^{48} 5^9$
-24	$2^{12} 3^6 17^3$	-55	$-3^{12} 5^6 11^3 29^3 41^3$	-84	$-2^{24} 3^{15} 47^3 59^3$
-27	$-2^{15} 3^{15} 5^3$	-56	$2^{24} 11^6 17^3 41^3$	-87	$3^{18} 5^{18} 23^3 53^3 59^3$
-28	$3^3 5^3 17^3$	-59	$-2^{48} 11^3$	-88	$2^{12} 3^6 5^6 17^3 41^3$
-31	$-3^9 11^3 17^3 23^3$	-60	$3^6 5^3 29^3 41^3$	-91	$-2^{30} 3^6 17^3$

<sup>1</sup><https://math.mit.edu/~drew/NormsOfSinguliModuli2000.pdf>



# Question

## Question

It seems like  $j\left(\frac{1+\sqrt{-11}}{2}\right) = -2^{15}$  is the only singular modulus that is an  $S$ -unit for  $S$  a singleton. Is this the case?

## Question

It seems like  $j\left(\frac{1+\sqrt{-11}}{2}\right) = -2^{15}$  is the only singular modulus that is an  $S$ -unit for  $S$  a singleton. Is this the case?

A. Sutherland checked this *conjecture* for discriminants  $D$  in  $] -10^5, -3]$  (private communication).

# Difference of singular moduli

# Difference of singular moduli

Fix  $S$  a finite set of prime numbers.

# Difference of singular moduli

Fix  $S$  a finite set of prime numbers.

Theorem (H–Menaes–Rivera–Letelier, 2021)

*Given a singular modulus  $j_2$ , there are at most finitely many singular moduli  $j_1$  such that  $j_1 - j_2$  is an  $S$ -unit.*

# Difference of singular moduli

Fix  $S$  a finite set of prime numbers.

Theorem (H–Menaes–Rivera–Letelier, 2021)

*Given a singular modulus  $j_2$ , there are at most finitely many singular moduli  $j_1$  such that  $j_1 - j_2$  is an  $S$ -unit.*

We use Habegger's original strategy together with the new ingredient that for every prime number  $p$ , singular moduli are  $p$ -adically disperse.

# Habegger's strategy (for singular units)



## Habegger's strategy (for singular units)

Habegger considered the absolute logarithmic Weil height

$$h(a) = \frac{1}{[K : \mathbb{Q}]} \sum_{v \in M_K} d_v \log \max\{1, |a|_v\}$$

for  $a$  in  $K$  a number field, where

- $M_K$  is the set of places of  $K$ ,
- $|\cdot|_v$  is a representative absolute value extending  $|\cdot|_p$  with  $p$  prime or  $\infty$  (the usual field norms on  $\mathbb{Q}$ ),
- $d_v = [K_v : \mathbb{Q}_p]$ .

## Habegger's strategy (for singular units)

Habegger considered the absolute logarithmic Weil height

$$h(a) = \frac{1}{[K : \mathbb{Q}]} \sum_{v \in M_K} d_v \log \max\{1, |a|_v\}$$

for  $a$  in  $K$  a number field, where

- $M_K$  is the set of places of  $K$ ,
- $|\cdot|_v$  is a representative absolute value extending  $|\cdot|_p$  with  $p$  prime or  $\infty$  (the usual field norms on  $\mathbb{Q}$ ),
- $d_v = [K_v : \mathbb{Q}_p]$ .

**First ingredient:** For  $j$  a singular modulus of discriminant  $D$  we have

$$h(j) \geq A \log |D| + B,$$

with  $A, B$  absolute constants,  $A > 0$ .

This follows from results of Colmez (1989), and Nakkajima and Taguchi (1991).



**Second ingredient:** A density estimate for the number of singular moduli around 0. Given  $\varepsilon > 0$  find  $r > 0$  small such that

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$

This follows from the following equidistribution theorem for CM points.

**Second ingredient:** A density estimate for the number of singular moduli around 0. Given  $\varepsilon > 0$  find  $r > 0$  small such that

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$

This follows from the following equidistribution theorem for CM points.

**Theorem (Duke (1988) + Clozel and Ullmo (2004))**

*When  $D \rightarrow -\infty$  we have*

$$\frac{1}{h(D)} \sum_{z \in \Lambda_D} \delta_z \rightarrow \frac{3}{\pi} \frac{dx dy}{y^2}$$

*weakly on  $\Gamma \backslash \mathbb{H}$ .*

**Second ingredient:** A density estimate for the number of singular moduli around 0. Given  $\varepsilon > 0$  find  $r > 0$  small such that

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$

This follows from the following equidistribution theorem for CM points.

**Theorem (Duke (1988) + Clozel and Ullmo (2004))**

*When  $D \rightarrow -\infty$  we have*

$$\frac{1}{h(D)} \sum_{z \in \Lambda_D} \delta_z \rightarrow \frac{3}{\pi} \frac{dx dy}{y^2}$$

*weakly on  $\Gamma \backslash \mathbb{H}$ .*

This step is not effective.



**Third ingredient:** An estimate for the Archimedean distance between a singular modulus and 0. For  $j$  a nonzero singular modulus of discriminant  $D$  we have

$$-\log |j| \leq c_\infty \log |D|,$$

with  $c_\infty > 0$  absolute constant.



**Third ingredient:** An estimate for the Archimedean distance between a singular modulus and 0. For  $j$  a nonzero singular modulus of discriminant  $D$  we have

$$-\log |j| \leq c_\infty \log |D|,$$

with  $c_\infty > 0$  absolute constant.

In the “ $(j - \alpha)$  version” of Habegger’s theorem ( $\alpha$  algebraic integer) one needs David and Hirata-Kohno’s deep lower bound for linear forms on  $n = 2$  elliptic logarithms (2009).

# Putting everything together

# Putting everything together

If  $j$  is a singular unit of discriminant  $D$ , then

$$\begin{aligned} A \log |D| + B &\leq h(j) \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K} d_v \log \max\{1, |a|_v\} \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty} d_v \log \max\{1, |a|_v\} \\ &= -\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty, |a|_v < 1} d_v \log |a|_v, \end{aligned}$$

by the product formula.

# Putting everything together

If  $j$  is a singular unit of discriminant  $D$ , then

$$\begin{aligned} A \log |D| + B &\leq h(j) \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K} d_v \log \max\{1, |a|_v\} \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty} d_v \log \max\{1, |a|_v\} \\ &= -\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty, |a|_v < 1} d_v \log |a|_v, \end{aligned}$$

by the product formula. For  $\varepsilon > 0$  convenient we get

$$-\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty, |a|_v < 1} d_v \log |a|_v \leq A_\varepsilon \log |D| + B_\varepsilon$$

with  $A_\varepsilon < A$ .

# Putting everything together

If  $j$  is a singular unit of discriminant  $D$ , then

$$\begin{aligned} A \log |D| + B &\leq h(j) \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K} d_v \log \max\{1, |a|_v\} \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty} d_v \log \max\{1, |a|_v\} \\ &= -\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty, |a|_v < 1} d_v \log |a|_v, \end{aligned}$$

by the product formula. For  $\varepsilon > 0$  convenient we get

$$-\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty, |a|_v < 1} d_v \log |a|_v \leq A_\varepsilon \log |D| + B_\varepsilon$$

with  $A_\varepsilon < A$ . Hence  $|D|$  is bounded and the result follows.

# Our proof (for singular $S$ -units)

## Our proof (for singular $S$ -units)

We use Habegger's strategy. For  $p$  prime, fix an extension of  $|\cdot|_p$  to  $\overline{\mathbb{Q}}$ .

**First ingredient:** For  $j$  a singular modulus of discriminant  $D$  we have

$$h(j) \geq A \log |D| + B,$$

with  $A, B$  absolute constants,  $A > 0$ .

## Our proof (for singular $S$ -units)

We use Habegger's strategy. For  $p$  prime, fix an extension of  $|\cdot|_p$  to  $\overline{\mathbb{Q}}$ .

**First ingredient:** For  $j$  a singular modulus of discriminant  $D$  we have

$$h(j) \geq A \log |D| + B,$$

with  $A, B$  absolute constants,  $A > 0$ .

**Second ingredient:** A  $p$ -adic density estimate for the number of singular moduli around 0. Given  $\varepsilon > 0$  find  $r > 0$  small such that

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B_p(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$



## Our proof (for singular $S$ -units)

We use Habegger's strategy. For  $p$  prime, fix an extension of  $|\cdot|_p$  to  $\overline{\mathbb{Q}}$ .

**First ingredient:** For  $j$  a singular modulus of discriminant  $D$  we have

$$h(j) \geq A \log |D| + B,$$

with  $A, B$  absolute constants,  $A > 0$ .

**Second ingredient:** A  $p$ -adic density estimate for the number of singular moduli around 0. Given  $\varepsilon > 0$  find  $r > 0$  small such that

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B_p(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$

**Third ingredient:** An estimate for the  $p$ -adic distance between a singular modulus and 0. For  $j$  a nonzero singular modulus of discriminant  $D$  we have

$$-\log |j|_p \leq c_p \log |D|,$$

with  $c_p > 0$  absolute constant.

# Putting everything together

# Putting everything together

If  $j$  is a singular unit of discriminant  $D$ , then

$$\begin{aligned} A \log |D| + B &\leq h(j) \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K} d_v \log \max\{1, |a|_v\} \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty \cup M_K^S} d_v \log \max\{1, |a|_v\} \\ &= -\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty \cup M_K^S, |a|_v < 1} d_v \log |a|_v, \end{aligned}$$

by the product formula.

# Putting everything together

If  $j$  is a singular unit of discriminant  $D$ , then

$$\begin{aligned} A \log |D| + B &\leq h(j) \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K} d_v \log \max\{1, |a|_v\} \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty \cup M_K^S} d_v \log \max\{1, |a|_v\} \\ &= -\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty \cup M_K^S, |a|_v < 1} d_v \log |a|_v, \end{aligned}$$

by the product formula. For  $\varepsilon > 0$  convenient we get

$$-\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty \cup M_K^S, |a|_v < 1} d_v \log |a|_v \leq A_{\varepsilon, S} \log |D| + B_{\varepsilon, S}$$

with  $A_{\varepsilon, S} < A$ .

# Putting everything together

If  $j$  is a singular unit of discriminant  $D$ , then

$$\begin{aligned} A \log |D| + B &\leq h(j) \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K} d_v \log \max\{1, |a|_v\} \\ &= \frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty \cup M_K^S} d_v \log \max\{1, |a|_v\} \\ &= -\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty \cup M_K^S, |a|_v < 1} d_v \log |a|_v, \end{aligned}$$

by the product formula. For  $\varepsilon > 0$  convenient we get

$$-\frac{1}{[K:\mathbb{Q}]} \sum_{v \in M_K^\infty \cup M_K^S, |a|_v < 1} d_v \log |a|_v \leq A_{\varepsilon, S} \log |D| + B_{\varepsilon, S}$$

with  $A_{\varepsilon, S} < A$ . Hence  $|D|$  is bounded and the result follows.

# Singular moduli are $p$ -adically disperse

# Singular moduli are $p$ -adically disperse

Theorem (H–Menares–Rivera–Letelier, 2021)

*Given  $\varepsilon > 0$  there exists  $r > 0$  small such that*

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B_p(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$

# Singular moduli are $p$ -adically disperse

## Theorem (H–Menares–Rivera–Letelier, 2021)

*Given  $\varepsilon > 0$  there exists  $r > 0$  small such that*

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B_p(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$

This follows from our identification of all limit measures of CM points in the  $p$ -adic setting.



# $p$ -adic distribution of CM points

## $p$ -adic distribution of CM points

For simplicity, restrict to  $D < 0$  fundamental discriminant.

We have  $j(\Lambda_D) \subset \overline{\mathbb{Q}} \subset \mathbb{C}_p \subset \mathbb{A}_{\text{Berk}}^1$ .

# $p$ -adic distribution of CM points

For simplicity, restrict to  $D < 0$  fundamental discriminant.

We have  $j(\Lambda_D) \subset \overline{\mathbb{Q}} \subset \mathbb{C}_p \subset \mathbb{A}_{\text{Berk}}^1$ .

$\mathbb{A}_{\text{Berk}}^1$  can be defined as the set of multiplicative semi-norms on  $\mathbb{C}_p[X]$  extending  $|\cdot|_p$  on  $\mathbb{C}_p$ , with a suitable topology (locally compact, arc-connected),

# $p$ -adic distribution of CM points

For simplicity, restrict to  $D < 0$  fundamental discriminant.

We have  $j(\Lambda_D) \subset \overline{\mathbb{Q}} \subset \mathbb{C}_p \subset \mathbb{A}_{\text{Berk}}^1$ .


$\mathbb{A}_{\text{Berk}}^1$  can be defined as the set of multiplicative semi-norms on  $\mathbb{C}_p[X]$  extending  $|\cdot|_p$  on  $\mathbb{C}_p$ , with a suitable topology (locally compact, arc-connected), and

$$\mathbb{C}_p \hookrightarrow \mathbb{A}_{\text{Berk}}^1, \quad z \mapsto \iota(z)$$

is defined by  $\iota(z)(f) = |f(z)|_p$  for  $f$  in  $\mathbb{C}_p[X]$ .


$\mathbb{C}_p$  is dense in  $\mathbb{A}_{\text{Berk}}^1$ .

Above the unit disc in  $\mathbb{C}_p$  we have the following picture<sup>2</sup>



<sup>2</sup>Illustration of Joe Silverman

Above the unit disc in  $\mathbb{C}_p$  we have the following picture<sup>2</sup>



At the top we have the Gauss point  $\zeta$  defined by

$$\zeta(a_0 + a_1X + \dots + a_nX^n) = \max\{|a_0|_p, |a_1|_p, \dots, |a_n|_p\}.$$

---

<sup>2</sup>Illustration of Joe Silverman

# Convergence towards the Gauss point

# Convergence towards the Gauss point

## Theorem (H–Menares–Rivera–Letelier, 2020)

- ① For fundamental discriminants  $D < 0$  with  $\left(\frac{D}{p}\right) = 1$  we have

$$\frac{1}{h(D)} \sum_{z \in \Lambda_D} \delta_{j(z)} \rightarrow \delta_\zeta$$

weakly on  $\mathbb{A}_{\text{Berk}}^1$ .

- ② This is not the case for fundamental discriminants  $D < 0$  with  $\left(\frac{D}{p}\right) \neq 1$ .



The case  $\left(\frac{D}{p}\right) \neq 1$

The case  $\left(\frac{D}{p}\right) \neq 1$

Let  $\mathcal{O}_D$  denote the ring of integers of  $\mathbb{Q}(\sqrt{D})$ . Then

$$\Lambda_D = \{E \text{ ell. curve over } \overline{\mathbb{Q}} \text{ with } \text{End}(E) \simeq \mathcal{O}_D\} \subset Y(\overline{\mathbb{Q}})$$

where  $Y(\overline{\mathbb{Q}})$  is the (open) moduli space of elliptic curves over  $\overline{\mathbb{Q}}$ .

The case  $\left(\frac{D}{p}\right) \neq 1$

Let  $\mathcal{O}_D$  denote the ring of integers of  $\mathbb{Q}(\sqrt{D})$ . Then

$$\Lambda_D = \{E \text{ ell. curve over } \overline{\mathbb{Q}} \text{ with } \text{End}(E) \simeq \mathcal{O}_D\} \subset Y(\overline{\mathbb{Q}})$$

where  $Y(\overline{\mathbb{Q}})$  is the (open) moduli space of elliptic curves over  $\overline{\mathbb{Q}}$ .

Let  $\mathfrak{D}$  be the  $p$ -adic discriminant of the ring of integers  $\mathcal{O}_{\mathfrak{D}}$  of  $\mathbb{Q}_p(\sqrt{D})$ .

The case  $\left(\frac{D}{p}\right) \neq 1$

Let  $\mathcal{O}_D$  denote the ring of integers of  $\mathbb{Q}(\sqrt{D})$ . Then

$$\Lambda_D = \{E \text{ ell. curve over } \overline{\mathbb{Q}} \text{ with } \text{End}(E) \simeq \mathcal{O}_D\} \subset Y(\overline{\mathbb{Q}})$$

where  $Y(\overline{\mathbb{Q}})$  is the (open) moduli space of elliptic curves over  $\overline{\mathbb{Q}}$ .

Let  $\mathfrak{D}$  be the  $p$ -adic discriminant of the ring of integers  $\mathcal{O}_{\mathfrak{D}}$  of  $\mathbb{Q}_p(\sqrt{D})$ . Then  $D \in \mathfrak{D}$ ,  $\mathcal{O}_D \subset \mathcal{O}_{\mathfrak{D}}$  and

$$\Lambda_D \subset \Lambda_{\mathfrak{D}} = \{E \text{ ell. curve over } \overline{\mathbb{Q}}_p \text{ with } \text{End}(\hat{E}) \simeq \mathcal{O}_{\mathfrak{D}}\} \subset Y(\overline{\mathbb{Q}}_p)$$

where  $\hat{E}$  is the *formal group* of  $E$ .

The case  $\left(\frac{D}{p}\right) \neq 1$

The case  $\left(\frac{D}{p}\right) \neq 1$

Every (fundamental) discriminant  $D < 0$  with  $\left(\frac{D}{p}\right) \neq 1$  belongs to some  $p$ -adic (fundamental) discriminant  $\mathfrak{D}$ .

The case  $\left(\frac{D}{p}\right) \neq 1$

Every (fundamental) discriminant  $D < 0$  with  $\left(\frac{D}{p}\right) \neq 1$  belongs to some  $p$ -adic (fundamental) discriminant  $\mathfrak{D}$ .

**Theorem (H–Menaes–Rivera-Letelier, 2021)**

*For a  $p$ -adic discriminant  $\mathfrak{D}$  the set  $\Lambda_{\mathfrak{D}}$  is compact and there exists a (unique) Borel probability measure  $\nu_{\mathfrak{D}}$  with support  $\Lambda_{\mathfrak{D}}$  such that for fundamental discriminants  $D < 0$  with  $D \in \mathfrak{D}$  we have*

$$\frac{1}{h(D)} \sum_{z \in \Lambda_D} \delta_{j(z)} \rightarrow \nu_{\mathfrak{D}}$$

*weakly on  $Y(\overline{\mathbb{Q}_p})$ .*

The case  $\left(\frac{D}{p}\right) \neq 1$

Every (fundamental) discriminant  $D < 0$  with  $\left(\frac{D}{p}\right) \neq 1$  belongs to some  $p$ -adic (fundamental) discriminant  $\mathfrak{D}$ .

**Theorem (H–Menaes–Rivera-Letelier, 2021)**

*For a  $p$ -adic discriminant  $\mathfrak{D}$  the set  $\Lambda_{\mathfrak{D}}$  is compact and there exists a (unique) Borel probability measure  $\nu_{\mathfrak{D}}$  with support  $\Lambda_{\mathfrak{D}}$  such that for fundamental discriminants  $D < 0$  with  $D \in \mathfrak{D}$  we have*

$$\frac{1}{h(D)} \sum_{z \in \Lambda_D} \delta_{j(z)} \rightarrow \nu_{\mathfrak{D}}$$

*weakly on  $Y(\overline{\mathbb{Q}_p})$ .*

There are 3 (for  $p > 2$ ) or 7 (for  $p = 2$ )  $p$ -adic fundamental discriminants.



# Singular moduli are $p$ -adically disperse

# Singular moduli are $p$ -adically disperse

Theorem (H–Menaes–Rivera–Letelier, 2021)

*None of the limit measures of CM points in the  $p$ -adic topology has an atom in  $\mathbb{C}_p$ .*

# Singular moduli are $p$ -adically disperse

Theorem (H–Menaes–Rivera–Letelier, 2021)

*None of the limit measures of CM points in the  $p$ -adic topology has an atom in  $\mathbb{C}_p$ .*

This implies our second ingredient.

# Singular moduli are $p$ -adically disperse

## Theorem (H–Menaes–Rivera–Letelier, 2021)

*None of the limit measures of CM points in the  $p$ -adic topology has an atom in  $\mathbb{C}_p$ .*

This implies our second ingredient.

## Theorem (H–Menaes–Rivera–Letelier, 2021)

*Given  $\varepsilon > 0$  there exists  $r > 0$  small such that*

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B_p(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$

# Singular moduli are $p$ -adically disperse

## Theorem (H–Menaes–Rivera–Letelier, 2021)

*None of the limit measures of CM points in the  $p$ -adic topology has an atom in  $\mathbb{C}_p$ .*

This implies our second ingredient.

## Theorem (H–Menaes–Rivera–Letelier, 2021)

*Given  $\varepsilon > 0$  there exists  $r > 0$  small such that*

$$\frac{1}{h(D)} (j(\Lambda_D) \cap B_p(0, r)) \leq \varepsilon \text{ for } D \rightarrow -\infty.$$

# The last ingredient

# The last ingredient

## Theorem (H–Menares–Rivera–Letelier, 2021)

*For  $j$  a nonzero singular modulus of discriminant  $D$  we have*

$$-\log |j|_p \leq c_p \log |D|,$$

*with  $c_p > 0$  absolute constant.*

# The last ingredient

## Theorem (H–Menaes–Rivera–Letelier, 2021)

*For  $j$  a nonzero singular modulus of discriminant  $D$  we have*

$$-\log |j|_p \leq c_p \log |D|,$$

*with  $c_p > 0$  absolute constant.*

The proof uses ideas of F. Charles (2018) and results of Gross (1986): deformation theory of elliptic curves, formal groups/modules, and canonical/quasi-canonical liftings.



# The last ingredient

## Theorem (H–Menares–Rivera–Letelier, 2021)

*For  $j$  a nonzero singular modulus of discriminant  $D$  we have*

$$-\log |j|_p \leq c_p \log |D|,$$

*with  $c_p > 0$  absolute constant.*

The proof uses ideas of F. Charles (2018) and results of Gross (1986): deformation theory of elliptic curves, formal groups/modules, and canonical/quasi-canonical liftings.

With the three main ingredients, we get the result!

# The last ingredient

## Theorem (H–Menares–Rivera–Letelier, 2021)

*For  $j$  a nonzero singular modulus of discriminant  $D$  we have*

$$-\log |j|_p \leq c_p \log |D|,$$

*with  $c_p > 0$  absolute constant.*

The proof uses ideas of F. Charles (2018) and results of Gross (1986): deformation theory of elliptic curves, formal groups/modules, and canonical/quasi-canonical liftings.

With the three main ingredients, we get the result!

The strategy is essentially the same for differences of singular moduli that are  $S$ -units.

# Final comments

# Final comments

In

*On singular moduli that are  $S$ -units (2020)*

F. Campagna shows that  $S_0 = \{p \text{ prime}, p \equiv 1 \pmod{3}\}$  every singular  $S$ -unit is a singular unit, hence there are none.

# Final comments

In

*On singular moduli that are  $S$ -units (2020)*

F. Campagna shows that  $S_0 = \{p \text{ prime}, p \equiv 1 \pmod{3}\}$  every singular  $S$ -unit is a singular unit, hence there are none.

We can use Campagna's result to extend ours to certain classes of infinite sets  $S$  of prime numbers (larger than  $S_0$ ).

# Other modular functions

---

<sup>3</sup>(private communication)

## Other modular functions

Habegger asked us<sup>3</sup>: What about the  $\lambda$ -invariants? These are Hauptmoduln for  $\Gamma(2)$ .

---

<sup>3</sup>(private communication)

## Other modular functions

Habegger asked us<sup>3</sup>: What about the  $\lambda$ -invariants? These are Hauptmoduln for  $\Gamma(2)$ .

**General question:** What about more general Hauptmoduln?

---

<sup>3</sup>(private communication)



## Other modular functions

Habegger asked us<sup>3</sup>: What about the  $\lambda$ -invariants? These are Hauptmoduln for  $\Gamma(2)$ .

**General question:** What about more general Hauptmoduln?

The method seems to extend without major difficulties to the case of differences of singular moduli that are  $S$ -units for any Hauptmodul of a genus zero subgroup of  $\mathrm{GL}_2^+(\mathbb{Q})$  that is algebraically related to the  $j$ -function.

---

<sup>3</sup>(private communication)

# Examples

# Examples

- ① The  $\lambda$ -invariants: there are six of them, they satisfy

$$2^8(1 - \lambda + \lambda^2)^3 - j\lambda^2(1 - \lambda)^2 = 0.$$

# Examples

- ① The  $\lambda$ -invariants: there are six of them, they satisfy

$$2^8(1 - \lambda + \lambda^2)^3 - j\lambda^2(1 - \lambda)^2 = 0.$$

In

*The lambda invariant at CM points (2018)*

Yang, Yin and Yu chose a particular lambda invariant  $\lambda_0$  and proved that  $\lambda_0(z)$  is an algebraic unit for infinitely many CM points  $z$ .

# Examples

- ① The  $\lambda$ -invariants: there are six of them, they satisfy

$$2^8(1 - \lambda + \lambda^2)^3 - j\lambda^2(1 - \lambda)^2 = 0.$$

In

*The lambda invariant at CM points (2018)*

Yang, Yin and Yu chose a particular lambda invariant  $\lambda_0$  and proved that  $\lambda_0(z)$  is an algebraic unit for infinitely many CM points  $z$ .

- ② Weber functions  $\mathbf{f}, \mathbf{f}_1, \mathbf{f}_2$  are roots of

$$(X^{24} \pm 16)^3 - X^{24}j = 0.$$

# Examples

- ① The  $\lambda$ -invariants: there are six of them, they satisfy

$$2^8(1 - \lambda + \lambda^2)^3 - j\lambda^2(1 - \lambda)^2 = 0.$$

In

*The lambda invariant at CM points (2018)*

Yang, Yin and Yu chose a particular lambda invariant  $\lambda_0$  and proved that  $\lambda_0(z)$  is an algebraic unit for infinitely many CM points  $z$ .

- ② Weber functions  $\mathbf{f}, \mathbf{f}_1, \mathbf{f}_2$  are roots of

$$(X^{24} \pm 16)^3 - X^{24}j = 0.$$

Weber's computations show that infinitely many singular moduli for these functions are algebraic units.

# Examples

- ① The  $\lambda$ -invariants: there are six of them, they satisfy

$$2^8(1 - \lambda + \lambda^2)^3 - j\lambda^2(1 - \lambda)^2 = 0.$$

In

*The lambda invariant at CM points (2018)*

Yang, Yin and Yu chose a particular lambda invariant  $\lambda_0$  and proved that  $\lambda_0(z)$  is an algebraic unit for infinitely many CM points  $z$ .

- ② Weber functions  $\mathbf{f}, \mathbf{f}_1, \mathbf{f}_2$  are roots of

$$(X^{24} \pm 16)^3 - X^{24}j = 0.$$

Weber's computations show that infinitely many singular moduli for these functions are algebraic units.

**Note that 0 is not a singular modulus for any of these functions.**

Fin

¡Muchas gracias!

