


[Problems](#)
[Submit](#)
[Runs Status](#)
[Rank List](#)
[Statistics](#)
[Clarifications](#)

## B. A Simple Problem

Time Limit: 1.0 Seconds Memory Limit: 65536K

Given a nonnegative integer  $a$ , and a positive integer  $N$ , we define:

$$f(a, 1) = a$$

$$f(a, k) = f(a, k-1) * f(a, k-1) \% N, k > 1$$

There may or may not exist some positive integer  $k$  satisfying  $f(a, k) = 0$ .

Your task is, given a positive integer  $N$ , to determine how many  $a$  ( $0 \leq a \leq N$ ) there are, such that for some positive integer  $k$ ,  $f(a, k) = 0$ .

### Input

The input contains an integer  $T$  on the first line, indicating the number of test cases. Each test case contains only one positive integer  $N$  ( $1 \leq N \leq 1000000000$ ) on a line.

### Output

For each test case, output the answer on a single line.

### Sample Input

```
6
2
12
50
180
245
361
```

### Sample Output

```
2
3
6
7
8
20
```

*Source: The 5th UESTC Programming Contest*

[Problem ID in problemset: 2793](#)

[Submit](#) [Back](#) [Runs](#) [Statistics](#) [Clarifications](#)

[Tianjin University Online Judge v1.2.4](#)