

# Lab Exercise #1

Lysyl Bess H. Talaban

2025-09-29

*#1a.*

```
vector1 <- c (-5:5)
vector1
```

```
## [1] -5 -4 -3 -2 -1 0 1 2 3 4 5
```

*#the output are numbers from -5 to 5*

*#1b.*

```
x <- 1:7
x
```

```
## [1] 1 2 3 4 5 6 7
```

*#the output are numbers from 1 to 7*

*#2*

```
seq1 <- seq(1, 3, by=0.2)
seq1
```

```
## [1] 1.0 1.2 1.4 1.6 1.8 2.0 2.2 2.4 2.6 2.8 3.0
```

*#the output is a sequence of numbers from 1 to 3 in steps of 0.2*

*#3*

```
ages <- c(34, 28, 22, 36, 27, 18, 52, 39, 42, 29, 35, 31, 27,
         22, 37, 34, 19, 20, 57, 49, 50, 37, 46, 25, 17, 37, 43, 53, 41, 51, 35,
         24, 33, 41, 53, 40, 18, 44, 38, 41, 48, 27, 39, 19, 30, 61, 54, 58, 26,
         18)
```

*#a. Access 3rd element, what is the value?*

```
ages[3]
```

```
## [1] 22
```

*#the value is 22*

*#b. Access 2nd and 4th element, what are the values?*

```
ages[c(2,4)]
```

```
## [1] 28 36
```

*#the values are 28 and 36*

*#c. Access all but the 1st element is not included. Write the R code and its output.*

```
ages[c(2:50)]
```

```
## [1] 28 22 36 27 18 52 39 42 29 35 31 27 22 37 34 19 20 57 49 50 37 46 25 17 37
## [26] 43 53 41 51 35 24 33 41 53 40 18 44 38 41 48 27 39 19 30 61 54 58 26 18
```

```
# OUTPUT
```

```
#> # 3b.
```

```
# > ages[c(2, 4)]
```

```
#[1] 28 36
```

```
#> # the values are 28 and 36
```

```
# >
```

```
# > # 3c.
```

```
# > ages[c(2:50)]
```

```
#[1] 28 22 36 27 18 52 39 42 29 35 31 27 22 37 34 19 20 57 49 50 37 46 25 17 37 43 53 41 51 35 24 33 41
```

```
#[39] 41 48 27 39 19 30 61 54 58 26 18
```

```
# 4.
```

```
x <- c("first"=3, "second"=0, "third"=9)
```

```
x
```

```
## first second third
```

```
##      3      0      9
```

```
# a. Print the results. Then access x[c("first", "third")]. Describe the output.
```

```
x[c("first", "third")]
```

```
## first third
```

```
##      3      9
```

```
# the output is displays a 2x2 table with the corresponding columns and rows
```

```
# b. Write the code and its output.
```

```
# OUTPUT
```

```
#x <- c("first"=3, "second"=0, "third"=9)
```

```
#> x
```

```
#first second third
```

```
#3      0      9
```

```
#>
```

```
# > # 4a.
```

```
# > x[c("first", "third")]
```

```
#first third
```

```
#3      9
```

```
#> # the output is displays a 2x2 table with the corresponding columns and rows
```

```
# >
```

```
# > # 4b.
```

```
# > x
```

```
#first second third
```

```
#3      0      9
```

```
# 5. Create a sequence x from -3:2.
```

```
x <- seq(-3:2)
```

```
# a. Modify 2nd element and change it to 0; Describe the output.
```

```
x[2] <- 0
```

```
x
```

```
## [1] 1 0 3 4 5 6
```

```
# 5b. Write the code and its output.
```

```
# OUTPUT
```

```
#> x[2] <- 0
#> x
#[1] 1 0 3 4 5 6
```

```
# 6.
```

```
Month <- c("Jan", "Feb", "March", "Apr", "May", "June")
Price_per_liter_PhP <- c(52.50, 57.25, 60.00, 65.00, 74.25, 54.00)
Purchase_quantity_Liters <- c(25, 30, 40, 50, 10, 45)
```

```
# a. Create a data frame for month, price per liter (php) and purchase-quantity (liter). Write the code.
```

```
df <- data.frame(
  Month = Month,
  Price_per_liter_PhP = Price_per_liter_PhP,
  Purchase_quantity_Liters = Purchase_quantity_Liters
)
```

```
df
```

```
##   Month Price_per_liter_PhP Purchase_quantity_Liters
## 1   Jan                52.50                    25
## 2   Feb                57.25                    30
## 3 March                60.00                    40
## 4   Apr                65.00                    50
## 5   May                74.25                    10
## 6   June                54.00                    45
```

```
# b. What is the average fuel expenditure of Mr. Cruz from Jan to June? Note: Use weighted.mean(liter, ,
weighted.mean(Price_per_liter_PhP, Purchase_quantity_Liters)
```

```
## [1] 59.2625
```

```
# 7.
```

```
# a. Type "rivers" in your R console. Create a vector data with 7
# elements, containing the number of elements (length) in rivers,
# their sum (sum), mean (mean), median (median), variance (var)
# standard deviation (sd), minimum (min) and maximum (max).
```

```
data <- c(length(rivers), sum(rivers), mean(rivers), median(rivers), var(rivers),
          sd(rivers), min(rivers), max(rivers))
data
```

```
## [1] 141.0000 83357.0000 591.1844 425.0000 243908.4086 493.8708
## [7] 135.0000 3710.0000
```

```
# b. What are the results?
```

```
# the result are 141, 83357, 591.1844, 425, 243908, 493, 135, and 3710 correspondingly
```

```
# c. Write the code and its outputs.
```

```
#data <- c(length(rivers), sum(rivers), mean(rivers), median(rivers), var(rivers),
#          + sd(rivers), min(rivers), max(rivers))
#> data <- c(length(rivers), sum(rivers), mean(rivers), median(rivers), var(rivers),
#          + sd(rivers), min(rivers), max(rivers))
```

```
#> data
```

```
#[1] 141.0000 83357.0000 591.1844 425.0000 243908.4086 493.8708 135.0000 3710.0000
```

```
# 8.
```

```
# a. Create vectors according to the above table. Write the codes.
```

```
Power_Ranking <- c(1:25)
```

```

Celebrity_Name <- c(
  "Tom Cruise", "Rolling Stones", "Oprah Winfrey", "U2", "Tiger Woods",
  "Steven Spielberg", "Howard Stern", "50 Cent", "Cast of the Sopranos",
  "Dan Brown", "Bruce Springsteen", "Donald Trump", "Muhammad Ali",
  "Paul McCartney", "George Lucas", "Elton John", "David Letterman",
  "Phil Mickelson", "J.K Rowling", "Bradd Pitt", "Peter Jackson",
  "Dr. Phil McGraw", "Jay Lenon", "Celine Dion", "Kobe Bryant"
)

Pay_Millions <- c(
  67, 90, 225, 110, 90, 332, 302, 41, 52, 88, 55, 44, 55,
  40, 233, 34, 40, 47, 75, 25, 39, 45, 32, 40, 31
)

# b. Modify the power ranking and pay of J.K. Rowling. Change power ranking to 15 and pay to 90. Write
Power_Ranking[19] <- 15
Pay_Millions[19] <- 90

# c. Interpret the data.

Modified_Data <- data.frame(
  Ranking = Power_Ranking,
  Celebrity = Celebrity_Name,
  Pay = Pay_Millions)

```

Modified\_Data

##	Ranking	Celebrity	Pay
## 1	1	Tom Cruise	67
## 2	2	Rolling Stones	90
## 3	3	Oprah Winfrey	225
## 4	4	U2	110
## 5	5	Tiger Woods	90
## 6	6	Steven Spielberg	332
## 7	7	Howard Stern	302
## 8	8	50 Cent	41
## 9	9	Cast of the Sopranos	52
## 10	10	Dan Brown	88
## 11	11	Bruce Springsteen	55
## 12	12	Donald Trump	44
## 13	13	Muhammad Ali	55
## 14	14	Paul McCartney	40
## 15	15	George Lucas	233
## 16	16	Elton John	34
## 17	17	David Letterman	40
## 18	18	Phil Mickelson	47
## 19	15	J.K Rowling	90
## 20	20	Bradd Pitt	25
## 21	21	Peter Jackson	39
## 22	22	Dr. Phil McGraw	45
## 23	23	Jay Lenon	32
## 24	24	Celine Dion	40

## 25      25      Kobe Bryant 31