



MACHINE LEARNING









CHAPTER 1: PRELIMINARY

Outlines

- Pattern Recognition
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theory
 - Entropy and Information
-

Example


Handwritten Digit Recognition



Outlines


- Pattern Recognition
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theory
 - Entropy and Information
-

Polynomial Curve Fitting




$$y(x, \mathbf{w}) = w_0 + w_1x + w_2x^2 + \dots + w_Mx^M = \sum_{j=0}^M w_jx^j$$

Sum-of-Squares Error Function




$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$


0th Order Polynomial




1st Order Polynomial




3rd Order Polynomial



9th Order Polynomial



Over-fitting




Root-Mean-Square (RMS) Error: $E_{\text{RMS}} = \sqrt{2E(\mathbf{w}^*)/N}$

Polynomial Coefficients

	$M = 0$	$M = 1$	$M = 3$	$M = 9$
w_0^*	0.19	0.82	0.31	0.35
w_1^*		-1.27	7.99	232.37
w_2^*			-25.43	-5321.83
w_3^*			17.37	48568.31
w_4^*				-231639.30
w_5^*				640042.26
w_6^*				-1061800.52
w_7^*				1042400.18
w_8^*				-557682.99
w_9^*				125201.43


Data Set Size: $N = 15$

9th Order Polynomial



Data Set Size: $N = 100$

9th Order Polynomial




Regularization


Penalize large coefficient values

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$


Regularization: $\ln \lambda = -18$



Regularization: $\ln \lambda = 0$



Regularization: E_{RMS} vs. $\ln \lambda$



Polynomial Coefficients


	$\ln \lambda = -\infty$	$\ln \lambda = -18$	$\ln \lambda = 0$
w_0^*	0.35	0.35	0.13
w_1^*	232.37	4.74	-0.05
w_2^*	-5321.83	-0.77	-0.06
w_3^*	48568.31	-31.97	-0.05
w_4^*	-231639.30	-3.89	-0.03
w_5^*	640042.26	55.28	-0.02
w_6^*	-1061800.52	41.32	-0.01
w_7^*	1042400.18	-45.95	-0.00
w_8^*	-557682.99	-91.53	0.00
w_9^*	125201.43	72.68	0.01

Outlines


- Pattern Recognition
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theories
 - Entropy and Information
-

Probability Theory

Apples and Oranges



Probability Theory



Marginal Probability

$$p(X = x_i) = \frac{c_i}{N}.$$


Joint Probability

$$p(X = x_i, Y = y_j) = \frac{n_{ij}}{N}$$

Conditional Probability

$$p(Y = y_j | X = x_i) = \frac{n_{ij}}{c_i}$$

Probability Theory



Sum Rule

$$\begin{aligned} p(X = x_i) &= \frac{c_i}{N} = \frac{1}{N} \sum_{j=1}^L n_{ij} \\ &= \sum_{j=1}^L p(X = x_i, Y = y_j) \end{aligned}$$

Product Rule

$$\begin{aligned} p(X = x_i, Y = y_j) &= \frac{n_{ij}}{N} = \frac{n_{ij}}{c_i} \cdot \frac{c_i}{N} \\ &= p(Y = y_j | X = x_i) p(X = x_i) \end{aligned}$$

The Rules of Probability

Sum Rule

$$p(X) = \sum_Y p(X, Y)$$

Product Rule

$$p(X, Y) = p(Y|X)p(X)$$


Bayes' Theorem

$$p(Y|X) = \frac{p(X|Y)p(Y)}{p(X)}$$

$$p(X) = \sum_Y p(X|Y)p(Y)$$

posterior \propto likelihood \times prior

Probability Densities




$$p(x \in (a, b)) = \int_a^b p(x) \, dx$$

$$P(z) = \int_{-\infty}^z p(x) \, dx$$

$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) \, dx = 1$$

Transformed Densities




$$\begin{aligned} p_y(y) &= p_x(x) \left| \frac{dx}{dy} \right| \\ &= p_x(g(y)) |g'(y)| \end{aligned}$$

Expectations

$$\mathbb{E}[f] = \sum_x p(x) f(x)$$

$$\mathbb{E}[f] = \int p(x) f(x) \, dx$$

$$\mathbb{E}_x[f|y] = \sum_x p(x|y) f(x)$$


Conditional Expectation
(discrete)

$$\mathbb{E}[f] \simeq \frac{1}{N} \sum_{n=1}^N f(x_n)$$

Approximate Expectation
(discrete and continuous)

Variances and Covariances


$$\text{var}[f] = \mathbb{E} \left[(f(x) - \mathbb{E}[f(x)])^2 \right] = \mathbb{E}[f(x)^2] - \mathbb{E}[f(x)]^2$$

$$\begin{aligned} \text{cov}[x, y] &= \mathbb{E}_{x,y} [\{x - \mathbb{E}[x]\} \{y - \mathbb{E}[y]\}] \\ &= \mathbb{E}_{x,y}[xy] - \mathbb{E}[x]\mathbb{E}[y] \end{aligned}$$

$$\begin{aligned} \text{cov}[\mathbf{x}, \mathbf{y}] &= \mathbb{E}_{\mathbf{x}, \mathbf{y}} [\{\mathbf{x} - \mathbb{E}[\mathbf{x}]\} \{\mathbf{y}^T - \mathbb{E}[\mathbf{y}^T]\}] \\ &= \mathbb{E}_{\mathbf{x}, \mathbf{y}}[\mathbf{x}\mathbf{y}^T] - \mathbb{E}[\mathbf{x}]\mathbb{E}[\mathbf{y}^T] \end{aligned}$$

The Gaussian Distribution

$$\mathcal{N}(x|\mu, \sigma^2) = \frac{1}{(2\pi\sigma^2)^{1/2}} \exp \left\{ -\frac{1}{2\sigma^2} (x - \mu)^2 \right\}$$



$$\mathcal{N}(x|\mu, \sigma^2) > 0$$

$$\int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) dx = 1$$

Gaussian Mean and Variance


$$\mathbb{E}[x] = \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) x \, dx = \mu$$

$$\mathbb{E}[x^2] = \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) x^2 \, dx = \mu^2 + \sigma^2$$

$$\text{var}[x] = \mathbb{E}[x^2] - \mathbb{E}[x]^2 = \sigma^2$$

The Multivariate Gaussian


$$\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\boldsymbol{\Sigma}|^{1/2}} \exp \left\{ -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) \right\}$$



Outlines

- Pattern Recognition
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theories
 - Entropy and Information
-

Gaussian Parameter Estimation



$$p(\mathbf{x} | \mu, \sigma^2) = \prod_{n=1}^N \mathcal{N}(x_n | \mu, \sigma^2)$$

Maximum (Log) Likelihood

$$\ln p(\mathbf{x}|\mu, \sigma^2) = -\frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu)^2 - \frac{N}{2} \ln \sigma^2 - \frac{N}{2} \ln(2\pi)$$


$$\mu_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N x_n \qquad \sigma_{\text{ML}}^2 = \frac{1}{N} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2$$

Properties of μ_{ML} and σ_{ML}^2


$$\mathbb{E}[\mu_{\text{ML}}] = \mu$$

$$\mathbb{E}[\sigma_{\text{ML}}^2] = \left(\frac{N-1}{N}\right) \sigma^2$$

$$\begin{aligned} \tilde{\sigma}^2 &= \frac{N}{N-1} \sigma_{\text{ML}}^2 \\ &= \frac{1}{N-1} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2 \end{aligned}$$



Curve Fitting Re-visited



(t, x) : training data $\Rightarrow \mathbf{w}, \beta$ (\mathbf{w}, β, x_0) : $\Rightarrow p(t|x_0, \mathbf{w}, \beta)$

Maximum Likelihood

$$p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta) = \prod_{n=1}^N \mathcal{N}(t_n | y(x_n, \mathbf{w}), \beta^{-1})$$


$$\ln p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta) = - \underbrace{\frac{\beta}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2}_{\beta E(\mathbf{w})} + \frac{N}{2} \ln \beta - \frac{N}{2} \ln(2\pi)$$

Determine \mathbf{w}_{ML} by minimizing sum-of-squares error, $E(\mathbf{w})$.

$$\frac{1}{\beta_{\text{ML}}} = \frac{1}{N} \sum_{n=1}^N \{y(x_n, \mathbf{w}_{\text{ML}}) - t_n\}^2$$

Predictive Distribution

$$p(t|x, \mathbf{w}_{\text{ML}}, \beta_{\text{ML}}) = \mathcal{N}(t|y(x, \mathbf{w}_{\text{ML}}), \beta_{\text{ML}}^{-1})$$



MAP: A Step towards Bayes

$$p(\mathbf{w}|\alpha) = \mathcal{N}(\mathbf{w}|\mathbf{0}, \alpha^{-1}\mathbf{I}) = \left(\frac{\alpha}{2\pi}\right)^{(M+1)/2} \exp\left\{-\frac{\alpha}{2}\mathbf{w}^T\mathbf{w}\right\}$$

$$p(\mathbf{w}|\mathbf{x}, \mathbf{t}, \alpha, \beta) \propto p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta)p(\mathbf{w}|\alpha)$$

$$\beta\tilde{E}(\mathbf{w}) = \frac{\beta}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\alpha}{2}\mathbf{w}^T\mathbf{w}$$

Determine \mathbf{w}_{MAP} by minimizing regularized sum-of-squares error, $\tilde{E}(\mathbf{w})$.

Bayesian Curve Fitting


$$p(t|x, \mathbf{x}, \mathbf{t}) = \int p(t|x, \mathbf{w})p(\mathbf{w}|\mathbf{x}, \mathbf{t}) d\mathbf{w} = \mathcal{N}(t|m(x), s^2(x))$$

$$m(x) = \beta \phi(x)^T \mathbf{S} \sum_{n=1}^N \phi(x_n) t_n \qquad s^2(x) = \beta^{-1} + \phi(x)^T \mathbf{S} \phi(x)$$

$$\mathbf{S}^{-1} = \alpha \mathbf{I} + \beta \sum_{n=1}^N \phi(x_n) \phi(x_n)^T \qquad \phi(x_n) = (x_n^0, \dots, x_n^M)^T$$


Bayesian Predictive Distribution

$$p(t|x, \mathbf{x}, \mathbf{t}) = \mathcal{N}(t|m(x), s^2(x))$$



Model Selection


Cross-Validation




Outlines

- Pattern Recognition
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theory
 - Entropy and Information
-


Curse of Dimensionality



$D = 1$



$D = 2$




$D = 3$

Curse of Dimensionality

Polynomial curve fitting, $M = 3$

$$y(\mathbf{x}, \mathbf{w}) = w_0 + \sum_{i=1}^D w_i x_i + \sum_{i=1}^D \sum_{j=1}^D w_{ij} x_i x_j + \sum_{i=1}^D \sum_{j=1}^D \sum_{k=1}^D w_{ijk} x_i x_j x_k$$


Gaussian Densities in
higher dimensions



Reduction of Dimensionality

Data Basis Coefficients

$Y = AX$




$$\min_{A_i} A_i^T \text{VAR}(Y) A_i$$

$$A_i^T \text{VAR}(Y) A_i = \lambda_i$$

$$s.t. \quad A_i^T A_i = 1 \quad E[Y_i] = \mathbf{0}$$

Feature Extraction



$$\mathcal{L}(f_1, f_2) = t\|f_1 - f_2\|^2 + (1 - t)[m - \|f_1 - f_2\|^2]_+$$

$t=1$: two vectors belong to the same category; $[]_+$: non-negative

Outlines

- Pattern Recognition
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theory
 - Entropy and Information
-

Decision Theory


Inference step

Determine either $p(t|\mathbf{x})$ or $p(\mathbf{x}, t)$.


Decision step

For given \mathbf{x} , determine optimal t .


Receiver Operating Characteristic Curve




TP	FP
FN	TN
1	1




Bimodal Distribution




Decision Threshold




True Positive




False Positive




True Negative




False Negative





Moving the Threshold: Right




Moving the Threshold: Left



ROC Curve



Minimum Misclassification Rate



$$\begin{aligned} p(\text{mistake}) &= p(\mathbf{x} \in \mathcal{R}_1, \mathcal{C}_2) + p(\mathbf{x} \in \mathcal{R}_2, \mathcal{C}_1) \\ &= \int_{\mathcal{R}_1} p(\mathbf{x}, \mathcal{C}_2) d\mathbf{x} + \int_{\mathcal{R}_2} p(\mathbf{x}, \mathcal{C}_1) d\mathbf{x}. \end{aligned}$$

Minimum Expected Loss

Example: classify medical images as 'cancer' or 'normal'

		Decision	
		cancer	normal
Truth	cancer	0	1000
	normal	1	0

False
Positive

False
Negatives

The diagram illustrates the relationship between the truth and the decision in a classification task. The matrix shows the loss values for different combinations of truth and decision. The 'False Positive' box is connected to the cell where the truth is 'normal' and the decision is 'cancer' (loss 1). The 'False Negatives' box is connected to the cell where the truth is 'cancer' and the decision is 'normal' (loss 1000).


Minimum Expected Loss

$$\mathbb{E}[L] = \sum_k \sum_j \int_{\mathcal{R}_j} L_{kj} p(\mathbf{x}, \mathcal{C}_k) d\mathbf{x}$$

Regions \mathcal{R}_j are chosen to minimize

$$\mathbb{E}[L] = \sum_k L_{kj} p(\mathcal{C}_k | \mathbf{x})$$

Reject Option



Why Separate Inference and Decision?

- Minimizing risk (loss matrix may change over time)
- Reject option
- Unbalanced class priors
- Combining models

Decision Theory for Regression

Inference step

Determine $p(\mathbf{x}, t)$.


Decision step

For given \mathbf{x} , make optimal prediction, $y(\mathbf{x})$, for t .

Loss function: $\mathbb{E}[L] = \iint L(t, y(\mathbf{x}))p(\mathbf{x}, t) \, d\mathbf{x} \, dt$


The Expected Squared Loss Function

$$\mathbb{E}[L] = \iint \{y(\mathbf{x}) - t\}^2 p(\mathbf{x}, t) \, d\mathbf{x} \, dt$$



$$\begin{aligned} \{y(\mathbf{x}) - t\}^2 &= \{y(\mathbf{x}) - \mathbb{E}[t|\mathbf{x}] + \mathbb{E}[t|\mathbf{x}] - t\}^2 \\ &= \{y(\mathbf{x}) - \mathbb{E}[t|\mathbf{x}]\}^2 + 2\{y(\mathbf{x}) - \mathbb{E}[t|\mathbf{x}]\}\{\mathbb{E}[t|\mathbf{x}] - t\} + \{\mathbb{E}[t|\mathbf{x}] - t\}^2 \end{aligned}$$

$$\mathbb{E}[L] = \int \{y(\mathbf{x}) - \mathbb{E}[t|\mathbf{x}]\}^2 p(\mathbf{x}) \, d\mathbf{x} + \int \text{var}[t|\mathbf{x}] p(\mathbf{x}) \, d\mathbf{x}$$



$$y(\mathbf{x}) = \mathbb{E}[t|\mathbf{x}]$$

predictor

noise

$y(x)$: an estimator of the mean of t for given \mathbf{x}

Generative vs Discriminative

Generative approach:



Model $p(t, \mathbf{x}) = p(\mathbf{x}|t)p(t)$

Use Bayes' theorem $p(t|\mathbf{x}) = \frac{p(\mathbf{x}|t)p(t)}{p(\mathbf{x})}$

Discriminative approach:

Model $p(t|\mathbf{x})$ directly

t : category



Outlines

- Pattern Recognition
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theories
 - Entropy and Information
-

Entropy

$$H[x] = - \sum_x p(x) \log_2 p(x)$$

Important quantity in

- coding theory
- statistical physics
- machine learning

Entropy

Coding theory: x discrete with 8 possible states; how many bits to transmit the state of x ?

All states equally likely

$$H[x] = -8 \times \frac{1}{8} \log_2 \frac{1}{8} = 3 \text{ bits.}$$

Entropy

x	a	b	c	d	e	f	g	h
$p(x)$	$\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{16}$	$\frac{1}{64}$	$\frac{1}{64}$	$\frac{1}{64}$	$\frac{1}{64}$
code	0	10	110	1110	111100	111101	111110	111111

$$\begin{aligned} H[x] &= -\frac{1}{2} \log_2 \frac{1}{2} - \frac{1}{4} \log_2 \frac{1}{4} - \frac{1}{8} \log_2 \frac{1}{8} - \frac{1}{16} \log_2 \frac{1}{16} - \frac{4}{64} \log_2 \frac{1}{64} \\ &= 2 \text{ bits} \end{aligned}$$

$$\begin{aligned} \text{average code length} &= \frac{1}{2} \times 1 + \frac{1}{4} \times 2 + \frac{1}{8} \times 3 + \frac{1}{16} \times 4 + 4 \times \frac{1}{64} \times 6 \\ &= 2 \text{ bits} \end{aligned}$$

Entropy



In how many ways can N identical objects be allocated M bins?

$$W = \frac{N!}{\prod_i n_i!}$$

$$H = \frac{1}{N} \ln W \simeq - \lim_{N \rightarrow \infty} \sum_i \left(\frac{n_i}{N} \right) \ln \left(\frac{n_i}{N} \right) = - \sum_i p_i \ln p_i$$

Entropy maximized when $\forall i : p_i = \frac{1}{M}$

Entropy



Differential Entropy

Put bins of width Δ along the real line

$$\lim_{\Delta \rightarrow 0} \left\{ - \sum_i p(x_i) \Delta \ln p(x_i) \right\} = - \int p(x) \ln p(x) dx$$

Differential entropy maximized (for fixed σ^2) when

$$p(x) = \mathcal{N}(x|\mu, \sigma^2)$$

in which case

$$H[x] = \frac{1}{2} \{ 1 + \ln(2\pi\sigma^2) \} .$$

Conditional Entropy

$$H[\mathbf{y}|\mathbf{x}] = - \iint p(\mathbf{y}, \mathbf{x}) \ln p(\mathbf{y}|\mathbf{x}) \, d\mathbf{y} \, d\mathbf{x}$$

$$H[\mathbf{x}, \mathbf{y}] = H[\mathbf{y}|\mathbf{x}] + H[\mathbf{x}]$$

The Kullback-Leibler Divergence

$$\begin{aligned} \text{KL}(p\|q) &= \overset{\text{Cross Entropy } C(p\|q)}{-\int p(\mathbf{x}) \ln q(\mathbf{x}) d\mathbf{x}} - \overset{\text{Entropy } H(p)}{\left(-\int p(\mathbf{x}) \ln p(\mathbf{x}) d\mathbf{x}\right)} \\ &= -\int p(\mathbf{x}) \ln \left\{ \frac{q(\mathbf{x})}{p(\mathbf{x})} \right\} d\mathbf{x} \end{aligned}$$

$$\text{KL}(p\|q) \simeq \frac{1}{N} \sum_{n=1}^N \left\{ \overset{\text{Cross Entropy}}{-\ln q(\mathbf{x}_n|\boldsymbol{\theta})} + \overset{\text{Negative Entropy}}{\ln p(\mathbf{x}_n)} \right\}$$

$$\text{KL}(p\|q) \geq 0 \qquad \text{KL}(p\|q) \neq \text{KL}(q\|p)$$

KL divergence describes a distance between model p and model q

Cross Entropy for Machine Learning

Goal of Machine Learning: $p(\text{real data}) \approx p(\text{model} / \theta)$

we assume: $p(\text{training data}) \approx p(\text{real data})$

Operation of Machine Learning: $p(\text{training data}) \approx p(\text{model} / \theta)$

$$\begin{aligned} & \min_{\theta} \text{KL}(p(\text{training data}) || p(\text{model} | \theta)) \\ \Leftrightarrow & \min_{\theta} C(p(\text{training data}) || p(\text{model} | \theta)) \quad \text{as } H(p(\text{training data})) \text{ is fixed} \end{aligned}$$

Bernoulli model: $p(\text{model} / \theta) = \prod_n \rho^{t_n} (1 - \rho)^{1 - t_n}$ t_n : training data

Cross entropy: $C = -\sum_n t_n \ln \rho + (1 - t_n) \ln(1 - \rho)$ ρ : model parameter

Mutual Information

$$\begin{aligned} I[\mathbf{x}, \mathbf{y}] &\equiv \text{KL}(p(\mathbf{x}, \mathbf{y}) \| p(\mathbf{x})p(\mathbf{y})) \\ &= - \iint p(\mathbf{x}, \mathbf{y}) \ln \left(\frac{p(\mathbf{x})p(\mathbf{y})}{p(\mathbf{x}, \mathbf{y})} \right) d\mathbf{x} d\mathbf{y} \end{aligned}$$

$$I[\mathbf{x}, \mathbf{y}] = H[\mathbf{x}] - H[\mathbf{x}|\mathbf{y}] = H[\mathbf{y}] - H[\mathbf{y}|\mathbf{x}]$$

Mutual information describes the degree of dependence between \mathbf{x} and \mathbf{y}

Information Gain



$H[\mathbf{x}]$: uncertain of balls

$H[\mathbf{x}|\mathbf{y}]$:
uncertain of balls after
weighing once

\mathbf{x} : one ball lighter

\mathbf{y} : weighing once

$\mathbf{x}|\mathbf{y}$: one ball lighter
after weighing once

$$I[\mathbf{x}, \mathbf{y}] = H[\mathbf{x}] - H[\mathbf{x}|\mathbf{y}] = \log_2 3$$

$$H[\mathbf{x}] = \log_2 N$$

After weighing $\frac{N}{3}$ times, all the uncertainties can be removed

Independent Signal Separation




Independent Component Analysis

Mixture Coefficients

Data


Components

$$Y = AX$$


$$\min_A I([X_1, X_2, \dots, X_M] | A, Y)$$

After optimization, the components of X become as much independent as possible

ICA Illustration



Summary

- Pattern Recognition
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theory
 - Entropy and Information
-