

# One-Dimensional Probability Distribution for Random Walker Position

Lucas Brito

October 15, 2021

Our model is as follows: say the probability of finding the walker at position  $x_0$  at a time  $t_0 + \tau$   $P(x_0, t_0 + \tau)$ . There are three possibilities for the walker's position at  $t = t_0$ : the walker was positioned some small distance  $\delta$  to the left (then moved to the right), the walker was positioned some small distance  $\delta$  to the right (then moved to the left), or the walker did not move. With the probability that the walker moves to the right being  $p_r$  and the probability the walker moves to the left  $p_\ell$ , one can set up the equation

$$P(x_0, t_0 + \tau) = p_\ell \cdot P(x_0 + \delta, t_0) + (1 - p_r - p_\ell) \cdot P(x_0, t_0) + p_r \cdot P(x_0 - \delta, t_0)$$

We will use a multivariable Taylor expansion about  $x_0, t_0$ . On the left hand side, we will naturally evaluate the expansion at  $(x_0, t_0 + \tau)$ ; all derivatives with respect to  $x$  evaluate to zero for they are in powers of  $(x_0 - x_0)$ . The left hand side is then

$$\begin{aligned} P(x_0, t_0) + \frac{\partial P}{\partial t} \Big|_{t=t_0} (t_0 - t_0 + \tau) + \frac{1}{4} \frac{\partial^2 P}{\partial t^2} \Big|_{t=t_0} (t_0 - t_0 + \tau)^2 + \dots \\ = P(x_0, t_0) + \frac{\partial P}{\partial t} \Big|_{t=t_0} \tau + \frac{1}{4} \frac{\partial^2 P}{\partial t^2} \Big|_{t=t_0} \tau^2 + \dots \end{aligned}$$

The right hand side, evaluated similarly, is

$$\begin{aligned} p_\ell \left[ P(x_0, t_0) + \frac{\partial P}{\partial x} \Big|_{x=x_0} \delta + \frac{1}{4} \frac{\partial^2 P}{\partial x^2} \Big|_{x_0} \delta^2 + \dots \right] \\ + (1 - p_r - p_\ell) \left[ P(x_0, t_0) + \frac{\partial P}{\partial x} \Big|_{x=x_0} (x_0 - x_0) + \frac{\partial^2 P}{\partial x^2} \Big|_{x=x_0} (x_0 - x_0)^2 + \dots \right] \\ + p_r \left[ P(x_0, t_0) - \frac{\partial P}{\partial x} \Big|_{x=x_0} \delta + \frac{1}{4} \frac{\partial^2 P}{\partial x^2} \Big|_{x_0} \delta^2 + \dots \right] \end{aligned}$$

Every other term of the right hand side evidently disappears and we are left with

$$P(x_0, t_0) + \frac{1}{4} (p_\ell + p_r) \frac{\partial^2 P}{\partial x^2} \Big|_{x=x_0} \delta^2 + \dots$$

With  $\delta$  and  $\tau$  small, we return to the equation and drop terms above the second order.

$$P(x_0, t_0) + \frac{\partial P}{\partial t} \Big|_{t=t_0} \tau + \frac{1}{4} \frac{\partial^2 P}{\partial t^2} \Big|_{t=t_0} \tau^2 = P(x_0, t_0) + \frac{1}{4} (p_\ell + p_r) \frac{\partial^2 P}{\partial x^2} \Big|_{x=x_0} \delta^2$$

With  $x_0$  and  $t_0$  arbitrary, we are free to drop the evaluation notation:

$$\begin{aligned} P(x_0, t_0) + \frac{\partial P}{\partial t} \tau + \frac{1}{4} \frac{\partial^2 P}{\partial t^2} \tau^2 &= P(x_0, t_0) + \frac{1}{4} (p_\ell + p_r) \frac{\partial^2 P}{\partial x^2} \delta^2 \\ \frac{\partial P}{\partial t} \tau + \frac{1}{4} \frac{\partial^2 P}{\partial t^2} \tau^2 &= \frac{1}{4} (p_\ell + p_r) \frac{\partial^2 P}{\partial x^2} \delta^2 \end{aligned}$$

where we've additionally cancelled the constant term. With  $\tau$  and  $\delta$  being arbitrary small quantities, we are free to set  $\tau = \delta^2$  such that

$$\frac{\partial P}{\partial t} \delta^2 + \frac{1}{4} \frac{\partial^2 P}{\partial t^2} \delta^4 = \frac{1}{4} (p_\ell + p_r) \frac{\partial^2 P}{\partial x^2} \delta^2$$

We previously argued that  $\delta^4$  terms are too small to be considered, so we find

$$\frac{\partial P}{\partial t} \delta^2 = \frac{1}{4} (p_\ell + p_r) \frac{\partial^2 P}{\partial x^2} \delta^2$$

and, dividing both sides by  $\delta^2$ ,

$$\boxed{\frac{\partial P}{\partial t} = \frac{1}{4} (p_\ell + p_r) \frac{\partial^2 P}{\partial x^2}}$$

The one-dimensional heat equation!