

# Problem Set 4

Lucas Z. Brito  
PHYS2470

October 25, 2022

## Goldenfeld 7-1

---

Consider the Landau free energy

$$L = \int d^d x \left\{ \frac{1}{2}(\nabla\phi)^2 + \frac{1}{2}r_0\phi^2 + \frac{u_n}{n!}\phi^n \right\}$$

- a) Use the Ginzburg criterion or dimensional analysis to find the upper critical dimension.
- b) Comment on the accuracy of the tricritical exponents which were calculated in exercise 5-2, as a function of dimension.
- c) Show that higher powers of  $\nabla\phi$  and higher derivatives of  $\phi$  are negligible as  $T \rightarrow T_c$ .

## Solution

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Phasellus non libero est. Nunc semper odio lacinia ultrices volutpat. In et dignissim sem. Sed a est dui. Praesent consectetur sodales suscipit. Donec nisl lacus, elementum quis auctor eget, luctus a risus. Morbi est augue, interdum nec sem ut, volutpat pretium dui. Nullam et dui id ipsum egestas fermentum sed ac massa. Mauris ultrices congue diam sed finibus. Mauris congue ante a felis luctus, ut imperdiet purus rhoncus. Nunc rhoncus porttitor erat. Maecenas in nisl justo. Duis dapibus justo magna. Aenean non sapien ultrices, vestibulum massa eget, feugiat odio. Praesent erat dolor, mollis eu neque id, interdum tempus lorem. Praesent blandit aliquet dictum.

$$(i\cancel{\not{D}} - m) \underbrace{\psi(\cancel{x})}_{=1} = 0 \tag{1}$$

See equation (1).